

MVE165/MMG630, Applied Optimization
Lecture 15
Multiobjective optimization and optimization
under uncertainty

Ann-Brith Strömberg


2009-05-04

Applied optimization

- ▶ Many practical optimization problems have several objectives
 - ▶ Some goals cannot be reduced to a common scale of cost/profit \Rightarrow trade-offs must be addressed
 - ▶ **Examples:** financial investments (risk/return), engine design (efficiency/ NO_x /soot), investment cost vs. future emissions (Ass 3a), radio therapy (cure vs. undesired effects, Ass 3b)
- \Rightarrow *Multiple objectives*
- ▶ Decisions may have to be made before information is known
 - ▶ **Examples:** investments (Ass 3a), hydro and wind power production, maintenance planning (Ass 2), energy systems (Ass 1), ...
 - ▶ Represent uncertain data by (discrete) probability distributions
 - ▶ Consider decisions to make after the information is revealed
- \Rightarrow *Optimization under uncertainty (stochastic programming)*

Optimization of multiple objectives

- ▶ Consider the minimization of $f(x) = (x - 1)^2$ subject to $0 \leq x \leq 3$
- ▶ Optimal solution: $x^* = 1$




Optimization of multiple objectives

- ▶ Consider then two objectives:
minimize $[f_1(x), f_2(x)]$ subject to $0 \leq x \leq 3$

$$f_1(x) = (x - 1)^2$$



$$f_2(x) = 3(x - 2)^2$$

- ▶ How can we define an optimal solution?
 - ▶ A solution is **Pareto optimal** if **no other** feasible solution has a better value in **all** objectives
- ⇒ All points $x \in [1, 2]$ are Pareto optimal



Pareto optimal solutions in the objective space

- ▶ minimize $[f_1(x), f_2(x)]$ subject to $0 \leq x \leq 3$
where $f_1(x) = (x - 1)^2$ and $f_2(x) = 3(x - 2)^2$
- ▶ A solution is **Pareto optimal** if **no other** feasible solution has a better value in **all** objectives




- ▶ Pareto optima \Leftrightarrow nondominated points \Leftrightarrow efficient frontier

Efficient points

- ▶ Consider a bi-objective linear program:

$$\begin{array}{ll} \text{maximize} & 3x_1 + x_2 \\ \text{maximize} & -x_1 + 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & 0 \leq x_1 \leq 3 \\ & 0 \leq x_2 \leq 3 \end{array}$$




- ▶ The solutions in the green cone are better w.r.t. both objectives
- ▶ The point $x = (2, 2)$ is an *efficient* solution

Dominated points



$$\begin{array}{ll} \text{maximize} & 3x_1 + x_2 \\ \text{maximize} & -x_1 + 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & 0 \leq x_1 \leq 3 \\ & 0 \leq x_2 \leq 3 \end{array}$$




- ▶ The point $x = (3, 0)$ is *dominated* by the solutions in the green cone
- ▶ Feasible solutions exist that are better w.r.t. both objectives

Dominated points



maximize $3x_1 + x_2$
maximize $-x_1 + 2x_2$
subject to $x_1 + x_2 \leq 4$
 $0 \leq x_1 \leq 3$
 $0 \leq x_2 \leq 3$




- ▶ The point $x = (1, 1)$ is dominated by the solutions in the green cone
- ▶ Feasible solutions exist that are better w.r.t. both objectives

The efficient frontier—the set of Pareto optimal solutions



$$\begin{array}{ll} \text{maximize} & 3x_1 + x_2 \\ \text{maximize} & -x_1 + 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & 0 \leq x_1 \leq 3 \\ & 0 \leq x_2 \leq 3 \end{array}$$




- ▶ The set of efficient solutions is given by

$$\left\{ \mathbf{x} \in \mathbb{R}^2 \mid \mathbf{x} = \alpha \begin{pmatrix} 3 \\ 1 \end{pmatrix} + (1 - \alpha) \begin{pmatrix} 1 \\ 3 \end{pmatrix}, 0 \leq \alpha \leq 1 \right\} \cup \left\{ \mathbf{x} \in \mathbb{R}^2 \mid \mathbf{x} = \alpha \begin{pmatrix} 1 \\ 3 \end{pmatrix} + (1 - \alpha) \begin{pmatrix} 0 \\ 3 \end{pmatrix}, 0 \leq \alpha \leq 1 \right\}$$

The Pareto optimal set in the objective space



$$\begin{aligned} &\text{maximize} && f_1(\mathbf{x}) := 3x_1 + x_2 \\ &\text{maximize} && f_2(\mathbf{x}) := -x_1 + 2x_2 \\ &\text{subject to} && x_1 + x_2 \leq 4 \\ &&& 0 \leq x_1 \leq 3 \\ &&& 0 \leq x_2 \leq 3 \end{aligned}$$





- ▶ The set of Pareto optimal objective values is given by

$$\left\{ (f_1, f_2) \in \mathbb{R}^2 \mid \mathbf{f} = \alpha \begin{pmatrix} 10 \\ -1 \end{pmatrix} + (1 - \alpha) \begin{pmatrix} 6 \\ 5 \end{pmatrix}, 0 \leq \alpha \leq 1 \right\} \cup \left\{ (f_1, f_2) \in \mathbb{R}^2 \mid \mathbf{f} = \alpha \begin{pmatrix} 6 \\ 5 \end{pmatrix} + (1 - \alpha) \begin{pmatrix} 3 \\ 6 \end{pmatrix}, 0 \leq \alpha \leq 1 \right\}$$

Mapping from the decision space to the objective space

maximize $[3x_1 + x_2; -x_1 + 2x_2]$
subject to $x_1 + x_2 \leq 4, \quad 0 \leq x_1 \leq 3, \quad 0 \leq x_2 \leq 3$



Solutions methods for multiobjective optimization

- ▶ Construct the efficient frontier by treating one objective as a constraint and optimizing for the other:

$$\begin{array}{ll} \text{maximize} & 3x_1 + x_2 \\ \text{subject to} & -x_1 + 2x_2 \geq \varepsilon \\ & x_1 + x_2 \leq 4 \\ & 0 \leq x_1 \leq 3 \\ & 0 \leq x_2 \leq 3 \end{array}$$

- ▶ Here, let $\varepsilon \in [-1, 6]$. Why?
- ▶ What if the number of objectives is > 2 ?
- ▶ How many programs do we have to solve for seven objectives and ten values of ε_k for each objective f_k ?

Solution methods: preemptive optimization


- ▶ Consider one objective at a time—the most important first
- ▶ Solve for the first objective
- ▶ Solve for the second objective over the solution set for the first
- ▶ Solve for the third objective over the solution set for the second
- ▶ ...

- ▶ The solution is an efficient point
- ▶ Different orderings of the objectives yield different solutions

- ▶ Exercise: solve the previous example using preemptive optimization on different orderings

Solution methods: weighted sums of objectives

- ▶ Give each maximization (minimization) objective a positive (negative) weight
- ▶ Solve a single objective maximization problem
- ⇒ Yields an efficient solution
- ▶ Well spread weights do not necessarily produce solutions that are well spread on the efficient frontier (ex: $\{\frac{1}{10}, \frac{1}{2}, 1, 2, 10\}$)
- ▶ If the objectives are not concave (maximization) or the feasible set is not convex, then not all points on the efficient frontier may be possible to detect using weighted sums of objectives



Solution methods: soft constraints

- ▶ Consider the multiobjective optimization problem to

$$\text{maximize } [f_1(\mathbf{x}), \dots, f_K(\mathbf{x})] \text{ subject to } \mathbf{x} \in X$$

- ▶ Define a target value t_k and a deficiency variable $d_k \geq 0$ for each objective f_k
- ▶ Construct a soft constraint for each objective:

$$\text{maximize } f_k(\mathbf{x}) \quad \Rightarrow \quad f_k(\mathbf{x}) + d_k \geq t_k, \quad k = 1, \dots, K$$

- ▶ Minimize the sum of deficiencies:

$$\begin{aligned} & \text{minimize} && \sum_{k \in K} d_k \\ & \text{subject to} && f_k(\mathbf{x}) + d_k \geq t_k, \quad k = 1, \dots, K \\ & && d_k \geq 0, \quad k = 1, \dots, K \\ & && \mathbf{x} \in X \end{aligned}$$


- ▶ Important: Find first a common scale for f_k , $k = 1, \dots, K$

Deterministic optimization models

- ▶ In a *deterministic* optimization model, *uncertain* parameters are represented by, e.g., (empirical) averages
- ▶ The weakness is that the *prediction is considered as a truth* and decisions are made as if the future was completely known
- ▶ Optimization tends to *augment errors in the data* when uncertain data is replaced by predictions — “*Do something as good as possible*”
- ▶ We need a methodology for handling optimization under uncertainty


Scenario approaches vs. optimization under uncertainty

- ▶ **Common approach:** Handle deficiency of deterministic optimization models by identifying *scenarios* representing the uncertainty—solve one deterministic model for each scenario
- ⇒ May yield some information about the variations of the solution,
- ▶ But each decision proposal presumes *perfect information* about the future
- ▶ **Optimization under uncertainty** (*stochastic programming*)
- ▶ Uncertain parameters represented by *stochastic variables*
- ▶ Discretization necessary for the solvability of the models ⇒ discrete (approximate) probability distributions




A decision must be made prior all data being known

- ▶ Several time stages \Rightarrow a scenario *tree*



Two realizations \leftrightarrow two branches

A scenario tree
with three
realizations
each time step:



- ▶ Stochastic variables are realized *between* decision time points.
- ▶ Goal: optimize the *expected value over the scenario tree* of, e.g., the revenue

Deterministic optimization vs. optimization under uncertainty

Deterministic optimization


- ▶ All parameters and conditions are assumed known for sure

Optimization under uncertainty


- ▶ Decisions are based on observations (previous outcomes) and under *uncertainty of future outcomes*
- ▶ A decision *must not depend* on outcomes not yet revealed
- ▶ A node in the tree \iff a vector of decision variables
- ▶ In a specific node, the remaining future uncertainty is represented by the sub-tree rooted in that node
- ▶ Typically, only the decision associated to the root node is implemented—next time stage a new model is solved—a so called *rolling horizon*
- ▶ The magnitude of an optimization problem *increases* when uncertainties are modeled explicitly

The LEGO furniture factory revisited

Small block, purchase price: 100



Large block, purchase price: 200



Chair: $\left[\begin{array}{l} \text{demand: } 50 + \xi_2 \\ \text{sales price: } (5 + \eta_2) \cdot 100 \\ \text{prod. cost: } 50 \end{array} \right]$









Table: $\left[\begin{array}{l} \text{demand: } 100 + \xi_1 \\ \text{sales price: } (8 + \eta_1) \cdot 100 \\ \text{prod. cost: } 100 \end{array} \right]$



The *stochastic parameters* η_1 , η_2 , ξ_1 , and ξ_2 are assumed to have discrete probability functions

Discrete probability functions

- ▶ The values $\eta_1 \in \{-1, 0, 0.5\}$, $\eta_2 \in \{-0.5, 0.5, 1\}$, $\xi_1 \in \{-20, 0, 20\}$, $\xi_2 \in \{-10, 0, 5\}$ are achieved with probabilities $\{\frac{1}{4}, \frac{1}{2}, \frac{1}{4}\}$



- ▶ **Assumption here:** Low/medium/high demand levels correspond to low/medium/high price levels
- ⇒ The four stochastic parameters are *dependent*
- ⇒ Totally three scenarios

Definition of variables

- ▶ The demand and the selling prices are *not known* when the blocks are purchased
- ▶ The purchase budget is 80000
- ▶ Variables:
 - x_1 = # of large blocks purchased
 - x_2 = # of small blocks purchased
 - y_1 = # tables produced
 - y_2 = # chairs produced
 - v_1 = # tables sold
 - v_2 = # chairs sold
- ▶ The values of the purchase variables x_1 and x_2 must be *decided on before* the demand and selling prices are known
- ▶ Production (y_1 and y_2) and sales (v_1 and v_2) are decided on later

Mathematical model

Minimize *purchase cost plus production cost minus sales revenue*

$$\begin{aligned} \text{minimize}_{x,y,v} \quad & z := 100 \cdot [2x_1 + x_2 + y_1 - (8 + \eta_1)v_1 + 0.5y_2 - (5 + \eta_2)v_2] \\ \text{subject to} \quad & 2x_1 + x_2 \leq 800 && \text{(budget)} \\ & x_1 - 2y_1 - y_2 \geq 0 && \text{(large blocks used } \leq \text{ purchased)} \\ & x_2 - 2y_1 - 2y_2 \geq 0 && \text{(small blocks used } \leq \text{ purchased)} \\ & y_1 - v_1 \geq 0 && \text{(tables sold } \leq \text{ produced)} \\ & y_2 - v_2 \geq 0 && \text{(chairs sold } \leq \text{ produced)} \\ & v_1 \leq 100 + \xi_1 && \text{(tables sold } \leq \text{ demand)} \\ & v_2 \leq 50 + \xi_2 && \text{(chairs sold } \leq \text{ demand)} \\ & x_1, x_2, y_1, v_1, y_2, v_2 \geq 0 && \text{(integer)} \end{aligned}$$

Deterministic (expected value) solution

- ▶ Assume that the stochastic parameters attain their respective expected values:
 - ▶ $E(\eta_1) = \frac{1}{4} \cdot (-1) + \frac{1}{2} \cdot 0 + \frac{1}{4} \cdot 0.5 = -0.125$
 - ▶ $E(\eta_2) = \frac{1}{4} \cdot (-0.5) + \frac{1}{2} \cdot 0.5 + \frac{1}{4} \cdot 1 = 0.375$
 - ▶ $E(\xi_1) = \frac{1}{4} \cdot (-20) + \frac{1}{2} \cdot 0 + \frac{1}{4} \cdot 20 = 0$
 - ▶ $E(\xi_2) = \frac{1}{4} \cdot (-10) + \frac{1}{2} \cdot 0 + \frac{1}{4} \cdot 5 = -1.25$
- ▶ Replace the stochastic parameters in the mathematical model by their expected values.

The expected value solution

$$\begin{array}{llllll} \text{minimize}_{x,y,z} & z := 100 \cdot [2x_1 + x_2 + y_1 - 7.875v_1 + 0.5y_2 - 5.375v_2] & & & & \\ \text{subject to} & 2x_1 + x_2 & & & & \leq 800 \\ & x_1 & -2y_1 & -y_2 & & \geq 0 \\ & & x_2 & -2y_1 & -2y_2 & \geq 0 \\ & & & y_1 - v_1 & & \geq 0 \\ & & & & y_2 - v_2 & \geq 0 \\ & & & & v_1 & \leq 100 \\ & & & & & v_2 \leq 48.75 \\ & x_1, & x_2, & y_1, v_1, & y_2, v_2 & \geq 0 \quad (\text{integer}) \end{array}$$

- ▶ Solution: $x_1 = 248.75$, $x_2 = 297.5$, $y_1 = v_1 = 100$,
 $y_2 = v_2 = 48.75$, $z := -13016$ (minus the profit)
- ▶ **Deterministic solution:**
 - ▶ Purchase ≈ 249 large and ≈ 298 small blocks
 - ▶ Produce and sell ≈ 100 tables and ≈ 49 chairs
 - ▶ Profit: 13016

OBS: Infeasible at the lowest demand scenario (80 tables, 40 chairs)!

A hedging deterministic optimization model

- ▶ Choose $\eta_1 = -1$, $\eta_2 = -0.5$, $\xi_1 = -20$, and $\xi_2 = -10$

$$\text{minimize}_{x,y,z} \quad z := 100 \cdot [2x_1 + x_2 + y_1 - 7v_1 + 0.5y_2 - 4.5v_2]$$

$$\begin{aligned} \text{subject to} \quad & 2x_1 + x_2 && \leq 800 \\ & x_1 & -2y_1 & -y_2 \geq 0 \\ & & x_2 & -2y_1 & -2y_2 \geq 0 \\ & & & y_1 - v_1 & \geq 0 \\ & & & & y_2 - v_2 \geq 0 \\ & & & & v_1 & \leq 80 \\ & & & & & v_2 \leq 40 \\ & x_1, & x_2, & y_1, v_1, & y_2, v_2 & \geq 0 \quad (\text{integer}) \end{aligned}$$

- ▶ Solution: $x_1 = 152.1$, $x_2 = 181.8$, $y_1 = v_1 = 61.22$,
 $y_2 = v_2 = 29.67$, $z = 0$
- ▶ **Deterministic solution:**
 - ▶ Purchase ≈ 152 large and ≈ 182 small blocks
 - ▶ Produce and sell ≈ 61 tables and ≈ 30 chairs
 - ▶ Profit: 0

A stochastic optimization model

Stage 1: The purchase decision takes the possible outcomes of demand and selling prices into consideration, with their respective probabilities, and the corresponding decisions on production/sales to make later on

- ▶ Three different *scenarios*: low/medium/high level of prices and demand: $\eta^1 = [-1, -0.5]$, $\xi^1 = [-20, -10]$, $\eta^2 = [0, 0.5]$, $\xi^2 = [0, 0]$, $\eta^3 = [0.5, 1]$, $\xi^3 = [20, 5]$

Stage 2: When the decisions on production/sales are to be made, the levels of prices and demand will be revealed

- ▶ Also the decided purchase of raw material is known
- ⇒ Optimize with respect to the outcome of the stochastic parameters and the decisions from stage 1 (*recourse*)

The first stage decision

- ▶ Minimize the purchase cost minus the *expected future profit*
- ▶ Decide on how many blocks to purchase (x)
- ▶ Consider the possible future outcomes of the demand (ξ) and price (η) levels and the decisions on the production ($y(x, \xi, \eta)$) and sales ($v(x, \xi, \eta)$)

$$\begin{array}{ll} \text{minimize}_x & z := 100 \cdot [2x_1 + x_2 - E_{\xi, \eta} Q(x, \xi, \eta)] \quad (\text{convex in } x) \\ \text{subject to} & 2x_1 + x_2 \leq 800 \quad (\text{purchase} \leq \text{budget}) \\ & x_1, x_2 \geq 0 \quad (\text{integer}) \end{array}$$

- ▶ $E_{\xi, \eta} Q(x, \xi, \eta)$ denotes the *expected value of the future profit*, which is computed in stage 2

The second stage decisions

- ▶ Maximize future profit (sales revenues minus production costs)
- ▶ Decide on production and sales for each outcome of the price (η) and demand (ξ) and for each purchase decision (x)

$$Q(x, \xi, \eta) = \left(\begin{array}{l} \text{maximize}_{y,v} \quad -y_1 + (8 + \eta_1)v_1 - 0.5y_2 + (5 + \eta_2)v_2 \\ \text{subject to} \quad 2y_1 \quad +y_2 \leq x_1 \\ \quad \quad \quad 2y_1 \quad +2y_2 \leq x_2 \\ \quad \quad \quad v_1 \quad \quad \quad \leq y_1 \\ \quad \quad \quad \quad \quad v_2 \leq y_2 \\ \quad \quad \quad v_1 \quad \quad \quad \leq 100 + \xi_1 \\ \quad \quad \quad \quad \quad v_2 \leq 50 + \xi_2 \\ y_1, v_1 \quad y_2, v_2 \geq 0 \quad \quad \quad \text{(integer)} \end{array} \right)$$

Expected future profits—the second stage decisions

$$E_{\xi, \eta} Q(x, \xi, \eta) = \frac{1}{4} Q(x, \xi^1, \eta^1) + \frac{1}{2} Q(x, \xi^2, \eta^2) + \frac{1}{4} Q(x, \xi^3, \eta^3)$$

$$= \frac{1}{4} \begin{pmatrix} \max -y_1^1 + 7v_1^1 - 0.5y_2^1 + 4.5v_2^1 \\ \text{subject to} & 2y_1^1 + y_2^1 \leq x_1 \\ & 2y_1^1 + 2y_2^1 \leq x_2 \\ & v_1^1 \leq \min\{y_1^1, 80\} \\ & v_2^1 \leq \min\{y_2^1, 40\} \\ & v_1^1, v_2^1, y_1^1, y_2^1 \geq 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} \max -y_1^2 + 8v_1^2 - 0.5y_2^2 + 5.5v_2^2 \\ \text{subject to} & 2y_1^2 + y_2^2 \leq x_1 \\ & 2y_1^2 + 2y_2^2 \leq x_2 \\ & v_1^2 \leq \min\{y_1^2, 100\} \\ & v_2^2 \leq \min\{y_2^2, 50\} \\ & v_1^2, v_2^2, y_1^2, y_2^2 \geq 0 \end{pmatrix} + \frac{1}{4} \begin{pmatrix} \max -y_1^3 + 8.5v_1^3 - 0.5y_2^3 + 6v_2^3 \\ \text{subject to} & 2y_1^3 + y_2^3 \leq x_1 \\ & 2y_1^3 + 2y_2^3 \leq x_2 \\ & v_1^3 \leq \min\{y_1^3, 120\} \\ & v_2^3 \leq \min\{y_2^3, 55\} \\ & v_1^3, v_2^3, y_1^3, y_2^3 \geq 0 \end{pmatrix}$$

Solution to the optimization model that takes the uncertainty into consideration

- ▶ First stage solution: $x = (200, 250)$
- ⇒ Objective value (minus expected profit): $z = -10687$
- ▶ Second stage solution: $y^1 = v^1 = (80, 40)$,
 $y^2 = v^2 = y^3 = v^3 = (75, 50)$
 - ▶ If scenario 1 occurs (low) the profit becomes:
 $-100 \cdot (2 \cdot 200 + 250 + 80 - 7 \cdot 80 + 0.5 \cdot 40 - 4.5 \cdot 40) = -1000$
 - ▶ If scenario 2 occurs (medium) the profit becomes:
 $-100 \cdot (2 \cdot 200 + 250 + 75 - 8 \cdot 75 + 0.5 \cdot 50 - 5.5 \cdot 50) = 12500$
 - ▶ If scenario 3 occurs (high) the profit becomes:
 $-100 \cdot (2 \cdot 200 + 250 + 75 - 8.5 \cdot 75 + 0.5 \cdot 50 - 6 \cdot 50) = 18750$
 - ▶ Expected profit: $\frac{-1000}{4} + \frac{12500}{2} + \frac{18750}{4} = 10687$

What if we would solve the deterministic model (expected value solution) for the first stage decision (x) and adjust the second stage solution ((y, v)) to the actual scenario observed in the second stage?

1. Solve the deterministic model $\Rightarrow \bar{x}_1 = 248.75, \bar{x}_2 = 297.5$
2. Compute the future profit $Q(\bar{x}, \xi, \eta)$ for each scenario
3. The expected value of the expected value (deterministic) solution is: $z = 100 \cdot [2\bar{x}_1 + \bar{x}_2 - E_{\xi, \eta} Q(\bar{x}, \xi, \eta)]$ (next page)
4. Second stage solutions (three different scenarios):
 $y^1 = v^1 = (80, 40)$ $y^2 = v^2 = (100, 48.75)$
 $y^3 = v^3 = (100, 48.75)$
 \Rightarrow The expected profit of the expected value solution: 9140
5. The value of the stochastic solution: $10687 - 9140 = 1547 > 0$

The expected profit increases when taking the uncertainties under consideration already in the formulation of the model

Expected future profit for $\bar{x} = (248.75, 297.5)$

$$E_{\xi, \eta} Q(\bar{x}, \xi, \eta) = \frac{1}{4} Q(\bar{x}, \xi^1, \eta^1) + \frac{1}{2} Q(\bar{x}, \xi^2, \eta^2) + \frac{1}{4} Q(\bar{x}, \xi^3, \eta^3)$$

$$= \frac{1}{4} \begin{pmatrix} \max -y_1^1 + 7v_1^1 - 0.5y_2^1 + 4.5v_2^1 \\ \text{subject to } 2y_1^1 + y_2^1 \leq 248.75 \\ 2y_1^1 + 2y_2^1 \leq 297.5 \\ v_1^1 \leq \min\{y_1^1, 80\} \\ v_2^1 \leq \min\{y_2^1, 40\} \\ v_1^1, v_2^1, y_1^1, y_2^1 \geq 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} \max -y_1^2 + 8v_1^2 - 0.5y_2^2 + 5.5v_2^2 \\ \text{subject to } 2y_1^2 + y_2^2 \leq 248.75 \\ 2y_1^2 + 2y_2^2 \leq 297.5 \\ v_1^2 \leq \min\{y_1^2, 100\} \\ v_2^2 \leq \min\{y_2^2, 50\} \\ v_1^2, v_2^2, y_1^2, y_2^2 \geq 0 \end{pmatrix} + \frac{1}{4} \begin{pmatrix} \max -y_1^3 + 8.5v_1^3 - 0.5y_2^3 + 6v_2^3 \\ \text{subject to } 2y_1^3 + y_2^3 \leq 248.75 \\ 2y_1^3 + 2y_2^3 \leq 297.5 \\ v_1^3 \leq \min\{y_1^3, 120\} \\ v_2^3 \leq \min\{y_2^3, 55\} \\ v_1^3, v_2^3, y_1^3, y_2^3 \geq 0 \end{pmatrix}$$