

MVE165/MMG630, Applied Optimization

Lecture 2

Ann-Brith Strömberg

2009-03-17


Intersections of linear (in)equalities are convex sets

- ▶ A set S is convex if, for any elements $\mathbf{x}, \mathbf{y} \in S$ it holds that


$$\alpha \mathbf{x} + (1 - \alpha) \mathbf{y} \in S \text{ for all } 0 \leq \alpha \leq 1$$

- ▶ Examples:

Convex sets



Non-convex sets




Linear functions are convex


- ▶ A function f is convex on the set S if, for any elements $\mathbf{x}, \mathbf{y} \in S$ it holds that

$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}) \text{ for all } 0 \leq \alpha \leq 1$$

Convex function



Non-convex function



Global solutions of convex programs

- ▶ Let \mathbf{x}^* be a *local* minimizer of a *convex function* over a *convex set*. Then \mathbf{x}^* is also a *global* minimizer.
 - ▶ Linear functions are convex and polyhedra are convex sets
- ⇒ Every local optimum of a linear program is a global optimum
- ▶ If a linear program has any optimal solutions, at least one optimal solution is at an extreme point of the feasible set

A general linear program

$$\text{min or max } c_1x_1 + \dots + c_nx_n$$

$$\text{subject to } a_{i1}x_1 + \dots + a_{in}x_n \begin{cases} \leq \\ = \\ \geq \end{cases} b_i, \quad i = 1, \dots, m$$

$$x_j \geq 0, \quad j = 1, \dots, n$$



The standard form and the simplex method for linear programs

- ▶ Every linear program can be reformulated such that:
 - ▶ all constraints are expressed as equalities with non-negative right hand sides and
 - ▶ all variables are non-negative
- ▶ Referred to as the *standard form*
- ▶ These requirements streamline the simplex method calculations
- ▶ Commercial solvers can handle also inequality constraints and “free” variables—the reformulations are automatically taken care of



The simplex method—reformulations

- ▶ The lego example:

$$\begin{bmatrix} 2x_1 & +x_2 & \leq & 6 \\ 2x_1 & +2x_2 & \leq & 8 \\ & x_1, x_2 & \geq & 0 \end{bmatrix} \Leftrightarrow \begin{bmatrix} 2x_1 & +x_2 & +s_1 & = & 6 \\ 2x_1 & +2x_2 & & +s_2 & = & 8 \\ & x_1, x_2, s_1, s_2 & \geq & 0 \end{bmatrix}$$

- ▶ s_1 and s_2 are called *slack variables*—they “fill out” the (positive) distances between the left and right hand sides
- ▶ *Surplus variable* s_3 :

$$\begin{bmatrix} x_1 & + & x_2 & \geq & 800 \\ & x_1, x_2 & \geq & 0 \end{bmatrix} \Leftrightarrow \begin{bmatrix} x_1 & + & x_2 & - & s_3 & = & 800 \\ & x_1, x_2, s_3 & \geq & 0 \end{bmatrix}$$



The simplex method—reformulations, cont.

- ▶ Non-negative right hand side:

$$\begin{bmatrix} x_1 - x_2 & \leq & -23 \\ & x_1, x_2 & \geq & 0 \end{bmatrix} \Leftrightarrow \begin{bmatrix} -x_1 + x_2 & \geq & 23 \\ & x_1, x_2 & \geq & 0 \end{bmatrix} \Leftrightarrow \begin{bmatrix} -x_1 + x_2 - s_4 & = & 23 \\ & x_1, x_2, s_4 & \geq & 0 \end{bmatrix}$$


- ▶ Free variables:

$$\begin{bmatrix} x_1 + x_2 & \leq & 10 \\ & x_1 & \geq & 0 \end{bmatrix} \Leftrightarrow \begin{bmatrix} x_1 + x_2^1 - x_2^2 & \leq & 10 \\ & x_1, x_2^1, x_2^2 & \geq & 0 \end{bmatrix} \Leftrightarrow \begin{bmatrix} x_1 + x_2^1 - x_2^2 + s_5 & = & 10 \\ & x_1, x_2^1, x_2^2, s_5 & \geq & 0 \end{bmatrix}$$



Basic feasible solutions


- ▶ Consider m equations of n variables, where $m \leq n$
- ▶ Set $n - m$ variables to zero and solve (if possible) the remaining $(m \times m)$ system of equations
- ▶ If the solution is *unique*, it is called a *basic solution*
- ▶ Such a solution corresponds to an intersection (feasible or infeasible) of m hyperplanes in \mathbb{R}^m
- ▶ Each extreme point of the feasible set is an intersection of m hyperplanes with all variable values ≥ 0
- ▶ Basic feasible solution \Leftrightarrow extreme point of the feasible set



Lecture 2 Applied Optimization

Basic and non-basic variables

basic variables	basic solution			non-basic variables (0, 0)	point	feasible?
s_1, s_2, s_3	23	6	85	x_1, x_2	A	yes
s_1, s_2, x_1	$-5\frac{1}{3}$	$4\frac{1}{9}$	$28\frac{1}{3}$	s_3, x_2	H	no
s_1, s_2, x_2	23	$-4\frac{5}{8}$	$10\frac{5}{8}$	x_1, s_3	C	no
s_1, x_1, s_3	-67	90	-185	s_2, x_2	I	no
s_1, x_2, s_3	23	6	37	s_2, x_1	B	yes
x_1, s_2, s_3	23	$4\frac{7}{15}$	16	s_1, x_2	G	yes
x_2, s_2, s_3	-	-	-	s_1, x_1	-	-
x_1, x_2, s_1	15	5	8	s_2, s_3	D	yes
x_1, x_2, s_2	23	2	$2\frac{7}{15}$	s_1, s_3	F	yes
x_1, x_2, s_3	23	$4\frac{7}{15}$	$-19\frac{11}{15}$	s_1, s_2	E	no



Lecture 2 Applied Optimization

Basic feasible solutions, example

- ▶ Constraints:


$$\begin{aligned} x_1 &\leq 23 & (1) \\ 0.067x_1 + x_2 &\leq 6 & (2) \\ 3x_1 + 8x_2 &\leq 85 & (3) \\ x_1, x_2 &\geq 0 \end{aligned}$$

- ▶ Add slack variables:

$$\begin{aligned} x_1 + s_1 &= 23 & (1) \\ 0.067x_1 + x_2 + s_2 &= 6 & (2) \\ 3x_1 + 8x_2 + s_3 &= 85 & (3) \\ x_1, x_2, s_1, s_2, s_3 &\geq 0 \end{aligned}$$


$m = 3$

$n = 5$



Lecture 2 Applied Optimization

Basic feasible solutions correspond to solutions to the system of equations that fulfil non-negativity



$$\begin{bmatrix} x_1 & +s_1 & = 23 \\ 0.067x_1 & +x_2 & +s_2 & = 6 \\ 3x_1 & +8x_2 & +s_3 & = 85 \end{bmatrix}$$

$$A: x_1 = x_2 = 0 \Rightarrow \begin{bmatrix} s_1 & = 23 \\ s_2 & = 6 \\ s_3 & = 85 \end{bmatrix}$$

$$B: x_1 = s_2 = 0 \Rightarrow \begin{bmatrix} s_1 & = 23 \\ x_2 & = 6 \\ 8x_2 & +s_3 & = 85 \end{bmatrix}$$


$$D: s_3 = s_2 = 0 \Rightarrow \begin{bmatrix} x_1 & +s_1 & = 23 \\ 0.067x_1 & +x_2 & = 6 \\ 3x_1 & +8x_2 & = 85 \end{bmatrix}$$

$$F: s_3 = s_1 = 0 \Rightarrow \begin{bmatrix} x_1 & +s_1 & = 23 \\ 0.067x_1 & +x_2 & +s_2 & = 6 \\ 3x_1 & +8x_2 & = 85 \end{bmatrix}$$

$$G: x_2 = s_1 = 0 \Rightarrow \begin{bmatrix} x_1 & = 23 \\ 0.067x_1 & +s_2 & = 6 \\ 3x_1 & +s_3 & = 85 \end{bmatrix}$$

Lecture 2 Applied Optimization

Basic **infeasible** solutions correspond to solutions to the system of equations



$$H: x_2 = s_3 = 0 \Rightarrow \begin{cases} x_1 + s_1 = 23 \\ 0.067x_1 + s_2 = 6 \\ 3x_1 = 85 \end{cases}$$

$$C: x_1 = s_3 = 0 \Rightarrow \begin{cases} s_1 = 23 \\ x_2 + s_2 = 6 \\ 8x_2 = 85 \end{cases}$$

$$I: s_2 = x_2 = 0 \Rightarrow \begin{cases} x_1 + s_1 = 23 \\ 0.067x_1 + s_3 = 6 \\ 3x_1 = 85 \end{cases}$$

$$\therefore s_1 = x_1 = 0 \Rightarrow \begin{cases} s_1 = 23 \\ x_2 + s_2 = 6 \\ 8x_2 + s_3 = 85 \end{cases}$$

$$E: s_1 = s_2 = 0 \Rightarrow \begin{cases} x_1 + s_1 = 23 \\ 0.067x_1 + x_2 = 6 \\ 3x_1 + 8x_2 + s_3 = 85 \end{cases}$$

Basic feasible solutions and the simplex method

- Express the m basic variables in terms of the $n - m$ non-basic variables
- Example: Start at $x_1 = x_2 = 0 \Rightarrow s_1, s_2, s_3$ are basic

$$\begin{cases} x_1 + s_1 = 23 \\ \frac{1}{15}x_1 + x_2 + s_2 = 6 \\ 3x_1 + 8x_2 + s_3 = 85 \end{cases}$$

- Express $s_1, s_2,$ and s_3 in terms of x_1 and x_2 :

$$\begin{cases} s_1 = 23 - x_1 \\ s_2 = 6 - \frac{1}{15}x_1 - x_2 \\ s_3 = 85 - 3x_1 - 8x_2 \end{cases}$$

- Express the objective in terms of the non-basic variables:
 $z = 2x_1 + 3x_2 \Leftrightarrow z - 2x_1 - 3x_2 = 0$

Basic feasible solutions and the simplex method

- The first basic solution can be represented as

$$\begin{array}{rcl} -z + 2x_1 + 3x_2 & = & 0 \quad (0) \\ x_1 + s_1 & = & 23 \quad (1) \\ \frac{1}{15}x_1 + x_2 + s_2 & = & 6 \quad (2) \\ 3x_1 + 8x_2 + s_3 & = & 85 \quad (3) \end{array}$$

- Marginal values for increasing the non-basic variables x_1 and x_2 from zero: 2 and 3, resp.

\Rightarrow Choose x_2 — let x_2 enter the basis DRAW GRAPH!!

- One basic variable ($s_1, s_2,$ or s_3) must leave the basis. Which?
- The value of x_2 can increase until some basic variable reaches the value 0:

$$\left. \begin{array}{l} (2): s_2 = 6 - x_2 \geq 0 \Rightarrow x_2 \leq 6 \\ (3): s_3 = 85 - 8x_2 \geq 0 \Rightarrow x_2 \leq 10\frac{5}{8} \end{array} \right\} \Rightarrow \begin{array}{l} s_2 = 0 \text{ when} \\ x_2 = 6 \\ \text{(and } s_3 = 37) \end{array}$$

- s_2 will leave the basis

Change basis through row operations

- Eliminate s_2 from the basis, let x_2 enter the basis using row operations:

$$\begin{array}{rcl} -z + 2x_1 + 3x_2 & = & 0 \quad (0) \\ x_1 + s_1 & = & 23 \quad (1) \\ \frac{1}{15}x_1 + x_2 + s_2 & = & 6 \quad (2) \\ 3x_1 + 8x_2 + s_3 & = & 85 \quad (3) \end{array}$$

$$\begin{array}{rcl} -z + \frac{9}{5}x_1 - 3s_2 & = & -18 \quad (0) - 3 \cdot (2) \\ x_1 + s_1 & = & 23 \quad (1) - 0 \cdot (2) \\ \frac{1}{15}x_1 + x_2 + s_2 & = & 6 \quad (2) \\ \frac{37}{15}x_1 - 8s_2 + s_3 & = & 37 \quad (3) - 8 \cdot (2) \end{array}$$

- Corresponding basic solution: $s_1 = 23, x_2 = 6, s_3 = 37$.
- Nonbasic variables: $x_1 = s_2 = 0$
- The marginal value of x_1 is $\frac{9}{5} > 0$. Let x_1 enter the basis
- Which should leave? $s_1, x_2,$ or s_3 ?

Change basis ...

$$\begin{array}{rcll}
 -z & +\frac{9}{5}x_1 & & -3s_2 & = & -18 & (0) \\
 & x_1 & & +s_1 & = & 23 & (1) \\
 & \frac{1}{15}x_1 & +x_2 & & +s_2 & = & 6 & (2) \\
 & \frac{37}{15}x_1 & & -8s_2 & +s_3 & = & 37 & (3)
 \end{array}$$

- ▶ The value of x_1 can increase until some basic variable reaches the value 0:

$$\left. \begin{array}{l}
 (1) : s_1 = 23 - x_1 \geq 0 \Rightarrow x_1 \leq 23 \\
 (2) : x_2 = 6 - \frac{1}{15}x_1 \geq 0 \Rightarrow x_1 \leq 90 \\
 (3) : s_3 = 37 - \frac{37}{15}x_1 \geq 0 \Rightarrow x_1 \leq 15
 \end{array} \right\} \Rightarrow \begin{array}{l} s_3 = 0 \text{ when} \\ x_1 = 15 \end{array}$$

- ▶ x_1 enters the basis and s_3 will leave the basis


- ▶ Perform row operations:

$$\begin{array}{rcll}
 -z & & +2.84s_2 & -0.73s_3 & = & -45 & (0) - (3) \cdot \frac{15}{37} \cdot \frac{9}{5} \\
 & s_1 & +3.24s_2 & -0.41s_3 & = & 8 & (1) - (3) \cdot \frac{15}{37} \\
 & x_2 & +1.22s_2 & -0.03s_3 & = & 5 & (2) - (3) \cdot \frac{15}{37} \cdot \frac{1}{15} \\
 & x_1 & -3.24s_2 & +0.41s_3 & = & 15 & (3) \cdot \frac{15}{37}
 \end{array}$$

Optimal basic solution

$$\begin{array}{rcll}
 -z & & -0.87s_1 & -0.37s_3 & = & -52 \\
 & & 0.31s_1 & +s_2 & -0.12s_3 & = & 2.47 \\
 & x_2 & -0.37s_1 & & +0.12s_3 & = & 2 \\
 & x_1 & & +s_1 & & = & 23
 \end{array}$$

- ▶ No marginal value is positive. No improvement can be made
- ▶ The optimal basis is given by $s_2 = 2.47$, $x_2 = 2$, and $x_1 = 23$
- ▶ Non-basic variables: $s_1 = s_3 = 0$
- ▶ Optimal value: $z = 52$



Change basis ...

$$\begin{array}{rcll}
 -z & & +2.84s_2 & -0.73s_3 & = & -45 & (0) \\
 & s_1 & +3.24s_2 & -0.41s_3 & = & 8 & (1) \\
 & x_2 & +1.22s_2 & -0.03s_3 & = & 5 & (2) \\
 & x_1 & -3.24s_2 & +0.41s_3 & = & 15 & (3)
 \end{array}$$

- ▶ Let s_2 enter the basis (marginal value > 0)
- ▶ The value of s_2 can increase until some basic variable = 0:

$$\left. \begin{array}{l}
 (1) : s_1 = 8 - 3.24s_2 \geq 0 \Rightarrow s_2 \leq 2.47 \\
 (2) : x_2 = 5 - 1.22s_2 \geq 0 \Rightarrow s_2 \leq 4.10 \\
 (3) : x_1 = 15 + 3.24s_2 \geq 0 \Rightarrow s_2 \geq -4.63
 \end{array} \right\} \Rightarrow \begin{array}{l} s_1 = 0 \text{ when} \\ s_2 = 2.47 \end{array}$$

- ▶ s_2 enters the basis and s_1 will leave the basis

- ▶ Perform row operations:

$$\begin{array}{rcll}
 -z & & -0.87s_1 & -0.37s_3 & = & -52 & (0) - (1) \cdot \frac{2.84}{3.24} \\
 & & 0.31s_1 & +s_2 & -0.12s_3 & = & 2.47 & (1) \cdot \frac{1}{3.24} \\
 & x_2 & -0.37s_1 & & +0.12s_3 & = & 2 & (2) - (1) \cdot \frac{1.22}{3.24} \\
 & x_1 & & +s_1 & & = & 23 & (3) + (1)
 \end{array}$$

Summary of the solution course

basis	-z	x_1	x_2	s_1	s_2	s_3	RHS
-z	1	2	3	0	0	0	0
s_1	0	1	0	1	0	0	23
s_2	0	0.067	1	0	1	0	6
s_3	0	3	8	0	0	1	85
-z	1	1.80	0	0	-3	0	-18
s_1	0	1	0	1	0	0	23
x_2	0	0.07	1	0	1	0	6
s_3	0	2.47	0	0	-8	1	37
-z	1	0	0	0	2.84	-0.73	-45
s_1	0	0	0	1	3.24	-0.41	8
x_2	0	0	1	0	1.22	-0.03	5
x_1	0	1	0	0	-3.24	0.41	15
-z	1	0	0	-0.87	0	-0.37	-52
s_2	0	0	0	0.31	1	-0.12	2.47
x_2	0	0	1	-0.37	0	0.12	2
x_1	0	1	0	1	0	0	23

Summary of the simplex method

- ▶ **Optimality condition:** The *entering* variable in a maximization (minimization) problem should have the largest positive (negative) marginal value (reduced cost).

The entering variable determines a direction in which the objective value increases (decreases).

If all reduced costs are negative (positive), the current basis is optimal.

- ▶ **Feasibility condition:** The *leaving* variable is the one with smallest nonnegative ratio.

Corresponds to the constraint that is “reached first”



Simplex search for linear (minimization) programs

1. **Initialization:** Choose any feasible basis, construct the corresponding basic solution \mathbf{x}^0 , let $t = 0$
2. **Step direction:** Select a variable to enter the basis using the optimality condition (negative marginal value). Stop if no entering variable exists
3. **Step length:** Select a leaving variable using the feasibility condition (smallest non-negative ratio)
4. **New iterate:** Compute the new basic solution \mathbf{x}^{t+1} by performing matrix operations.
Let $t := t + 1$ and repeat from 2



Solve the lego problem using the simplex method!

$$\begin{aligned} \text{maximize } z &= 1600x_1 + 1000x_2 \\ \text{subject to } & 2x_1 + x_2 \leq 6 \\ & 2x_1 + 2x_2 \leq 8 \\ & x_1, x_2 \geq 0 \end{aligned}$$


ON THE BOARD!!



Degeneracy

- ▶ If the smallest nonnegative ratio is zero, the value of a basic variable will become zero in the next iteration
- ▶ The solution is *degenerate*
- ▶ The objective value will *not* improve in this iteration
- ▶ Risk for *cycling* around (non-optimal) bases
- ▶ Reason: a *redundant* constraint “touches” the feasible set
- ▶ Example:

$$\begin{aligned} x_1 + x_2 &\leq 6 \\ x_2 &\leq 3 \\ x_1 + 2x_2 &\leq 9 \\ x_1, x_2 &\geq 0 \end{aligned}$$



- ▶ Computational rules to prevent from cycling



Unbounded solutions

- ▶ If all ratios are negative, the variable entering the basis may increase infinitely
- ▶ The feasible set is unbounded
- ▶ In a real application this would probably be due to some incorrect assumption

▶ Example:

$$\begin{array}{ll} \text{minimize } z = & -x_1 - 2x_2 \\ \text{subject to} & -x_1 + x_2 \leq 2 \\ & -2x_1 + x_2 \leq 1 \\ & x_1, x_2 \geq 0 \end{array}$$

DRAW GRAPH!!



Starting solution—finding an initial base

- ▶ Example:

$$\begin{array}{ll} \text{minimize } z = & 2x_1 + 3x_2 \\ \text{subject to} & 3x_1 + 2x_2 = 14 \\ & 2x_1 - 4x_2 \geq 2 \\ & 4x_1 + 3x_2 \leq 19 \\ & x_1, x_2 \geq 0 \end{array}$$

DRAW GRAPH!!

- ▶ Add slack and surplus variables

$$\begin{array}{ll} \text{minimize } z = & 2x_1 + 3x_2 \\ \text{subject to} & 3x_1 + 2x_2 = 14 \\ & 2x_1 - 4x_2 - s_1 = 2 \\ & 4x_1 + 3x_2 + s_2 = 19 \\ & x_1, x_2, s_1, s_2 \geq 0 \end{array}$$

- ▶ How finding an initial basis? Only s_2 is obvious!



Unbounded solutions

- ▶ A feasible basis is given by $x_1 = 1$, $x_2 = 3$, with corresponding tableau:

Homework: Find this basis using the simplex method.

basis	-z	x_1	x_2	s_1	s_2	RHS
-z	1	0	0	5	-3	7
x_1	0	1	0	1	-1	1
x_2	0	0	1	2	-1	3

- ▶ Entering variable is s_2
- ▶ Row 1: $x_1 = 1 + s_2 \geq 0 \Rightarrow s_2 \geq -1$
- ▶ Row 2: $x_2 = 3 + s_2 \geq 0 \Rightarrow s_2 \geq -3$
- ▶ No leaving variable can be found, since no constraint will prevent s_2 from increasing infinitely



Artificial variables

- ▶ Add artificial variables a_1 and a_2 to the first and second constraints, respectively
- ▶ Solve an artificial problem: minimize $a_1 + a_2$

$$\begin{array}{ll} \text{minimize } w = & a_1 + a_2 \\ \text{subject to} & 3x_1 + 2x_2 + a_1 = 14 \\ & 2x_1 - 4x_2 - s_1 + a_2 = 2 \\ & 4x_1 + 3x_2 + s_2 = 19 \\ & x_1, x_2, s_1, s_2, a_1, a_2 \geq 0 \end{array}$$

- ▶ The “phase one” problem
- ▶ An initial basis is given by $a_1 = 14$, $a_2 = 2$, and $s_2 = 19$:

basis	-w	x_1	x_2	s_1	s_2	a_1	a_2	RHS
-w	1	-5	2	1	0	0	0	-16
a_1	0	3	2	0	0	1	0	14
a_2	0	2	-4	-1	0	0	1	2
s_2	0	4	3	0	1	0	0	19



Find an initial solution using artificial variables

- ▶ x_1 enters $\Rightarrow a_2$ leaves (then $x_2 \Rightarrow s_2$, then $s_1 \Rightarrow a_1$)

basis	-w	x_1	x_2	s_1	s_2	a_1	a_2	RHS
-w	1	-5	2	1	0	0	0	-16
a_1	0	3	2	0	0	1	0	14
a_2	0	2	-4	-1	0	0	1	2
s_2	0	4	3	0	1	0	0	19
<hr/>								
-w	1	0	-8	-1.5	0	0		-11
a_1	0	0	8	1.5	0	1		11
x_1	0	1	-2	-0.5	0	0		1
s_2	0	0	11	2	1	0		15
<hr/>								
-w	1	0	0	-0.045	0.727	0		-0.091
a_1	0	0	0	0.045	-0.727	1		0.091
x_1	0	1	0	-0.136	0.182	0		3.727
x_2	0	0	1	0.182	0.091	0		1.364
<hr/>								
-w	1	0	0	0	0			0
s_1	0	0	0	1	-16			2
x_1	0	1	0	0	-2			4
x_2	0	0	1	0	3			1

- ▶ A feasible basis is given by $x_1 = 4$, $x_2 = 1$, and $s_1 = 2$

Infeasible linear programs

- ▶ If the solution to the “phase one” problem has optimal value = 0, a feasible basis has been found
 - ⇒ Start optimizing the original objective function z from this basis (*homework*)
 - ▶ If the solution to the “phase one” problem has optimal value $w > 0$, no feasible solutions exist
 - ▶ What would this mean in a real application?
 - ▶ Alternative: M -method (Big- M method): Add the artificial variables to the original objective—with a large coefficient
- Example:

$$\text{minimize } z = 2x_1 + 3x_2$$

$$\Rightarrow \text{minimize } z_a = 2x_1 + 3x_2 + Ma_1 + Ma_2$$

Alternative optimal solutions

- ▶ Example:

$$\begin{aligned} \text{minimize } z &= 2x_1 + 4x_2 \\ \text{subject to } &x_1 + 2x_2 \leq 5 \\ &x_1 + x_2 \leq 4 \\ &x_1, x_2 \geq 0 \end{aligned}$$

DRAW GRAPH!!

- ▶ The extreme points $(0, \frac{5}{2})$ and $(3, 1)$ have the same optimal value $z = 10$
- ▶ All solutions that are positive linear (convex) combinations of these are optimal:

$$(x_1, x_2) = \alpha \cdot (0, \frac{5}{2}) + (1 - \alpha) \cdot (3, 1), \quad 0 \leq \alpha \leq 1$$