

## Fö 8.2

Alla dessa integraler.

1. Envariabelintegral:  $\int_I f dx = \int_a^b f(x) dx = \lim_{N \rightarrow \infty} \sum_{i=1}^N f(x_i) \Delta x_i$

Variabelbyte:  $x = g(u), dx = g'(u) du$


$$\int_a^b f(x) dx = \int_{g(a)}^{g(b)} f(g(u)) g'(u) du$$

Fundamentalsatsen:  $\int_a^b Df(x) dx = [f(x)]_a^b$

Kurvintegral:  $r = r(t), t \in [a, b]$   
 $ds = |r'(t)| dt$

$$\int_C f ds = \int_a^b f(r(t)) |r'(t)| dt$$

Längd:  $L = \int_C ds$



Tangentkurvintegral:

$$dr = r'(t) dt = \frac{r'(t)}{|r'(t)|} |r'(t)| dt = \hat{T} ds$$

$$\int_C F \cdot dr = \int_C F \cdot \hat{T} ds = \int_a^b F(r(t)) \cdot r'(t) dt$$

Tolkning: arbete.

Fundamentalsats: om  $F = \nabla \phi$  så är

$$\int_C F \cdot dr = \int_C \nabla \phi \cdot dr = \phi(r(b)) - \phi(r(a))$$



2. Dubbelintegral:  $\iint_R f dA = \lim \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \Delta x_i \Delta y_j$

Upprepad integration:

$$\iint_R f dA = \int_a^b \int_{c(x)}^{d(x)} f(x, y) dy dx$$



eller

$$\iint_R f dA = \int_c^d \int_{a(y)}^{b(y)} f(x, y) dx dy$$



Area:  $A = \iint_R dA$

Variabelbyte:  $\begin{cases} x = x(u, v), \\ y = y(u, v), \end{cases} \quad (u, v) \in D, \quad R = R(u, v)$





$$dA = \left| \frac{\partial R}{\partial u} \times \frac{\partial R}{\partial v} \right| du dv = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

Koordinatkurvor, tangenter, jacobideterminant.

$$\begin{aligned} \iint_R f dA &= \iint_D f(R(u, v)) \left| \frac{\partial R}{\partial u} \times \frac{\partial R}{\partial v} \right| du dv \\ &= \iint_D f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv \end{aligned}$$

Yta:  $\mathcal{S} : \mathbb{R}^3 = \mathbb{R}(u, v), (u, v) \in D$

$$\varphi : \begin{cases} x = x(u, v), \\ y = y(u, v), \\ z = z(u, v), \end{cases} (u, v) \in D$$



Tangenter:  $\frac{\partial \mathbb{R}}{\partial u}, \frac{\partial \mathbb{R}}{\partial v}$

Normalvektor:  $N = \pm \frac{\partial \mathbb{R}}{\partial u} \times \frac{\partial \mathbb{R}}{\partial v}$

Enhetsnormal:  $\hat{N} = \frac{N}{|N|}$

$$dS = |N| du dv = \left| \frac{\partial \mathbb{R}}{\partial u} \times \frac{\partial \mathbb{R}}{\partial v} \right| du dv$$

Ytintegral:  $\iint_{\mathcal{S}} f dS = \iint_D f(\mathbb{R}(u, v)) \left| \frac{\partial \mathbb{R}}{\partial u} \times \frac{\partial \mathbb{R}}{\partial v} \right| du dv$


Area:  $A = \iint_{\mathcal{S}} dS$

Flödesintegral (normalytintegral):

$$\iint_{\mathcal{S}} \mathbb{F} \cdot d\mathbb{S} = \iint_{\mathcal{S}} \mathbb{F} \cdot \hat{N} dS = \pm \iint_D \mathbb{F}(\mathbb{R}(u, v)) \cdot \left( \frac{\partial \mathbb{R}}{\partial u} \times \frac{\partial \mathbb{R}}{\partial v} \right) du dv$$

Fundamentalsats: Stokes

$$\iint_{\mathcal{S}} (\nabla \times \mathbb{F}) \cdot \hat{N} dS = \int_{\mathcal{C}} \mathbb{F} \cdot d\mathbb{R}$$




Exempel Graf.  $y = f(x), x \in [a, b]$

$$\begin{cases} x = t \\ y = f(t), t \in [a, b] \end{cases}$$

$$r'(t) = i + f'(t)j$$

$$ds = \sqrt{1 + f'(t)^2} dt$$



Exempel Graf.  $z = f(x, y), x, y \in D$

$$\begin{cases} x = u \\ y = v \\ z = f(u, v) \end{cases}$$


$$\frac{\partial r}{\partial u} = i + f'_u k$$

$$\frac{\partial r}{\partial v} = j + f'_v k$$

$$\frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} = \begin{vmatrix} i & j & k \\ 1 & 0 & f'_u \\ 0 & 1 & f'_v \end{vmatrix} = -f'_u i - f'_v j + k$$

$$dS = \sqrt{1 + f'_u{}^2 + f'_v{}^2} du dv$$

$$d\mathcal{S} = \hat{N} dS = \pm (-f'_u i - f'_v j + k) du dv$$




Viktiga koordinater

Polära  $(r, \theta)$   $\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$


Cylindriska  $(r, \theta, z)$   $\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases}$

Sfäriska  $(\rho, \phi, \theta)$   $\begin{cases} x = \rho \sin \phi \cos \theta \\ y = \rho \sin \phi \sin \theta \\ z = \rho \cos \phi \end{cases}$




Trippelintegral:  $\iiint_R f dV = \lim \sum_i \sum_j \sum_k f(x_i, y_j, z_k) \Delta x_i \Delta y_j \Delta z_k$

Upprepad integration:



$$\begin{aligned} \iiint_R f dV &= \int_a^b \iint_{D(x)} f(x, y, z) dA dx = \\ &= \int_a^b \int_{c(x)}^{d(x)} \int_{h(x,y)}^{k(x,y)} f(x, y, z) dz dy dx \end{aligned}$$




Volym:  $V = \iiint_R dV$

Tyngdpunkt:  $\bar{x} = \frac{1}{V} \iiint_R x dV$ , samma för  $\bar{y}, \bar{z}$ .

Variabelbyte:  $\mathcal{R} = \mathcal{R}(u, v, w), (u, v, w) \in D$


$$\iiint_R f dV = \iiint_D f(\mathcal{R}(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw$$

$$dV = \left| \left( \frac{\partial \mathcal{R}}{\partial u} \times \frac{\partial \mathcal{R}}{\partial v} \right) \cdot \frac{\partial \mathcal{R}}{\partial w} \right| du dv dw$$



Fundamentalsats: Gauss

$$\iiint_D \nabla \cdot \mathbf{F} dV = \iint_{\mathcal{P}} \mathbf{F} \cdot \hat{\mathbf{N}} dS$$



Ga igenom tenta 130116 nr 5,6