

MATHEMATICS, Chalmers and Göteborg University

Tentamen i Financial Derivative and Stochastic Analysis(CTH-TMA285) (GU-MAM695)

December 16 2005, 8.30 to 12.30 at *Väg och vatten*.

Telephone back-up: Torbjörn Lundh, 0731-526320.

Allowable handbook: β eta. No calculators are allowed.

1. (3p) We will study a sequence of dice-tosses. Let D_i be the outcome of the toss number i , and let us define the random variable

$$X(n) := \sum_{i=1}^n D_i.$$

- (a) Compute

$$\mathbb{E}(X(n)|\mathcal{F}_{n-1}).$$

- (b) Construct a random variable $Z(n)$ based on $X(n)$ such that $Z(n) \rightarrow W(1)$ as $n \rightarrow \infty$, where $W(t)$ is the usual Brownian motion.
- (c) Compare your method for estimating $W(1)$ with the standard coin tossing method to generate an approximation of $W(1)$.
2. (3p) (Vasiček's interest rate model)

Let

$$dR(t) = (\alpha - \beta R(t)) dt + \sigma dW(t),$$

where α , β , and σ are positive constants.

- (a) Compute and simplify $d(e^{\beta t}R(t))$ to a form not including $R(t)$.
- (b) Integrate the equation you got above and solve for $R(t)$.
- (c) Use the formula above to compute $\mathbb{E}[R(t)]$.
3. (3p) Suppose we have a model of market including two stocks with values $S_1(t)$ and $S_2(t)$ at time t . Let us also assume that the interest rate, r , is constant in this model. The two stocks are modeled in the standard way, i.e.

$$dS_1(t) = \alpha_1 S_1(t) dt + \sigma_1 S_1(t) dW_1(t),$$

$$dS_2(t) = \alpha_2 S_2(t) dt + \sigma_2 S_2(t) dW_2(t),$$

where α_i and σ_i are constants. Furthermore, in this model we also have that $\alpha_1 = \alpha_2$, $\sigma_2 = 2\sigma_1$ and $W_1(t) = W_2(t)$ for all $t \geq 0$.

- (a) Define risk-neutral probability measure.
- (b) Show that in the market model described above, there does not exist any risk-neutral probability measure.
4. (3p) Let $X(t) = at + bW(t)$, where a and b are non-zero constants and $W(t)$ a Brownian motion. Let f be a real valued function which is (at least) two times differentiable and where $f(X(t))$ is a martingale. Describe the class of such functions.

5. (3p) Let $v(t, S(t))$ denote the price at time t of an up-and-out call with barrier B , strike price K , and maturity date T . Furthermore, suppose that the volatility σ and the interest rate r are constant. At a given time $t_0 \in (0, T]$, we assume that the underlying asset $S(t)$ is in $(0, B)$ and that the Greeks are known, i.e. $v_x(t_0, x_0) = \delta$, $v_t(t_0, x_0) = \theta$, and $v_{xx}(t_0, x_0) = \gamma$, where $x_0 = S(t_0)$.

- (a) What is the value of the up-and-out call at time t_0 ?
 (b) Suppose $t_0 = T$. Show that

$$\frac{\delta^2}{\gamma\theta} \geq 2\sigma^2$$

when $S(T) < K$ and

$$(\delta - 1)^2 r^2 \geq 2\gamma\sigma^2(Kr + \theta)$$

when $S(T) > K$. (In this case we naturally use the left derivative for θ at T .)

6. (4p) (Quadratic variation of an Itô-process)

- (a) Define the quadratic variation of a function $f(t)$ defined for all $t \in [0, T]$.
 (b) Let

$$X(t) = X(0) + \int_0^t \Delta(u) dW(u) + \int_0^t \Theta(u) du,$$

where $W(t)$ is a Brownian motion, $X(0)$ is nonrandom, and $\Delta(u)$ and $\Theta(u)$ are adapted stochastic processes. Derive the quadratic variation of the process $X(t)$.

7. (4p) (Girsanov's Theorem) State and prove Girsanov's Theorem in one dimension. (You do not need to prove other theorems or lemmas which you might use in your proof as long as you clearly state the results you are using.)