

Lecture 3: Lagrangian duality

Michael Patriksson

2010-09-06

The Relaxation Theorem

- ▶ Problem: find

$$f^* = \infimum_{\mathbf{x}} f(\mathbf{x}), \quad (1a)$$

$$\text{subject to } \mathbf{x} \in S, \quad (1b)$$

where $f : \mathbb{R}^n \mapsto \mathbb{R}$ is a given function and $S \subseteq \mathbb{R}^n$

- ▶ A *relaxation* to (1a)–(1b) has the following form: find

$$f_R^* = \infimum_{\mathbf{x}} f_R(\mathbf{x}), \quad (2a)$$

$$\text{subject to } \mathbf{x} \in S_R, \quad (2b)$$

where $f_R : \mathbb{R}^n \mapsto \mathbb{R}$ is a function with $f_R \leq f$ on S and $S_R \supseteq S$

Relaxation example (maximization)

- ▶ Binary knapsack problem:

$$\begin{aligned} z^* = \text{maximize} \quad & 7x_1 + 4x_2 + 5x_3 + 2x_4 \\ & \mathbf{x} \in \{0,1\}^4 \\ \text{subject to} \quad & 3x_1 + 3x_2 + 4x_3 + 2x_4 \leq 5 \end{aligned}$$

- ▶ Optimal solution: $\mathbf{x}^* = (1, 0, 0, 1)$, $z^* = 9$

- ▶ Continuous relaxation:

$$\begin{aligned} z_{\text{LP}}^* = \text{maximize} \quad & 7x_1 + 4x_2 + 5x_3 + 2x_4 \\ & \mathbf{x} \in [0,1]^4 \\ \text{subject to} \quad & 3x_1 + 3x_2 + 4x_3 + 2x_4 \leq 5 \end{aligned}$$

- ▶ Optimal solution: $\mathbf{x}_R^* = (1, \frac{2}{3}, 0, 0)$, $z_R^* = 9\frac{2}{3} > z^*$

- ▶ \mathbf{x}_R^* is *not feasible* in the binary problem

The relaxation theorem

1. [relaxation] $f_R^* \leq f^*$
2. [infeasibility] *If (2) is infeasible, then so is (1)*
3. [optimal relaxation]
If the problem (2) has an optimal solution $\mathbf{x}_R^ \in S$ for which*

$$f_R(\mathbf{x}_R^*) = f(\mathbf{x}_R^*),$$

then \mathbf{x}_R^ is an optimal solution to (1) as well*

- *Proof portion.* For 3., note that

$$f(\mathbf{x}_R^*) = f_R(\mathbf{x}_R^*) \leq f_R(\mathbf{x}) \leq f(\mathbf{x}), \quad \mathbf{x} \in S$$

Lagrangian relaxation, I

- ▶ Consider the optimization problem:

$$f^* = \underset{\mathbf{x}}{\text{infimum}} f(\mathbf{x}), \quad (3a)$$

$$\text{subject to } \mathbf{x} \in X, \quad (3b)$$

$$g_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m, \quad (3c)$$

where $f : \mathbb{R}^n \mapsto \mathbb{R}$ and $g_i : \mathbb{R}^n \mapsto \mathbb{R}$ ($i = 1, 2, \dots, m$) are given functions, and $X \subseteq \mathbb{R}^n$

- ▶ Here we assume that

$$-\infty < f^* < \infty, \quad (4)$$

that is, that f is bounded from below and that the problem has at least one feasible solution

- ▶ For a vector $\boldsymbol{\mu} \in \mathbb{R}^m$, we define the *Lagrange function*

$$L(\mathbf{x}, \boldsymbol{\mu}) = f(\mathbf{x}) + \sum_{i=1}^m \mu_i g_i(\mathbf{x}) = f(\mathbf{x}) + \boldsymbol{\mu}^T \mathbf{g}(\mathbf{x})$$

- ▶ We call the vector $\boldsymbol{\mu}^* \in \mathbb{R}^m$ a *Lagrange multiplier* if it is non-negative and if $f^* = \inf_{\mathbf{x} \in X} L(\mathbf{x}, \boldsymbol{\mu}^*)$ holds

Lagrange multipliers and global optima

- ▶ Let μ^* be a Lagrange multiplier.
Then, \mathbf{x}^* is an optimal solution to

$$f^* = \inf \{ f(\mathbf{x}) \mid \mathbf{x} \in X, g_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \},$$

if and only if it is feasible and

$$\mathbf{x}^* \in \arg \min_{\mathbf{x} \in X} L(\mathbf{x}, \mu^*), \quad \text{and} \quad \mu_i^* g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, m$$

- ▶ Notice the resemblance to the KKT conditions:
 - ▶ If $X = \mathbb{R}^n$ and all functions are in C^1 then
“ $\mathbf{x}^* \in \arg \min_{\mathbf{x} \in X} L(\mathbf{x}, \mu^*)$ ” \Leftrightarrow “force equilibrium condition”,
i.e., the first row of the KKT conditions
 - ▶ The second item, “ $\mu_i^* g_i(\mathbf{x}^*) = 0$ for all i ” \Leftrightarrow complementarity conditions

The Lagrangian dual problem associated with the Lagrangian relaxation

- ▶ The *Lagrangian dual function* is

$$q(\boldsymbol{\mu}) = \inf_{\mathbf{x} \in X} L(\mathbf{x}, \boldsymbol{\mu})$$

- ▶ The *Lagrangian dual problem* is to

$$q^* = \underset{\boldsymbol{\mu} \geq \mathbf{0}^m}{\text{maximize}} \quad q(\boldsymbol{\mu}) \quad (5)$$

- ▶ For some $\boldsymbol{\mu}$, $q(\boldsymbol{\mu}) = -\infty$ is possible. If this is true for all $\boldsymbol{\mu} \geq \mathbf{0}^m$ then

$$q^* = \underset{\boldsymbol{\mu} \geq \mathbf{0}^m}{\text{supremum}} \quad q(\boldsymbol{\mu}) = -\infty$$

The Lagrangian dual problem, cont'd

- ▶ The *effective domain* of q is $D_q = \{ \boldsymbol{\mu} \in \mathbb{R}^m \mid q(\boldsymbol{\mu}) > -\infty \}$

[Theorem] D_q is convex, and q is concave on D_q □

- ▶ Very good news: The Lagrangian dual problem is always convex!
- ▶ Maximize a concave function (even continuous as long as $D_q = \mathbb{R}^m$)
- ▶ Need still to show how a Lagrangian dual optimal solution can be used to generate a primal optimal solution

Weak Duality Theorem

Let \mathbf{x} and $\boldsymbol{\mu}$ be feasible in

$$f^* = \inf\{f(\mathbf{x}) \mid \mathbf{x} \in X, g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$$

and

$$q^* = \max\{q(\boldsymbol{\mu}) \mid \boldsymbol{\mu} \geq \mathbf{0}^m\},$$

respectively. Then,

$$q(\boldsymbol{\mu}) \leq f(\mathbf{x})$$

In particular,

$$q^* \leq f^*$$

If $q(\boldsymbol{\mu}) = f(\mathbf{x})$, then the pair $(\mathbf{x}, \boldsymbol{\mu})$ is optimal in the respective problem and

$$q^* = q(\boldsymbol{\mu}) = f(\mathbf{x}) = f^*$$

□

Weak Duality Theorem, cont'd

- ▶ Weak duality is also a consequence of the Relaxation Theorem: For any $\boldsymbol{\mu} \geq \mathbf{0}^m$, let

$$\begin{aligned}S &= X \cap \{ \mathbf{x} \in \mathbb{R}^n \mid \mathbf{g}(\mathbf{x}) \leq \mathbf{0}^m \}, \\S_R &= X, \\f_R &= L(\boldsymbol{\mu}, \cdot)\end{aligned}$$

Apply the Relaxation Theorem

- ▶ If $q^* = f^*$, there is *no duality gap*
- ▶ If there exists a Lagrange multiplier vector, then by the weak duality theorem, there is no duality gap

Global optimality conditions

- ▶ The vector $(\mathbf{x}^*, \boldsymbol{\mu}^*)$ is a pair of an optimal primal solution and a Lagrange multiplier if and only if

$$\boldsymbol{\mu}^* \geq \mathbf{0}^m, \quad (\text{Dual feasibility}) \quad (6a)$$

$$\mathbf{x}^* \in \arg \min_{\mathbf{x} \in X} L(\mathbf{x}, \boldsymbol{\mu}^*), \quad (\text{Lagrangian optimality}) \quad (6b)$$

$$\mathbf{x}^* \in X, \quad \mathbf{g}(\mathbf{x}^*) \leq \mathbf{0}^m, \quad (\text{Primal feasibility}) \quad (6c)$$

$$\mu_i^* g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, m \quad (\text{Complementary slackness}) \quad (6d)$$

- ▶ If $\exists(\mathbf{x}^*, \boldsymbol{\mu}^*)$ that fulfil (6), then there is a zero duality gap and Lagrange multipliers exist

Saddle points

- ▶ *The vector $(\mathbf{x}^*, \boldsymbol{\mu}^*)$ is a pair of an optimal primal solution and a Lagrange multiplier if and only if $\mathbf{x}^* \in X$, $\boldsymbol{\mu}^* \geq \mathbf{0}^m$, and $(\mathbf{x}^*, \boldsymbol{\mu}^*)$ is a saddle point of the Lagrangian function on $X \times \mathbb{R}_+^m$, that is,*

$$L(\mathbf{x}^*, \boldsymbol{\mu}) \leq L(\mathbf{x}^*, \boldsymbol{\mu}^*) \leq L(\mathbf{x}, \boldsymbol{\mu}^*), \quad (\mathbf{x}, \boldsymbol{\mu}) \in X \times \mathbb{R}_+^m,$$

holds

- ▶ If $\exists(\mathbf{x}^*, \boldsymbol{\mu}^*)$, equivalent to the global optimality conditions, the existence of Lagrange multipliers, and a zero duality gap

Strong duality for convex programs, introduction

- ▶ Convexity of the dual problem comes with very few assumptions on the original, primal problem
- ▶ The characterization of the primal–dual set of optimal solutions is also quite easily established
- ▶ To establish *strong duality*—sufficient conditions under which there is no duality gap—takes much more
- ▶ In particular—as with the KKT conditions—we need regularity conditions (constraint qualifications) and separation theorems

Strong duality theorem

- ▶ Consider the problem (3), that is,

$$f^* = \inf\{f(\mathbf{x}) \mid \mathbf{x} \in X, g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\},$$

where $f : \mathbb{R}^n \mapsto \mathbb{R}$ and g_i ($i = 1, \dots, m$) are *convex* and $X \subseteq \mathbb{R}^n$ is a *convex* set

- ▶ Introduce the following constraint qualification (CQ):

$$\exists \mathbf{x} \in X \text{ with } \mathbf{g}(\mathbf{x}) < \mathbf{0}^m \tag{7}$$

Strong duality theorem

Suppose that $-\infty < f^ < \infty$, and that the CQ (7) holds for the (convex) problem (3)*

- (a) There is no duality gap and there exists at least one Lagrange multiplier μ^* . Moreover, the set of Lagrange multipliers is bounded and convex*
- (b) If infimum in (3) is attained at some \mathbf{x}^* , then the pair (\mathbf{x}^*, μ^*) satisfies the global optimality conditions (6)*
- (c) If the functions f and g_i are in C^1 and X is open (for example, $X = \mathbb{R}^n$) then (6) equals the KKT conditions*

If all constraints are linear we can remove the CQ (7)

Example I: An explicit, differentiable dual problem

- ▶ Consider the problem to

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && f(\mathbf{x}) := x_1^2 + x_2^2, \\ & \text{subject to} && x_1 + x_2 \geq 4, \\ & && x_j \geq 0, \quad j = 1, 2 \end{aligned}$$

- ▶ Let

$$g(\mathbf{x}) = -x_1 - x_2 + 4$$

and

$$X = \{ (x_1, x_2) \mid x_j \geq 0, j = 1, 2 \} = \mathbb{R}_+^2$$

Example I, cont'd

- ▶ The Lagrangian dual function is

$$\begin{aligned}q(\mu) &= \min_{\mathbf{x} \in X} L(\mathbf{x}, \mu) := f(\mathbf{x}) + \mu(-x_1 - x_2 + 4) \\ &= 4\mu + \min_{\mathbf{x} \geq \mathbf{0}} \{x_1^2 + x_2^2 - \mu x_1 - \mu x_2\} \\ &= 4\mu + \min_{x_1 \geq 0} \{x_1^2 - \mu x_1\} + \min_{x_2 \geq 0} \{x_2^2 - \mu x_2\}, \quad \mu \geq 0\end{aligned}$$

- ▶ For a fixed $\mu \geq 0$, the minimum is attained at $x_1(\mu) = \frac{\mu}{2}, x_2(\mu) = \frac{\mu}{2}$
- ▶ Substituting this expression into $q(\mu) \Rightarrow$
 $q(\mu) = f(\mathbf{x}(\mu)) + \mu(-x_1(\mu) - x_2(\mu) + 4) = 4\mu - \frac{\mu^2}{2}$
- ▶ Note that q is *strictly concave*, and it is differentiable everywhere (since f, g are differentiable and $\mathbf{x}(\mu)$ is unique)

Example I, cont'd

- ▶ Recall the dual problem

$$q^* = \max_{\mu \geq 0} q(\mu) = \max_{\mu \geq 0} \left(4\mu - \frac{\mu^2}{2} \right)$$

- ▶ We have that $q'(\mu) = 4 - \mu = 0 \iff \mu = 4$
As $4 \geq 0$, this is the optimum in the dual problem!

$$\Rightarrow \mu^* = 4 \text{ and } \mathbf{x}^* = (x_1(\mu^*), x_2(\mu^*))^T = (2, 2)^T$$


- ▶ Also: $f(\mathbf{x}^*) = q(\mu^*) = 8$
- ▶ Here, the dual function is *differentiable*. The optimum \mathbf{x}^* is also unique and automatically given by $\mathbf{x}^* = \mathbf{x}(\mu^*)$

Example II: Implicit non-differentiable dual problem

- ▶ Consider the linear programming problem to

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && f(\mathbf{x}) := -x_1 - x_2, \\ & \text{subject to} && 2x_1 + 4x_2 \leq 3, \\ & && 0 \leq x_1 \leq 2, \\ & && 0 \leq x_2 \leq 1 \end{aligned}$$

- ▶ The optimal solution is $\mathbf{x}^* = (3/2, 0)^T$, $f(\mathbf{x}^*) = -3/2$




Example II: Lagrangian relax the first constraint

$$L(\mathbf{x}, \mu) = -x_1 - x_2 + \mu(2x_1 + 4x_2 - 3);$$

$$q(\mu) = -3\mu + \min_{0 \leq x_1 \leq 2} \{(-1 + 2\mu)x_1\} + \min_{0 \leq x_2 \leq 1} \{(-1 + 4\mu)x_2\}$$

$$= \begin{cases} -3 + 5\mu, & 0 \leq \mu \leq 1/4, & \Leftrightarrow x_1(\mu) = 2, x_2(\mu) = 1 \\ -2 + \mu, & 1/4 \leq \mu \leq 1/2, & \Leftrightarrow x_1(\mu) = 2, x_2(\mu) = 0 \\ -3\mu, & 1/2 \leq \mu & \Leftrightarrow x_1(\mu) = x_2(\mu) = 0 \end{cases}$$



$$\mu^* = \frac{1}{2}, \quad q(\mu^*) = -\frac{3}{2}$$

Example II, cont'd

- ▶ For linear (convex) programs strong duality holds, but how obtain \mathbf{x}^* from μ^* ?
- ▶ q is non-differentiable at $\mu^* \Rightarrow$ Utilize characterization in (6)
- ▶ The subproblem solution set at μ^* is
$$X(\mu^*) = \left\{ \begin{pmatrix} 2\alpha \\ 0 \end{pmatrix} \mid 0 \leq \alpha \leq 1 \right\}$$
- ▶ Among the subproblem solutions, we next have to find one that is primal feasible as well as complementary
- ▶ Primal feasibility means that $2 \cdot 2\alpha + 4 \cdot 0 \leq 3 \iff \alpha \leq 3/4$
- ▶ Complementarity means that
$$\mu^* \cdot (2x_1^* + 4x_2^* - 3) = 0 \iff \alpha = 3/4, \text{ since } \mu^* \neq 0$$
- ▶ Conclusion: the only primal vector \mathbf{x} that satisfies the system (6) together with the dual solution $\mu^* = 1/2$ is $\mathbf{x}^* = (3/2, 0)^T$
- ▶ Observe finally that $f^* = q^*$

A theoretical argument for $\mu^* = 1/2$

- ▶ Due to the global optimality conditions, the optimal solution must in this convex case be among the subproblem solutions
- ▶ Since x_1^* is not in one of the “corners” of X ($0 < x_1^* < 2$), the value of μ^* must be such that the cost term for x_1 in $L(\mathbf{x}, \mu^*)$ is zero! That is, $-1 + 2\mu^* = 0 \Rightarrow \mu^* = 1/2!$
- ▶ A non-coordinability phenomenon—a non-unique subproblem solution means that the optimal solution is not obtained automatically
- ▶ In non-convex cases (e.g., integrality constraints) the optimal solution may not be among the points in $X(\mu^*)$ (the set of subproblem solutions at μ^*)
- ▶ What do we do then??