

Optimality conditions for unconstrained and convexly constrained optimization.

Niclas Andréasson

29 januari 2004

Uppvärmning

Övning 1. Find the rectangular parallelepiped of unit volume that has the minimum surface area.

□

Unconstrained optimization

Öving 2. Consider the parametric minimization problem to

$$(1) \quad \underset{x_1, x_2}{\text{minimize}} \quad \frac{3}{2}(x_1^2 + x_2^2) + (1 + a)x_1x_2 - (x_1 + x_2) + b,$$

where a and b are some unknown real-valued parameters.

Find all possible values of a and b such that the problem (1) possesses a

unique globally optimal solution. Write down this solution (in terms of the

parameters a and b).

□

Övning 3. Let A be a symmetric $n \times n$ matrix. For $x \in \mathbb{R}^n$ such that $x \neq \mathbf{0}_n$, consider the function

$$p(x) = \frac{x^T A x}{x^T x},$$

and the related optimization problem to

$$\text{minimize}_{x \neq \mathbf{0}_n} p(x).$$

Determine all the stationary points as well as the global minima in the minimization problem (2).

□

Constrained optimization

Övning 4. Among all rectangles contained in a given circle, show that the one that has maximal area must be a square. \square

Övning 5. Consider the positive orthant

$$S = \{ \mathbf{x} \in \mathbb{R}^n \mid \mathbf{x} \geq \mathbf{0} \}.$$

Derive necessary optimality conditions for the problem to

minimize $f(\mathbf{x})$

subject to $\mathbf{x} \in S$,

where $f \in C^1$.

□

Övning 6. Consider the problem to

maximize $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$

subject to $\sum_{i=1}^n x_i = 1$, $x_i \geq 0$, $i = 1, \dots, n$,

where a_i are given positive scalars. Find a global maximum and show that it is unique. \square