

Lecture 5: Primal-dual optimality conditions

Overview

- Want to establish that \boldsymbol{x}^* local minimum of f over S implies that a well-defined condition holds that we can easily check
- This is possible when constraints are linear, since the set of feasible directions then can be stated simply
- With non-linear constraints things become more complicated
- *Constraint qualifications* CQ are needed to make sure that the *well-defined* condition is a necessary condition for local optimality (rule out strange cases)
- Under convexity, the condition turns out to also always (under no CQ) be sufficient for global optimality
- Called the *Karush–Kuhn–Tucker* conditions
- Karush: master's student at Univ. of Chicago, 1939
Tucker/Kuhn: prof./Ph.D. student at Princeton Univ., 1951

- Of course, a globally optimal solution must then satisfy the KKT conditions. But it is *not* practical to search for all KKT points and pick the best. Its use is for checking that an algorithm has found the right solution
- Compare checking for every x with $f'(x) = 0$ in \mathbb{R} !
- The user has all the responsibility!

Cautions needed!

- Costly errors can be made if one ignores that KKT conditions are *necessary*, but not always *sufficient*
- US Air Force's B-2 Stealth bomber program: Reaganism, 1980s
- Design variables: various dimensions, distribution of volume between wing and fuselage, flying speed, thrust, fuel consumption, drag, lift, air density, etc
- Objective: maximum range on full tank
- Model from the 1940s which had produced B-29, B-52, etc
- Solution to the KKT conditions found; specified design variable values that put almost all of the total volume in the wing, leading to the *flying wing design* for the B-2 bomber
- Billions of dollars later, found the design solution works, but its range too low in comparison with other bomber designs

- Review carried out. The model is correct!
- But ... The model was a nonconvex NLP; the review revealed a second solution to the KKT system
- Much less wing volume! Looks like an airplane! Maximizes range!
- In other words, the design implemented was the aerodynamically *worst* possible choice of configuration, leading to a very costly error
- Still flies. Why? Happens that it has good properties wrt. radar protection (stealth) ...

Nice photos, I



Nice photos, II



Overview, cont'd

- The condition must not only be easy to check, it should also state something useful
- It is easy to state some condition: *If x^* is a local minimum of f over S then it is also feasible*
- Completely useless, since it is satisfied for every feasible point
- That is what we end up with if we want something that is applicable to every problem. We need to get rid of some weird problems, and that is a main reason for introducing the CQs
- We begin by studying an abstract problem and provide a *geometric optimality condition*
- Next, we state the corresponding result for an explicit representation of S in terms of constraints. This is the *Fritz John condition*

- Introducing a CQ we then obtain the *Karush–Kuhn–Tucker* conditions
- There is more than one CQ, some more useful than others in particular cases
- *Linear independence of the equality constraints* is the classic one from the Lagrange multiplier rule. We extend it and show others

Geometric optimality conditions

Problem:

$$\begin{aligned} & \text{minimize } f(\mathbf{x}), \\ & \text{subject to } \mathbf{x} \in S, \end{aligned} \tag{1}$$

$S \subset \mathbb{R}^n$ nonempty, closed; $f : \mathbb{R}^n \rightarrow \mathbb{R}$ in C^1

- Idea: at a local minimum \mathbf{x}^* of f over S it is impossible to draw a curve from \mathbf{x}^* such that it is feasible and f decreases along it
- Cannot work with f itself; descent is measured in terms of directional derivatives. Linearize f
- We must also “linearize” S . Reason: the cone of feasible directions may be too small to be useful; also, it is difficult to state it explicitly. We replace the cone of feasible directions with the *tangent cone* to S at \mathbf{x}^*

- The *cone of feasible directions* for S at $\mathbf{x} \in \mathbb{R}^n$ is

$$R_S(\mathbf{x}) := \{ \mathbf{p} \in \mathbb{R}^n \mid \exists \tilde{\delta} > 0 \text{ such that } \mathbf{x} + \delta \mathbf{p} \in S, 0 \leq \delta \leq \tilde{\delta} \}$$

- The *tangent cone* for S at $\mathbf{x} \in \mathbb{R}^n$ is

$$T_S(\mathbf{x}) := \{ \mathbf{p} \in \mathbb{R}^n \mid \exists \{ \mathbf{x}_k \} \subset S, \{ \lambda_k \} \subset (0, \infty) : \lim_{k \rightarrow \infty} \mathbf{x}_k = \mathbf{x}, \\ \lim_{k \rightarrow \infty} \lambda_k (\mathbf{x}_k - \mathbf{x}) = \mathbf{p} \}$$

- $T_S(\mathbf{x})$ is closed; the set of tangents to sequences $\{ \mathbf{x}_k \} \subset S$
- It holds that $\text{cl } R_S(\mathbf{x}) \subset T_S(\mathbf{x})$ for every $\mathbf{x} \in \mathbb{R}^n$
- Suppose that for functions $g_i \in C^1$, $i = 1, \dots, m$:

$$S := \{ \mathbf{x} \in \mathbb{R}^n \mid g_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \}$$

- Two further cones:

$$\overset{\circ}{G}(\mathbf{x}) := \{ \mathbf{p} \in \mathbb{R}^n \mid \nabla g_i(\mathbf{x})^T \mathbf{p} < 0, i \in \mathcal{I}(\mathbf{x}) \},$$

and



$$G(\mathbf{x}) := \{ \mathbf{p} \in \mathbb{R}^n \mid \nabla g_i(\mathbf{x})^\top \mathbf{p} \leq 0, i \in \mathcal{I}(\mathbf{x}) \}$$

- For every $\mathbf{x} \in \mathbb{R}^n$ it holds that $\overset{\circ}{G}(\mathbf{x}) \subset R_S(\mathbf{x})$, and $T_S(\mathbf{x}) \subset G(\mathbf{x})$
- So, for every $\mathbf{x} \in \mathbb{R}^n$,

$$\overset{\circ}{G}(\mathbf{x}) \subset R_S(\mathbf{x}) \subset \text{cl } R_S(\mathbf{x}) \subset T_S(\mathbf{x}) \subset G(\mathbf{x})$$


Four examples, I

- $S = \{ \mathbf{x} \in \mathbb{R}^2 \mid -x_1 \leq 0, (x_1 - 1)^2 + x_2^2 \leq 1 \}$
- $R_S(\mathbf{0}^2) = \{ \mathbf{p} \in \mathbb{R}^2 \mid p_1 > 0 \}$
- $T_S(\mathbf{0}^2) = \{ \mathbf{p} \in \mathbb{R}^2 \mid p_1 \geq 0 \}$
- $T_S(\mathbf{0}^2) = \text{cl } R_S(\mathbf{0}^2)$

 S  $T_S(\mathbf{0}^2)$

Four examples, II



- $S = \{ \mathbf{x} \in \mathbb{R}^2 \mid -x_1 \leq 0, -x_2 \leq 0, x_1 x_2 \leq 0 \}$
- $R_S(\mathbf{0}^2) = T_S(\mathbf{0}^2) = S$



$$S = T_S(\mathbf{0}^2)$$



Four examples, III

- $S = \{ \mathbf{x} \in \mathbb{R}^2 \mid -x_1^3 + x_2 \leq 0, x_1^5 - x_2 \leq 0, -x_2 \leq 0 \}$
- $R_S(\mathbf{0}^2) = \emptyset$
- $T_S(\mathbf{0}^2) = \{ \mathbf{p} \in \mathbb{R}^2 \mid p_1 \geq 0, p_2 = 0 \}$

 S  $T_S(\mathbf{0}^2)$

Four examples, IV

- $S = \{ \mathbf{x} \in \mathbb{R}^2 \mid -x_2 \leq 0, (x_1 - 1)^2 + x_2^2 = 1 \}$
- $R_S(\mathbf{0}^2) = \emptyset$
- $T_S(\mathbf{0}^2) = \{ \mathbf{p} \in \mathbb{R}^2 \mid p_1 = 0, p_2 \geq 0 \}$



A geometric necessary optimality condition

- $\overset{\circ}{F}(\mathbf{x}^*) := \{ \mathbf{p} \in \mathbb{R}^n \mid \nabla f(\mathbf{x}^*)^T \mathbf{p} < 0 \}$
- Consider the problem (1). If $\mathbf{x}^* \in S$ is a local minimum of f over S then $\overset{\circ}{F}(\mathbf{x}^*) \cap T_S(\mathbf{x}^*) = \emptyset$
- This is an elegant criterion for checking whether a given point is a candidate for a local minimum. There is a catch though:
- The set $T_S(\mathbf{x}^*)$ is nearly impossible to compute in general!
- We will compute other cones that we hope will approximate $T_S(\mathbf{x}^*)$ well enough
- Specifically, we will use the cone $\overset{\circ}{G}(\mathbf{x})$

Example problem

- Consider the differentiable (linear) function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $f(\mathbf{x}) = x_1$
- Then, $\nabla f = (1, 0)^T$, and $\overset{\circ}{F}(\mathbf{0}^2) = \{\mathbf{p} \in \mathbb{R}^2 \mid p_1 < 0\}$
- $\mathbf{x}^* = \mathbf{0}^2$ is a local (in fact, even global) minimum in problem (1) with S given by either one of Examples I–IV
- Easy to check that the geometric necessary optimality condition $\overset{\circ}{F}(\mathbf{0}^2) \cap T_S(\mathbf{0}^2) = \emptyset$ is satisfied in all four examples (no surprise, in view of the above geometric theorem)

The Fritz John conditions

- If $\mathbf{x}^* \in S$ is a local minimum of f over S then there exist multipliers $\mu_0 \in \mathbb{R}$, $\boldsymbol{\mu} \in \mathbb{R}^m$ such that

$$\mu_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m \mu_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}^n, \quad (2a)$$

$$\mu_i g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, m, \quad (2b)$$

$$\mu_0, \mu_i \geq 0, \quad i = 1, \dots, m, \quad (2c)$$

$$(\mu_0, \boldsymbol{\mu}^T)^T \neq \mathbf{0}^{m+1} \quad (2d)$$

- Proof via the geometric necessary conditions and Farkas' Lemma
- What's bad about the Fritz John conditions? It may be possible to fulfill (2) at every feasible point by setting $\mu_0 = 0$! Then, f plays no role, which is bad. We will develop conditions (constraint qualifications) which ensure that $\mu_0 > 0$

Comments

- The vector $\boldsymbol{\mu}$ is a vector of *Lagrange multipliers*. Each of them is associated with a constraint, and will be shown to be a measure of the sensitivity of the solution to changes in the constraints
- Conditions (2a), (2c) are known as the *dual feasibility* conditions
- Condition (2b) is the *complementarity condition*. States that for inactive constraints $i \notin \mathcal{I}(\boldsymbol{x}^*)$, $\mu_i = 0$ must hold
- Will take a closer look at the Examples I–IV, but wait until the KKT conditions have been developed
- We do this by introducing conditions that bring either $\overset{\circ}{G}(\boldsymbol{x})$ or $G(\boldsymbol{x})$ to be tight enough approximations of $T_S(\boldsymbol{x})$

The Karush–Kuhn–Tucker conditions

- *Abadie's CQ:* At $\mathbf{x} \in S$ Abadie's constraint qualification holds if $G(\mathbf{x}) = T_S(\mathbf{x})$
- Satisfied by Example I and IV
- *Assume that at $\mathbf{x}^* \in S$ Abadie's CQ holds. If $\mathbf{x}^* \in S$ is a local minimum of f over S then there exists $\boldsymbol{\mu} \in \mathbb{R}^m$ such that*

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \mu_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}^n, \quad (3a)$$


$$\mu_i g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, m, \quad (3b)$$

$$\boldsymbol{\mu} \geq \mathbf{0}^m \quad (3c)$$

- Proof by first noting that $\overset{\circ}{F}(\mathbf{x}^*) \cap T_S(\mathbf{x}^*) = \emptyset$, which due to our CQ implies that $\overset{\circ}{F}(\mathbf{x}^*) \cap G(\mathbf{x}^*) = \emptyset$. Rest of the proof by Farkas' Lemma. [Note: case of $m = 0$!]

Comments

- The statement in (3a) is that \mathbf{x}^* is a stationary point to the Lagrangian function $\mathbf{x} \mapsto f(\mathbf{x}) + \boldsymbol{\mu}^\top \mathbf{g}(\mathbf{x}) = f(\mathbf{x}) + \sum_{i=1}^m \mu_i g_i(\mathbf{x})$
- The condition (3) is that $-\nabla f(\mathbf{x}^*) \in N_S(\mathbf{x}^*)$ holds. The normal cone $N_S(\mathbf{x}^*)$ is spanned by the normals of the active constraints



Example I

- Abadie's CQ is fulfilled, therefore the KKT-system is solvable
Indeed, the system

$$\left\{ \begin{array}{l} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} -1 & -2 \\ 0 & 0 \end{pmatrix} \boldsymbol{\mu} = \mathbf{0}^2, \\ \boldsymbol{\mu} \geq \mathbf{0}^2, \end{array} \right.$$

possesses solutions $\boldsymbol{\mu} = (\mu_1, 2^{-1}(1 - \mu_1))^T$ for every $0 \leq \mu_1 \leq 1$.
Therefore, there are infinitely many multipliers, that all belong to a bounded set

- Case of a non-unique *dual* solution $\boldsymbol{\mu}$

Equality constraints

Additional constraints $h_j(\mathbf{x}) = 0, j = 1, \dots, \ell$

- KKT system:

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \mu_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^{\ell} \lambda_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}^n, \quad (4a)$$

$$\mu_i g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, m, \quad (4b)$$

$$\boldsymbol{\mu} \geq \mathbf{0}^m \quad (4c)$$

- $\mu_i \geq 0$ for the \leq -constraints; λ_j is sign free for $=$ -constraints
- Interpretation: The condition (4) is a force equilibrium condition
- $-\nabla f(\mathbf{x}^*)$ is a force to violate the active constraints
- The remaining terms equal this force. $\mu_i \geq 0$ must hold (force towards feasibility). λ_j ? Cannot determine before-hand in which direction the surface must move

Other constraint qualifications

- *Slater CQ—convex sets with interior points*: The feasible set is convex, and there exists a feasible point such that every inequality constraint is satisfied strictly
- *Linear independence CQ*: The gradients of all the active constraints are linearly independent
- *Linear constraints CQ*: All the constraints are affine/linear
- *Mangasarian–Fromowitz CQ*: The gradients of all the functions h_j are linearly independent, and the set $\overset{\circ}{G}(\mathbf{x}) \cap H(\mathbf{x})$ is nonempty, where

$$H(\mathbf{x}) := \{ \mathbf{p} \in \mathbb{R}^n \mid \nabla h_i(\mathbf{x})^\top \mathbf{p} = 0, \quad i = 1, \dots, \ell \}$$

- Some CQs are stronger than others: LICQ \implies MFCQ \implies Abadie; Slater \implies MFCQ; linear constraints \implies Abadie

Convexity implies sufficiency

- *Assume the problem (1) is convex, that is, f as well as g_i , $i = 1, \dots, m$, are convex, and h_j , $j = 1, \dots, \ell$, are affine; also, all functions are in C^1 . Assume further that for $\mathbf{x}^* \in S$ the KKT conditions (4) are satisfied. Then, \mathbf{x}^* is a globally optimal solution to the problem (1).*
- *Proof.*

- Check interesting applications in the book on the characterization of eigenvalues and eigenvectors!