

## Exercises in Functional Analysis, Week VI

1. Let  $\{x_n\}$  be a sequence in  $L^p([a, b])$ ,  $x \in L^p([a, b])$ ,  $1 < p < \infty$ . Show that  $x_n \rightarrow x$  weakly iff  $\sup_n \|x_n\| < \infty$  and  $\int_a^\tau x_n(t) dt \rightarrow \int_a^\tau x(t) dt$  for all  $\tau \in [a, b]$ .  
 Consider the function sequence  $\{e^{inx}\}_{n \in \mathbb{N}}$  in  $L^p([-\pi, \pi])$ . For which  $p$ ,  $1 \leq p \leq \infty$ , is it weakly or weakly\* convergent? (For  $L^1$  you consider only weak convergence).
2. Let  $(e_n)$  denote the "standard basis" in  $\ell^p$ . For which  $p$  with  $1 \leq p \leq \infty$  does  $e_n$  converge weakly or weakly\* as  $n \rightarrow \infty$ , and what is the limit? (Be extra careful with the cases  $p = 1$ ,  $p = \infty$ ).
3. We know that  $\ell^1 = c_0^*$  and  $(\ell^1)^* = \ell^\infty$ . Give an example of a sequence of vectors in  $\ell^1$  that converges weakly\* but not weakly, which means that  $f(x_n) \rightarrow f(x)$  for all  $f \in c_0$  but not for all  $f \in \ell^\infty$ .
4. (a) Prove that translation is continuous in  $L^p(\mathbb{R})$  for  $1 \leq p < \infty$ : If  $f_t(x) = f(x - t)$  with  $f \in L^p$  then  $f_t \rightarrow f_{t_0}$  in  $L^p$  as  $t \rightarrow t_0$ . (Hint: use the fact that the space of continuous functions with compact support, i.e. continuous functions which are zero outside of a compact set in  $\mathbb{R}$ , are dense in  $L^p$ ). Prove also that this is false for  $p = \infty$ .  
 (b) Show that in  $L^\infty$  the translation is weakly\* continuous, i.e. show that  $f_t \rightarrow f_{t_0}$  weakly\* in  $L^\infty = (L^1)^*$ .
5. Let  $X$  be a Banach space and  $X^*$  its dual. Assume that  $\{x_j\}$  is a sequence in  $X$  that converges to  $x \in X$ , either in norm or weakly. Similarly,  $\{\varphi_j\}$  is a sequence in  $X^*$  with a limit  $\varphi$ , and the convergence is assumed to be either in norm or weak\*. This gives four different cases of convergence assumption. The question is: does it follow that  $\varphi_j(x_j) \rightarrow \varphi(x)$ ? Answer this question in each of the four cases, by means of a proof or a counterexample. Hint: Use that a sequence that converges weakly or weakly\* is bounded in norm.
6. Let  $H$  be a Hilbert space,  $x, x_n \in H$ ,  $n \geq 1$ . Show that  $x_n \rightarrow x$  in  $H$  if  $x_n \rightarrow x$  weakly and either (a)  $\|x_n\| \rightarrow \|x\|$  or (b)  $\limsup \|x_n\| \leq \|x\|$ .
7. Let  $X$  be a reflexive Banach space and  $\{x_n\} \subset X$ . Prove that if  $\{f(x_n)\}$  is Cauchy for all  $f \in X^*$  then there exists  $x \in X$  such that  $x_n \rightarrow x$  weakly.
8. (An application of Banach-Alaoglu theorem): Any normed space  $X$  can be isometrically embedded into  $C(\Omega)$ , the Banach space of continuous functions on a compact  $\Omega$ . Hint: Take  $\Omega = \{f \in X^* : \|f\| \leq 1\}$  and consider the mapping  $X \ni x \mapsto \hat{x}|_\Omega \in C(\Omega)$ .
9. Let  $X$  be a reflexive Banach space. Show that any bounded sequence of elements in  $X$  has a weakly convergent subsequence. Hint: Use the following facts: (1) If  $X^*$  is

separable then so is  $X$  (in their respective norm topologies); (2) Closed subspaces of reflexive spaces are reflexive.