## MMA320 Introduction to Algebraic Geometry

Exercises for Chapter 6

- **6.1.** Suppose a hypersurface V=(f) of degree d>1 in  $\mathbb{P}^n$  contains a linear subspace of dimension  $r\geq n/2$ . Show that V is singular.
- **6.2**. Prove that there is at least one line through a singular point of a cubic surface (and 'in general' 6). Let  $X \subset \mathbb{P}^4$  be a non-singular cubic threefold. Show that through every point there is at least one line.
- **6.3**. Show that the surface  $V(X^n + Y^n + Z^n + T^n) \subset \mathbb{P}^3$  contains exactly  $3n^2$  lines  $(\operatorname{char} k = 0 \text{ and } k = \bar{k})$ .
- **6.4**. Find all 27 lines on Clebsch' diagonal surface:

$$X_0^3 + X_1^3 + X_2^3 + X_3^3 + X_4^3 = 0$$
  
$$X_0 + X_1 + X_2 + X_3 + X_4 = 0$$

All lines are real, 15 are easy to see as intersections with coordinate hyperplanes.

**6.5**. Find the singular points of the cubic surface

$$X_0 X_1 X_2 + X_0 X_1 X_3 + X_0 X_2 X_3 + X_1 X_2 X_3 = 0.$$

Determine all lines on the surface.

**6.6**. Find the singular point of the cubic surface

$$ZX^2 - Y^3 + TZ^2 = 0 .$$

Determine all lines on the surface.

**6.7**. Determine the number of lines on the singular cubic surface

$$XYZ - T^3 = 0.$$

- **6.8**. Let  $P_1, \ldots, P_6$  be 6 points on a nondegenerate conic  $C = (f) \subset \mathbb{P}^2$ . The linear system  $\mathbb{P}(S_3(P_1, \ldots, P_6))$  of cubics through these six points gives rise to a rational map  $\mathbb{P}^2 \dashrightarrow \mathbb{P}^3$ . Compute the equation of the image.
- **6.9**. Let  $\pi$  be the skew projection of a smooth cubic surface from two skew lines l and m on a plane through the transversal  $l_1$ . Show that  $\pi$  is a regular map and determine which lines are blown down.
- **6.10**. Let S be the cubic surface, which is the image of the rational map determined by the linear system of cubics in the plane through 6 points, no three on a line, not all on a conic. Describe the 27 lines in terms of points and curves in the plane.