Brief solutions to examination in algebra 2017-06-08⁻ (MMG 500-MVE 150)

1a).
$$\pi = (12)(23)(34) = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 4 & 3 \\ 1 & 3 & 4 & 2 \\ 2 & 3 & 4 & 1 \end{pmatrix} = (1234)$$

1b) $(1234)^2 = (13)(24)$, $(1234)^3 = (1432)$ and $(1234)^4 = id$.

The group generated by π will thus have {*id*, (1234), (13)(24), (1432)} as underlying set.

2 If $k \in \mathbb{N}$, then $h(gh)^k h^{-1} = (hg)^k hh^{-1}$ by the associative law. As $hh^{-1} = e$, we thus get $h(gh)^k h^{-1} = (hg)^k$. In particular, if $(gh)^k = e$, then $(hg)^k = heh^{-1} = e$. Conversely, by symmetry $(hg)^k = e \Longrightarrow (gh)^k = e$. The list of exponents of all $k \in \mathbb{N}$ with $(gh)^k = e$ will therefore coincide with the list of all exponents with $(hg)^k = e$. So gh and hg have the same order.

3. Let [*k*] be the congruence class (mod 10) of $k \in \mathbb{Z}$. Then, [2], [4], [5], [6] and [8] are zero divisors in \mathbb{Z}_{10} as [2] [5] = [4][5] = [6][5] = [8][5] = [0], while [1], [3], [7], [9] are invertible as $[1]^2 = [1]$, [3][7] = [1] and $[9]^2 = [1]$. So any element $[k] \neq [0]$ is either a zero divisor or invertible in \mathbb{Z}_{10} .

Further, no element [k] in \mathbb{Z}_{10} can be both a zero divisor and invertible, Indeed, if [j][k]=[0] and [k][l]=[1], then [j]=[j][1]=[j]([k]][l]=([j][k])[l]=[0][l]=[0]. There are thus no further zero divisors or invertible elements in \mathbb{Z}_{10} .

4 The elements $\neq 0$ in \mathbb{Z}_p form a multiplicative group with p-1 elements as \mathbb{Z}_p is a field. By a corollary of Lagrange's theorem we have thus that $x^{p-1}=1$ for all $x\neq 0$. in \mathbb{Z}_p . Hence $x^p = x$ for all $x \in \mathbb{Z}_p$. The equation $x^p - a = 0$ is thus equivalent to the equation x - a = 0. This means that for each $a \in \mathbb{Z}_p$, the equation $x^p - a = 0$ has exactly one solution in \mathbb{Z}_p , namely x = a.

- 5. See Durbin's book
- 6. See Durbin's book