

CHALMERS | GÖTEBORGS UNIVERSITET

PREPRINT

Bicoloured Dyck Paths, Segmented
Permutations, and Chebyshev
Polynomials

ANDERS CLAEISSON

Department of Mathematics
CHALMERS UNIVERSITY OF TECHNOLOGY
GÖTEBORG UNIVERSITY
Göteborg Sweden 2004

Preprint 2004:32

Bicoloured Dyck Paths, Segmented Permutations, and Chebyshev Polynomials

Anders Claesson

CHALMERS | GÖTEBORGS UNIVERSITET



Mathematics
Department of Mathematics
Chalmers University of Technology and Göteborg University
SE-412 96 Göteborg, Sweden
Göteborg, May 2004

NO 2004:32
ISSN 0347-2809

Matematiska Vetenskaper
Göteborg 2004

BICOLOURED DYCK PATHS, SEGMENTED PERMUTATIONS, AND CHEBYSHEV POLYNOMIALS

ANDERS CLAESSON

ABSTRACT. A bicoloured Dyck path is a Dyck path in which each up-step is assigned one of two colours, say, red and green. We say that a permutation π is σ -segmented if every occurrence o of σ in π is a segment-occurrence (i.e., o is a contiguous subword in π).

We show combinatorially the following results: The 132-segmented permutations of length n with k occurrences of 132 are in one-to-one correspondence with bicoloured Dyck paths of length $2n - 4k$ with k red up-steps. Similarly, the 123-segmented permutations of length n with k occurrences of 123 are in one-to-one correspondence with bicoloured Dyck paths of length $2n - 4k$ with k red up-steps, each of height less than 2.

We enumerate the permutations above by enumerating the corresponding bicoloured Dyck paths. More generally, we present a bivariate generating function for the number of bicoloured Dyck paths of length $2n$ with k red up-steps, each of height less than h . This generating function is expressed in terms of Chebyshev polynomials of the second kind.

1. INTRODUCTION

It is relatively straightforward to show that number of permutations of $[n] = \{1, 2, \dots, n\}$ avoiding a pattern of length 3 is the Catalan number, $C_n = \binom{2n}{n}/(n+1)$ (e.g., see [8] or [5]). In contrast, to count the permutations containing r occurrences of a fixed pattern of length 3, for a general r , is a very hard problem. The best result on this latter problem has been achieved by Mansour and Vainshtein [6]. They presented an algorithm that computes the generating function for the number of permutations with r occurrences of 132 for any $r \geq 0$. The algorithm has been implemented in C. It yields explicit results for $1 \leq r \leq 6$.

We say that an occurrence o of σ in π is a *segment-occurrence* if o is a segment (also called factor) of π , in other words, if o is a contiguous subword in π . Elizalde and Noy [2] presented exponential generating functions for the distribution of the number of segment-occurrences of any pattern of length 3. Related problems have also been studied by Kitaev [3] and by Kitaev and Mansour [4].

We say that π is σ -segmented if every occurrence of σ in π is a segment-occurrence. For instance, 4365172 contains 3 occurrences of 132, namely 465, 365, and 172. Of these occurrences, only 365 and 172 are segment-occurrences. Thus 4365172 is not 132-segmented. Note that if π is σ -avoiding then π is also σ -segmented. In this article we try to enumerate the σ -segmented permutations by length and by the number of occurrences of σ . In [5] Krattenthaler gave two bijections: one between 132-avoiding permutations and Dyck paths, and one between 123-avoiding permutations and Dyck paths. We obtain two new results by extending these bijections:

- The 132-segmented permutations of length n with k occurrences of 132 are in one-to-one correspondence with bicoloured Dyck paths of length $2n - 4k$ with k red up-steps.

Date: April 16, 2004.

- The 123-segmented permutations of length n with k occurrences of 123 are in one-to-one correspondence with bicoloured Dyck paths of length $2n - 4k$ with k red up-steps, each of height less than 2.

Here a bicoloured Dyck path is a Dyck path in which each up-step is assigned one of two colours, say, red and green. We enumerate the permutations above by enumerating the corresponding bicoloured Dyck paths. To be more precise, let $\mathcal{B}_{n,k}$ be the set of bicoloured Dyck path of length $2n$ with k red up-steps. Let $\mathcal{B}_{n,k}^{[h]}$ be the subset of $\mathcal{B}_{n,k}$ consisting of those paths where the height of each red up-step is less than h . It is plain that $|\mathcal{B}_{n,k}| = \binom{n}{k} C_n$. We show that


$$\sum_{n,k \geq 0} |\mathcal{B}_{n,k}^{[h]}| q^{k_t n} = \frac{C(t) - 2xqU_h(x)U_{h-1}(x)}{1 + q - qU_h^2(x)}, \quad x = \frac{1}{2\sqrt{(1+q)t}},$$

where $C(t) = (1 - \sqrt{1 - 4t})/(2t)$ is the generating function for the Catalan numbers, and U_n is the n th Chebyshev polynomial of the second kind. We also find formulas for $|\mathcal{B}_{n,k}^{[1]}|$ and $|\mathcal{B}_{n,k}^{[2]}|$.

2. BICOLOURED DYCK PATHS

By a *lattice path* we shall mean a path in \mathbb{Z}^2 with steps $(1, 1)$ and $(1, -1)$; the steps $(1, 1)$ and $(1, -1)$ will be called *up-* and *down-steps*, respectively. Furthermore, a lattice path that never falls below the x -axis will be called *nonnegative*.

Recall that a *Dyck path* of length $2n$ is a nonnegative lattice path from $(0, 0)$ to $(2n, 0)$. As an example, these are the 5 Dyck paths of length 6:



Letting u and d represent the steps $(1, 1)$ and $(1, -1)$, we code a Dyck path with a word over $\{u, d\}$. For example, the paths above are coded by

$$ududud \quad uduudd \quad uuddud \quad uuddud \quad uuuddd$$

Let \mathcal{D}_n be the language over $\{u, d\}$ obtained from Dyck paths of length $2n$ via this coding, and let $\mathcal{D} = \cup_{n \geq 0} \mathcal{D}_n$. In general, if \mathcal{A} is a language over some alphabet X , then the *characteristic series* of \mathcal{A} , also (by slight abuse of notation) denoted \mathcal{A} , is the element of $C\langle\langle X \rangle\rangle$ defined by

$$\mathcal{A} = \sum_{w \in \mathcal{A}} w.$$

A nonempty Dyck path β can be written uniquely as $u\beta_1 d\beta_2$ where β_1 and β_2 are Dyck paths. This decomposition is called the *first return decomposition* of β , because the d in $u\beta_1 d\beta_2$ corresponds to the first place, after $(0, 0)$, where the path touches the x -axis. By this decomposition, the characteristic series of \mathcal{D} is uniquely determined by the functional equation

$$\mathcal{D} = 1 + u\mathcal{D}d\mathcal{D}, \quad (1)$$

where 1 denotes the empty word.


In a similar vein, we now consider the language \mathcal{B} over $\{u, \bar{u}, d\}$ whose characteristic series is uniquely determined by the functional equation

$$\mathcal{B} = 1 + (u + \bar{u})\mathcal{B}d\mathcal{B}. \quad (2)$$

Let \mathcal{B}_n be the set of words in \mathcal{B} that are of length $2n$, and let $\mathcal{B}_{n,k}$ be the set of words in \mathcal{B}_n with k occurrences of \bar{u} . As an example, when $n = 3$ and $k = 1$ there are 15 such words, namely

$\bar{u}dudud \quad \bar{u}duudd \quad \bar{u}uddud \quad \bar{u}uddd \quad \bar{u}uuddd$
 $ud\bar{u}dud \quad ud\bar{u}udd \quad u\bar{u}ddud \quad u\bar{u}dudd \quad u\bar{u}uddd$
 $udud\bar{u}d \quad ud\bar{u}udd \quad uudd\bar{u}d \quad uudd\bar{u}d \quad uu\bar{u}ddd$

We may view the elements of \mathcal{B} as *bicoloured Dyck paths*. The words from the previous example are depicted below.



Here steps represented by double edges are, say, red, and steps represented by simple edges are, say, green.

Proposition 1. *With $C_n = \text{card } \mathcal{D}_n$, we have*

$$\text{card } \mathcal{B}_{n,k} = \binom{n}{k} C_n \quad \text{and} \quad \text{card } \mathcal{B}_n = 2^n C_n.$$

Proof. A bicoloured Dyck paths β of length $2n$ naturally breaks up into two parts: (a) The Dyck path obtained from β by replacing each red up-step with a green ditto. (b) The subset of $[n] = \{1, 2, \dots, n\}$ consisting of those integers i for which the i th up-step is red. \square

For $h \geq 1$, let $\mathcal{B}^{[h]}$ be the subset of \mathcal{B} whose characteristic series is the solution to

$$\mathcal{B}^{[h]} = 1 + (u + \bar{u})\mathcal{B}^{[h-1]}d\mathcal{B}^{[h]}, \quad (3)$$

with the initial condition $\mathcal{B}^{[0]} = \mathcal{D}$, where \mathcal{D} is defined as above. Let

$\mathcal{B}_n^{[h]}$ be the set of words in $\mathcal{B}^{[h]}$ that are of length $2n$, and let

$\mathcal{B}_{n,k}^{[h]}$ be the set of words in $\mathcal{B}_n^{[h]}$ with k occurrences of \bar{u} .

To translate these definitions in terms of lattice paths we define the *height* of a step in a (bicoloured) lattice path as the height above the x -axis of its left point. Then $\mathcal{B}^{[h]}$ is the set of bicoloured Dyck paths whose red up-steps all are of height less than h . As an example, there is exactly one element in $\mathcal{B}_{3,1}$ that is not in $\mathcal{B}^{[2]}$, namely



To count words of given length in \mathcal{D} , \mathcal{B} and $\mathcal{B}^{[h]}$, we will study the commutative counterparts of the functional equations (1), (2) and (3). Formally, we define the substitution $\mu : \mathbb{C}\langle\langle u, \bar{u}, d \rangle\rangle \rightarrow \mathbb{C}\llbracket q, t \rrbracket$ by

$$\mu = \{ u \mapsto 1, \bar{u} \mapsto q, d \mapsto t \}.$$

Let $C = \mu(\mathcal{D})$, $B = \mu(\mathcal{B})$, and $B^{[h]} = \mu(\mathcal{B}^{[h]})$. We then get

$$C = 1 + tC^2, \quad (4)$$

$$B = 1 + (1 + q)tB^2, \quad (5)$$

$$B^{[h]} = 1 + (1 + q)tB^{[h-1]}B^{[h]}, \quad B^{[0]} = C. \quad (6)$$

By an easy application of the Lagrange inversion formula it follows from (4) that

$$[t^n]C(t)^i = \frac{i}{i+n} \binom{2n+i-1}{n}. \quad (7)$$

In particular, we obtain that C is the familiar generating function of the Catalan numbers, $C_n = \frac{1}{n+1} \binom{2n}{n}$. Thus we have derived the well known fact that the number of Dyck paths of length $2n$ is the n th Catalan number. Furthermore, it follows from (5) that

$$B(q, t) = C((1 + q)t), \quad (8)$$

and it follows from (6) that

$$B^{[h]}(q, t) = \frac{1}{1 - (1 + q)tB^{[h-1]}C(t)}, \quad B^{[0]} = C. \quad (9)$$

From these series we generate the first few values of $|\mathcal{B}_{n,k}|$, $|\mathcal{B}_{n,k}^{[1]}|$ and $|\mathcal{B}_{n,k}^{[2]}|$; tables with these values are given in Section 5.

Recall that the *Chebyshev polynomials of the second kind*, denoted $U_n(x)$, are defined by

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta},$$

where n is an integer, $x = \cos \theta$, and $0 \leq \theta \leq \pi$. Equivalently, these polynomials can be defined as the solution to the linear difference equation

$$U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x),$$

with $U_{-1}(x) = 0$ and $U_0(x) = 1$.

Via a bijection between Dyck paths and 132-avoiding permutations due to Krattenthaler [5, Lemma Φ and Theorem 2] it follows by a result of Chow and West [1, Theorem 3.1] that

$$C^{[h]}(t) = \frac{U_h\left(\frac{1}{2\sqrt{t}}\right)}{\sqrt{t} \cdot U_{h+1}\left(\frac{1}{2\sqrt{t}}\right)} \quad (10)$$

is the generating function for Dyck paths that stay below height h . Note that, since $C^{[0]} = 1$ and $C^{[h]} = (1 - tC^{[h-1]})^{-1}$, this result is also easy to prove by induction on h .

Theorem 2. *With $B^{[h]}$ being the generating function for the number of Dyck paths whose red up-steps all are of height less than h , and U_n being the n th Chebyshev polynomial of the second kind we have*

$$B^{[h]}(q, t) = \frac{4x^2U_{h-1}(x) - 2xU_{h-2}(x)C(t)}{2xU_h(x) - U_{h-1}(x)C(t)} = \frac{C(t) - 2xqU_h(x)U_{h-1}(x)}{1 + q - qU_h^2(x)},$$

where $x = 1/(2\sqrt{(1+q)t})$, and $C(t) = (1 - \sqrt{1 - 4t})/(2t)$ is the generating function for the Catalan numbers.

Proof. We shall prove the first equality by induction. To this end, we let

$$F^{[h]}(q, t) = \frac{4x^2U_{h-1}(x) - 2xU_{h-2}(x)C(t)}{2xU_h(x) - U_{h-1}(x)C(t)}.$$

From $U_{-2}(x) = -1$, $U_{-1}(x) = 0$, and $U_0(x) = 1$ it readily follows that $F^{[0]}(q, t) = C(t) = B^{[0]}(q, t)$. If $B^{[h]} = F^{[h]}$, for some fixed $h \geq 0$, then

$$\begin{aligned} B^{[h+1]} &= \frac{1}{1 - (1+q)tB^{[h]}} \\ &= \frac{1}{1 - (1+q)tF^{[h]}} \\ &= \frac{2xU_h - U_{h-1}C}{2xU_h - U_{h-1}C - (1+q)t(4x^2U_{h-1} - 2xU_{h-2}C)} \\ &= \frac{2xU_h - U_{h-1}C}{2xU_h - (1+q)t4x^2U_{h-1} - (U_{h-1} - (1+q)t2xU_{h-2})C} \\ &= \frac{4x^2U_h - 2xU_{h-1}C}{2x(2xU_h - (1+q)t4x^2U_{h-1}) - (2xU_{h-1} - (1+q)t4x^2U_{h-2})C} \\ &= \frac{4x^2U_h - 2xU_{h-1}C}{2x(2xU_h - U_{h-1}) - (2xU_{h-1} - U_{h-2})C} \\ &= \frac{4x^2U_h - 2xU_{h-1}C}{2xU_{h+1} - U_hC} \\ &= F^{[h+1]}. \end{aligned}$$

This completes the induction step, and thus the first equality holds for all $h \geq 0$. The second equality is plain algebra/trigonometry. \square

Proposition 3. For $n, k \geq 0$ we have

$$\begin{aligned} \text{card } \mathcal{B}_{n,k}^{[1]} &= b(n+k, n-k) = \frac{2k+1}{n+k+1} \binom{2n}{n-k}, \\ \text{card } \mathcal{B}_n^{[1]} &= \binom{2n}{n}, \end{aligned}$$

where $b(n, k) = \frac{n-k+1}{n+1} \binom{n+k}{n}$ is a ballot number.

Proof. The ballot number $b(n, k)$ is the number of nonnegative lattice paths from $(0, 0)$ to $(n+k, n-k)$. Thus, the first claim of the lemma is that $|\mathcal{B}_{n,k}^{[1]}|$ equals the number of nonnegative lattice paths from $(0, 0)$ to $(2n, 2k)$; let $\mathcal{A}_{n,k}$ denote the language over $\{u, d\}$ obtained from these paths via the usual coding. In addition, let $\mathcal{A}_n = \cup_{k \geq 0} \mathcal{A}_{n,k}$ and $\mathcal{A} = \cup_{n \geq 0} \mathcal{A}_n$. The characteristic series of \mathcal{A} satisfies

$$\mathcal{A} = 1 + u\mathcal{D}(u+d)\mathcal{A}.$$

From (3) we also know that

$$\mathcal{B}^{[1]} = 1 + (u + \bar{u})\mathcal{D}d\mathcal{B}^{[1]}.$$

We exploit the obvious similarity between these two functional equations to define, by recursion, a length preserving bijection f from $\mathcal{B}^{[1]}$ onto \mathcal{A} such that $\beta \in \mathcal{B}^{[1]}$ has exactly k occurrences of \bar{u} precisely when $f(\beta) \in \mathcal{A}$ ends at height $2k$:

$$f(\beta) = \begin{cases} 1 & \text{if } \beta = 1, \\ u\beta_1df(\beta_2) & \text{if } \beta = u\beta_1d\beta_2, \beta_1 \in \mathcal{D}, \beta_2 \in \mathcal{B}^{[1]}, \\ u\beta_1uf(\beta_2) & \text{if } \beta = \bar{u}\beta_1d\beta_2, \beta_1 \in \mathcal{D}, \beta_2 \in \mathcal{B}^{[1]}. \end{cases}$$

For $\beta \in \mathcal{B}$, let $|\beta|_{\bar{u}}$ denote the number of occurrences of \bar{u} in β , and for $\alpha \in \mathcal{A}$ let $h(\alpha)$ denote the height at which α ends. To prove that f is length preserving, bijective, and that $2|\cdot|_{\bar{u}} = h \circ f$, we use induction on path-length: f trivially has these properties as a function from $\mathcal{B}_0^{[1]}$ to \mathcal{A}_0 . Let n be a positive integer and assume that f has the desired properties as a function from $\cup_{k=0}^{n-1} \mathcal{B}_k^{[1]}$ to $\cup_{k=0}^{n-1} \mathcal{A}_k$.

Any β in $\mathcal{B}_n^{[1]}$ can be written as $\beta = x\beta_1d\beta_2$ for some $x \in \{u, \bar{u}\}$, $\beta_1 \in \mathcal{D}$ and $\beta_2 \in \mathcal{B}^{[1]}$. Therefore,

$$|f(\beta)| = 2 + |\beta_1| + |f(\beta_2)| = 2 + |\beta_1| + |\beta_2| = |\beta|$$

and

$$(h \circ f)(\beta) = 2|x|_{\bar{u}} + (h \circ f)(\beta_2) = 2|x|_{\bar{u}} + 2|\beta_2|_{\bar{u}} = 2|\beta|_{\bar{u}}$$

To prove that f is injective, assume that $f(\beta) = f(\beta')$, where $\beta' = x'\beta'_1d\beta'_2$ for some $x' \in \{u, \bar{u}\}$, $\beta'_1 \in \mathcal{D}$, and $\beta'_2 \in \mathcal{B}^{[1]}$. Then

$$f(\beta) = u\beta_1yf(\beta_2) = u\beta'_1y'f(\beta'_2) = f(\beta'),$$

in which $y, y' \in \{u, d\}$. Thus $\beta_1 = \beta'_1$, $y = y'$, and $f(\beta_2) = f(\beta'_2)$. By the induction hypothesis, $f(\beta_2) = f(\beta'_2)$ implies that $\beta_2 = \beta'_2$, and hence $\beta = \beta'$.

To prove that f is surjective, take any $\alpha = u\alpha'y\alpha''$ in \mathcal{A}_n , where $y \in \{u, d\}$, $\alpha' \in \mathcal{D}$, and $\alpha'' \in \mathcal{A}$. By the induction hypothesis, there exists β'' in $\mathcal{B}^{[1]}$ such that $f(\alpha'') = \beta''$; so $f(u\alpha'y\beta'') = \alpha$. This concludes the proof of the first part of the lemma.

Given the first result, the second result may be formulated as saying that the central binomial coefficient $\binom{2n}{n}$ is the sum of the ballot numbers $b(n+k, n-k)$ for $k = 0, 1, \dots, n$. This is a known fact (see [7, p. 79]); indeed

$$\frac{2k+1}{n+k+1} \binom{2n}{n-k} = \binom{2n}{n-k} - \binom{2n}{n-k-1},$$

and hence the sum these numbers is telescoping.

For a bijective proof of the second part we consider the set of all lattice paths from $(0, 0)$ to $(2n, 0)$; let \mathcal{P}_n be the language over $\{u, d\}$ obtained from these $\binom{2n}{n}$ paths via the usual coding, and let $\mathcal{P} = \cup_{n \geq 0} \mathcal{P}_n$. The characteristic series of \mathcal{P} satisfies


$$\mathcal{P} = 1 + u\mathcal{D}d\mathcal{P} + d\widehat{\mathcal{D}}u\mathcal{P},$$

where $\widehat{\mathcal{D}}$ is the image of \mathcal{D} under the involution on $\mathbb{C}\langle\langle u, d \rangle\rangle$ defined by $u \mapsto d$ and $d \mapsto u$; this involution has the effect of reflecting a Dyck path in the x -axis. A length preserving bijection g from $\mathcal{B}^{[1]}$ onto \mathcal{P} is then recursively defined by


$$g(\beta) = \begin{cases} 1 & \text{if } \beta = 1, \\ u\beta_1dg(\beta_2) & \text{if } \beta = u\beta_1d\beta_2, \beta_1 \in \mathcal{D}, \beta_2 \in \mathcal{B}^{[1]}, \\ d\widehat{\beta}_1ug(\beta_2) & \text{if } \beta = \bar{u}\beta_1d\beta_2, \beta_1 \in \mathcal{D}, \beta_2 \in \mathcal{B}^{[1]}. \end{cases}$$

Again, by induction on path-length it follows that g is a bijection. \square

Example. As an illustration of the bijections in the proof of Proposition 3, we have



and



Proposition 4. For $n, k \geq 0$ we have

$$\text{card } \mathcal{B}_{n,k}^{[2]} = \sum_{i \geq 0} \frac{2k+i+1}{n+k+i+1} \binom{k-1}{k-i} \binom{2n+i}{n-k}.$$

Proof. From (9) it follows that

$$B^{[2]}(q, t) = \frac{1 - t(1+q)C(t)}{1 - t(1+q)(1+C(t))}.$$

Using (4) we rewrite this as

$$B^{[2]}(q, t) = \frac{(1 - qtC(t)^2)C(t)}{1 - (1+C(t))qtC(t)^2}, \quad (11)$$

and on expanding the right hand side as a geometric series we get

$$[q^k]B^{[2]}(q, t) = t^k C(t)^{2k+1} (1+C(t))^{k-1} (\delta_{k,0} + C(t)), \quad (12)$$

where $\delta_{k,0}$ is 1 if $k=0$, and 0 otherwise. The result is easy to check for $k=0$, so let us assume that $k \geq 1$. Then

$$[q^k]B^{[2]}(q, t) = t^k \sum_{i \geq 0} \binom{k-1}{i} C(t)^{3k-i+1} = t^k \sum_{i \geq 0} \binom{k-1}{3k-i} C(t)^{i+1}.$$

From (7) we get

$$\begin{aligned} [t^n q^k]B^{[2]}(q, t) &= \sum_{i \geq 0} \frac{i+1}{n-k+i+1} \binom{k-1}{3k-i} \binom{2n-2k+i}{n-k} \\ &= \sum_{i \geq 0} \frac{i+1}{n+k+i+1} \binom{k-1}{i-1} \binom{2n-i}{n-k}, \end{aligned}$$

which concludes the proof. \square

3. SEGMENTED PERMUTATIONS

Let $v = v_1 v_2 \cdots v_n$ be a word over \mathbb{N} without repeated letters. We define the *reduction* of v , denoted $\text{red}(v)$, by

$$\text{red}(v)(i) = \text{card}\{j : v_j \leq v_i\}.$$

In other words, $\text{red}(v)$ is the permutation of $[n]$ obtained from v by replacing the smallest letter in v with 1, the second smallest with 2, etc. For instance, $\text{red}(19453) = 15342$. We will also need a map that is a kind of inverse to red . For a finite subset V of \mathbb{N} , with $n = |V|$, and a permutation π of $[n]$, we denote by $\text{red}_V^{-1}(\pi)$ the word over V obtained from π by replacing i in π with the i th smallest element in V , for all i . Here is an example: If $V = \{1, 3, 4, 5, 9\}$ then $\text{red}_V^{-1}(15342) = 19453$.

Given π in \mathcal{S}_n and σ in \mathcal{S}_k , an *occurrence* of σ in π is a subword

$$o = \pi(i_1)\pi(i_2) \cdots \pi(i_k)$$

of π such that $\text{red}(o) = \sigma$. If, in addition, $i_r + 1 = i_{r+1}$ for each $r = 1, 2, \dots, k-1$, then o is a *segment-occurrence* of σ in π . We say that π is $(\sigma)^k$ -*segmented* if there are exactly k occurrences of σ in π , each of which is a segment-occurrence of σ in π . A $(\sigma)^0$ -segmented permutation is usually called σ -*avoiding*, and the set of σ -avoiding permutations of $[n]$ is denoted $\mathcal{S}_n(\sigma)$.

If π is $(\sigma)^k$ -segmented for some k , then we say that π is σ -*segmented*. We also define

$$\mathcal{R}_n^k(\sigma) = \{\pi \in \mathcal{S}_n : \pi \text{ is } (\sigma)^k\text{-segmented}\}$$

and $\mathcal{R}_n(\sigma) = \cup_{k \geq 0} \mathcal{R}_n^k(\sigma)$. In other words, $\mathcal{R}_n(\sigma)$ is the set of σ -segmented permutations of length n . Let

$$R(\sigma; q, t) = \sum_{k, n \geq 0} \text{card } \mathcal{R}_n^k(\sigma) q^k t^n.$$

The first nontrivial case is $\sigma = 12$. A permutation is 12-segmented if all its non-inversions are rises. For instance, the permutation 7653412 is 12-segmented while 7643512 is not (45 is a non-inversion, but not a rise).

Let $\pi \in \mathcal{R}_n(12)$ with $n \geq 1$. If the letter 1 precedes the letter b in π , then $1b$ is an occurrence of 12 in π . Thus, either 1 is the last letter in π , or 1 is the penultimate letter in π and 2 is the last letter in π . In terms of the generating function $R = R(12; q, t)$ this amounts to

$$R = 1 + tR + qt^2R.$$

So R is a rational function in t and q . Extracting coefficients we get

$$\text{card } \mathcal{R}_n^k(12) = \binom{n-k}{k} \quad \text{and} \quad \text{card } \mathcal{R}_n(12) = F_n,$$

where F_n is the n th Fibonacci number (i.e., $F_{n+1} = F_n + F_{n-1}$ with $F_0 = F_1 = 1$).

For the cases $\sigma = 123$ and $\sigma = 132$, we have the following result.

Theorem 5. *Let $k \geq 0$ and $n \geq 3k$.*

The 132-segmented permutations of length n with k occurrences of 132 are in one-to-one correspondence with bicoloured Dyck paths of length $2n - 4k$ with k red up-steps. Thus

$$\text{card } \mathcal{R}_n^k(132) = \text{card } \mathcal{B}_{n-2k, k} = \binom{n-2k}{k} C_{n-2k},$$

where the last equality is a consequence of Proposition 1.

The 123-segmented permutations of length n with k occurrences of 123 are in one-to-one correspondence with bicoloured Dyck paths of length $2n - 4k$ with k red up-steps, each of height less than 2. Thus

$$\text{card } \mathcal{R}_n^k(123) = \text{card } \mathcal{B}_{n-2k, k}^{[2]} = \sum_{i \geq 0} \frac{2k+i+1}{n-k+i+1} \binom{k-1}{k-i} \binom{2n-4k+i}{n-3k},$$

where the last equality is a consequence of Proposition 4.

First proof. Let n be a positive integer, and let π be a 132-segmented permutation of length n . If the letter n is not part of any occurrence of 132, then we can factor π as $\pi = \pi_1 n \pi_2$, where π_1 and π_2 are 132-segmented permutations, and $\pi_2 < \pi_1$ (i.e., every letter in π_2 is smaller than every letter in π_1). On the other hand, if n is part of an occurrence of 132, then we can factor π as

$$\pi = \pi_1 a n b \pi_2, \quad \text{where } \pi_2 < a < b < \pi_1,$$

and π_1 and π_2 are 132-segmented permutations. In particular, $a = |\pi_2| + 1$ and $b = a + 1$. Thus the generating function $R = R(132; q, t)$ satisfies the functional equation

$$R = 1 + (t + qt^3)R^2.$$

It follows that $R = C(t + qt^3)$, where $C(t)$ is the generating function for the Catalan numbers, and hence $[t^n q^k]R = \text{card } \mathcal{B}_{n-2k, k}$, as claimed.

Let $\pi \in \mathcal{R}_n^k(123)$ with $n \geq 1$. Then, either $k=0$ and π is 123-avoiding, or $k \geq 1$ and π contains at least one occurrence of 123. Let us focus on the latter case, and let

$$\pi = \pi_1 abc \pi_2,$$

where abc is the leftmost occurrence of 123 in π . Then $a\pi_2$ is $(123)^{k-1}$ -segmented and $\pi_1 c$ is 123-avoiding, with the additional restriction that $a\pi_2$ may not begin with an occurrence of 123. Moreover,

$$a\pi_2 < b < \pi_1 c,$$

or else a non segment-occurrence of 123 would be present. With regard to the generating function $R = R(123; q, t)$ this decomposition of 123-segmented permutations amounts to the functional equation

$$R = C + qt(C - 1)(\tilde{R} - 1), \quad (13)$$

where $C = C(t)$ is the generating function of the Catalan numbers, and the coefficient of $q^k t^n$ in $\tilde{R} = \tilde{R}(q, t)$ is the number of $(123)^k$ -segmented permutations of length n that do not begin with an occurrence of 123. Considering the decomposition above in the special case when π_1 is the empty word, we see that $t^2 q(\tilde{R} - 1)$ is the generating function of the number of 123-segmented permutations that begin with an occurrence of 123; so

$$R = \tilde{R} + qt^2(\tilde{R} - 1). \quad (14)$$

Solving equations (13) and (14) for R , eliminating \tilde{R} , we get

$$R = \frac{(1 - qt^3 C^2)C}{1 - (1 + C)qt^3 C^2}. \quad (15)$$

It follows from (11) that $R = B^{[2]}(qt^2, t)$, as claimed. \square

Second proof. We shall define a bijection

$$f : \mathcal{R}(132) \rightarrow \mathcal{B}^{[2]},$$

such that $|f(\pi)| = 2(n - 2k)$ and $|f(\pi)|_{\bar{u}} = k$ whenever $\pi \in \mathcal{R}_n^k(132)$. Our definition of f will be recursive and we start by defining that $f(\epsilon) = \epsilon$, where ϵ denotes the empty word. Now, assume that n is a positive integer, and let π be a 132-segmented permutation of length n . As in the first proof, if the letter n is not part of any occurrence of 132, then we can factor π as $\pi = \pi_1 n \pi_2$, where π_1 and π_2 are 132-segmented permutations, and $\pi_2 < \pi_1$; in this case we define

$$f(\pi) = u(f \circ \text{red})(\pi_1)d(f \circ \text{red})(\pi_2).$$

If n is part of an occurrence of 132, then we can factor π as $\pi = \pi_1 a n b \pi_2$ where $\pi_2 < a < b < \pi_1$ and π_1 and π_2 are 132-segmented permutations; in this case we define

$$f(\pi) = \bar{u}(f \circ \text{red})(\pi_1)d(f \circ \text{red})(\pi_2).$$

For any β in \mathcal{B} , let

$$\lambda(\beta) = \frac{1}{2}|\beta| + 2|\beta|_{\bar{u}} = |\beta|_u + 3|\beta|_{\bar{u}}.$$

Using induction, it is plain to show that the inverse of f is given by

$$\begin{aligned} f^{-1}(\epsilon) &= \epsilon; \\ f^{-1}(u\beta_1 d\beta_2) &= (\text{red}_{V_1}^{-1} \circ f^{-1})(\beta_1) n (\text{red}_{V_2}^{-1} \circ f^{-1})(\beta_2), \end{aligned}$$

where $n = \lambda(\beta_1) + \lambda(\beta_2) + 1$, $V_1 = [\lambda(\beta_2) + 1, n - 1]$, and $V_2 = [1, \lambda(\beta_2)]$;

$$f^{-1}(\bar{u}\beta_1 d\beta_2) = (\text{red}_{V_1}^{-1} \circ f^{-1})(\beta_1) a n b (\text{red}_{V_2}^{-1} \circ f^{-1})(\beta_2),$$

where $a = \lambda(\beta_2) + 1$, $b = a + 1$, $n = \lambda(\beta_1) + b + 1$, $V_1 = [b + 1, n - 1]$, and $V_2 = [1, a - 1]$.

To find a bijective proof of the second part of Theorem 9 we will first discuss a decomposition of paths in $\mathcal{B}^{[2]}$ which is similar to the decomposition of permutations in $\mathcal{R}(123)$ underlying (13). Let $\beta \in \mathcal{B}^{[2]}$. If there is a leftmost occurrence of \bar{u} in β then the height of that \bar{u} must be either 0 or 1. Thus we have

$$\mathcal{B}^{[2]} = \mathcal{D} + \mathcal{D}\bar{u}\mathcal{B}^{[1]}d\mathcal{B}^{[2]} + \mathcal{D}u\mathcal{D}\bar{u}d\mathcal{B}^{[1]}d\mathcal{B}^{[2]} \quad (16)$$




FIGURE 1. The bijection $\mathcal{B}^{[2]} \setminus \mathcal{D} \xrightarrow{\Phi} (\mathcal{D} \setminus \{\epsilon\}) \times (\tilde{\mathcal{B}}^{[2]} \setminus \{\epsilon\})$

whose commutative counterpart is

$$\begin{aligned} B^{[2]} &= C + qtCB^{[1]}B^{[2]} + qt^2C^3B^{[1]}B^{[2]} \\ &= C + qt^{-1}(tC + t^2C^3)tB^{[1]}B^{[2]}. \end{aligned} \quad (17)$$

Since $C = 1 + tC^2$, the factor $tC + t^2C^3$ simplifies to $C - 1$. Moreover, if we let $\tilde{\mathcal{B}}^{[2]}$ denote the set of paths in $\mathcal{B}^{[2]}$ whose first step is u (i.e., not \bar{u}), then

$$\tilde{\mathcal{B}}^{[2]} = 1 + u\mathcal{B}^{[1]}d\mathcal{B}^{[2]}, \quad (18)$$

and, as a consequence, $tB^{[1]}B^{[2]} = \tilde{\mathcal{B}}^{[2]} - 1$. Thus (17) can be rewritten as

$$B^{[2]} = C + qt^{-1}(C - 1)(\tilde{\mathcal{B}}^{[2]} - 1),$$

which should be compared to (13). This suggests that we should be able to uniquely decompose any path β in $\mathcal{B}^{[2]} \setminus \mathcal{D}$ into two nonempty paths $\beta' \in \mathcal{D}$ and $\beta'' \in \tilde{\mathcal{B}}^{[2]}$ such that $|\beta| = |\beta'| + |\beta''| - 1$ and $|\beta|_{\bar{u}} = |\beta''|_{\bar{u}} + 1$. Indeed, using (16), (18) and

$$\mathcal{D} = 1 + \mathcal{D}u\mathcal{D} + \mathcal{D}u\mathcal{D}u\mathcal{D}d\mathcal{D},$$

such a decomposition is defined by the map

$$\begin{aligned} \beta_1 \bar{u} \beta_2 d \beta_3 &\mapsto \langle \beta_1 u d, u \beta_2 d \beta_3 \rangle, \\ \beta_1 u \beta'_1 \bar{u} \beta''_1 d \beta_2 d \beta_3 &\mapsto \langle \beta_1 u \beta'_1 u \beta''_1 d d, u \beta_2 d \beta_3 \rangle, \end{aligned}$$

where $\beta_1, \beta'_1, \beta''_1 \in \mathcal{D}$, $\beta_2 \in \mathcal{B}^{[1]}$, and $\beta_3 \in \mathcal{B}^{[2]}$. We denote by Φ the inverse of this map; it is obtained by simply reversing the arrows. See Figure 3 for a schematic diagram of Φ .

Let h be any bijection from $\mathcal{S}_n(123)$ to \mathcal{D}_n . For definiteness, we can take h to be the bijection Ψ given by Krattenthaler in [5, p. 522]. (A description of Ψ can be found in the example following this proof.) We shall define a bijection

$$g : \mathcal{R}(123) \rightarrow \mathcal{B}^{[2]}$$

such that $|g(\pi)| = 2(n - 2k)$ and $|g(\pi)|_{\bar{u}} = k$, whenever $\pi \in \mathcal{R}_n^k(123)$. If π avoids 123 then let $g(\pi) = h(\pi)$. If π does not avoid 123 then, as in the first proof, we can

write $\pi = \pi_1 abc \pi_2$, where abc is the leftmost occurrence of 123 in π ; in this case, we let

$$g(\pi) = \Phi\langle (g \circ \text{red})(\pi_1 c), (g \circ \text{red})(a\pi_2) \rangle.$$

Proving that g is invertible is similar to proving that f is invertible. \square

We remark that the bijection f from the first part of the preceding proof maps 132-avoiding permutations onto Dyck paths. In fact, the restriction of f to $\mathcal{S}(132)$ is a bijection due to Krattenthaler [5, p. 512].

Example 6. The permutation 846572931 is 132-segmented. It has two occurrences of 132, namely 465 and 293. We illustrate the bijection f , from the first part of the preceding proof, by finding the image of 846572931 under f :

$$\begin{aligned} f(846572931) &= \bar{u}f(84657)df(1) = \bar{u}udf(4657)dud = \\ &= \bar{u}uduf(465)ddud = \bar{u}udu\bar{u}dddud. \end{aligned}$$

For convenience we have not reduced the permutations in the intermediate steps.

To give an example of how g , from the second part of the preceding proof, is applied, we first need to describe Krattenthaler's [5, p. 522] bijection Ψ from $\mathcal{S}_n(123)$ to \mathcal{D}_n . Let $\pi = a_1 a_2 \cdots a_n$ be a 123-avoiding permutation. A *right-to-left maximum* is an element a_i such that $a_i > a_j$ for all $j > i$. Let the right-to-left maxima in π be m_1, m_2, \dots, m_s , from right to left, so that

$$\pi = \pi_s m_s \cdots \pi_2 m_2 \pi_1 m_1,$$

where π_i is the subword of π between m_{i+1} and m_i . If there is an occurrence ab of 12 in π_i then abm_i is an occurrence of 123 in π . Therefore, the elements in π_i are in decreasing order. Moreover, we have $\pi_i < \pi_{i+1}$.

The Dyck path $\Psi(\pi)$ is generated from right to left: Read π from right to left. Any right-to-left maximum m_i is translated into $m_i - m_{i-1}$ up-steps (with the convention $m_0 = 0$). Any subword π_i is translated into $|\pi_i| + 1$ down-steps.

We are now ready for an illustration of g . The permutation 957841362 is 123-segmented. It has two occurrences of 123, namely 578 and 136. To find the image of 957841362 under g we proceed as follows:

$$\begin{aligned} g(957841362) &= \Phi\langle (g \circ \text{red})(98), (g \circ \text{red})(541362) \rangle, \\ (g \circ \text{red})(98) &= \Psi(21) = udud, \\ (g \circ \text{red})(541362) &= \Phi\langle (g \circ \text{red})(546), (g \circ \text{red})(12) \rangle, \\ (g \circ \text{red})(546) &= \Psi(213) = uuudd, \\ (g \circ \text{red})(12) &= \Psi(12) = uudd, \\ \Phi(uuudd, uudd) &= \bar{u}\bar{u}uddudd, \\ \Phi(udud, \bar{u}\bar{u}uddudd) &= u\bar{d}\bar{u}\bar{u}uddudd. \end{aligned}$$

Thus $g(957841362) = u\bar{d}\bar{u}\bar{u}uddudd$.

Corollary 7. For $k \geq 0$ and $n \geq 0$ we have

$$\text{card } R_n^k(123) \leq \text{card } \mathcal{R}_n^k(132).$$

Proof. The result follows immediately from $\mathcal{B}_{n,k}^{[2]} \subseteq \mathcal{B}_{n,k}$ and Theorem 9. \square

Corollary 8. The generating functions $R(132; q, t)$ and $R(123; q, t)$ admit the following continued fraction expansions:

$$R(132; q, t) = \frac{1}{1 - \frac{t + qt^3}{1 - \frac{t + qt^3}{1 - \frac{t + qt^3}{\ddots}}}}, \quad R(123; q, t) = \frac{1}{1 - \frac{t + qt^3}{1 - \frac{t}{1 - \frac{t}{1 - \frac{t}{\ddots}}}}}.$$

Proof. Using (9) and Theorem 9 the result follows from iterating the identity $C(t) = 1/(1 - tC(t))$. \square

Proposition 9. The generating function

$$R(123, 132; p, q, t) = \sum_{\pi \in \mathcal{R}(123) \cap \mathcal{R}(132)} p^{(123)} \pi q^{(132)} \pi t^{|\pi|}$$

counting $\{123, 132\}$ -segmented permutations by occurrences of 123 and 132 is the following rational function:

$$R(123, 132; p, q, t) = \frac{1-t}{1-2t-(p+q)t^3} = \frac{1}{1 - \frac{t + (p+q)t^3}{1-t}}.$$

First proof. Let n be a positive integer, and let π be a $\{123, 132\}$ -segmented permutation of length n . We distinguish between three cases:

- (a) If the letter n is not part of any occurrence of 123 or 132, then we can factor π as $\pi = \pi_1 n \pi_2$, where π_1 is 12-avoiding, π_2 is $\{123, 132\}$ -segmented, and $\pi_2 < \pi_1$.
- (b) If the letter n is part of an occurrence of 123, then we can factor π as $\pi = \pi_1 abn \pi_2$, where π_1 is 12-avoiding, π_2 is $\{123, 132\}$ -segmented, and $\pi_2 < a < b < \pi_1$.
- (c) If the letter n is part of an occurrence of 132, then we can factor π as $\pi = \pi_1 anb \pi_2$, where π_1 is 12-avoiding, π_2 is $\{123, 132\}$ -segmented, and $\pi_2 < a < b < \pi_1$.

It is clear that an occurrence of 123 can not overlap with an occurrence of 132 without creating a non-segment occurrence of 123 or 132. Therefore, the cases (b) and (c) are mutually exclusive. Thus the generating function $R = R(123, 132; p, q, t)$ satisfies

$$R = 1 + R(12; 0, t)(t + pt^3 + qt^3)R, \quad (19)$$

where $R(12; 0, t) = 1/(1-t)$ is the generating function for 12-avoiding permutations. Solving (19) for R we obtain the desired result. \square

Second proof. To give a bijective proof of Proposition 9 we consider lattice paths with three different types of up-steps: let \mathcal{T} be the language over $\{u, \bar{u}, \bar{d}\}$ whose characteristic series is implicitly given by

$$\mathcal{T} = 1 + (u + \bar{u} + \bar{d}) \cdot \frac{1}{1-ud} \cdot d\mathcal{T}.$$

We may think of a word in the language \mathcal{T} as a 3-coloured Dyck path whose u -steps are of height 0 or 1, and whose \bar{u} - and \bar{d} -steps are of height 0. Applying the substitution $\mu : \mathbb{C}\langle u, \bar{u}, \bar{d} \rangle \rightarrow \mathbb{C}[p, q, t]$ defined by

$$\mu = \{ u \mapsto 1, \bar{u} \mapsto p, \bar{d} \mapsto q, d \mapsto t \},$$

we see that $R(123, 132, p, q, t) = \mu(\mathcal{T})(pt^2, qt^2, t)$. We shall give a bijection

$$f : \mathcal{R}(123) \cap \mathcal{R}(132) \rightarrow \mathcal{T}$$

such that $|\pi| = |\beta|_u + 3(|\beta|_{\bar{u}} + |\beta|_{\bar{a}})$, where $\beta = f(\pi)$. Following the decomposition given in the first proof, we recursively define f as follows:

$$\begin{aligned} f(\epsilon) &= \epsilon, \\ f(\pi_1 n \pi_2) &= u(f \circ \text{red})(\pi_1) d(f \circ \text{red})(\pi_2), \\ f(\pi_1 a b n \pi_2) &= \bar{u}(f \circ \text{red})(\pi_1) d(f \circ \text{red})(\pi_2), \\ f(\pi_1 a n b \pi_2) &= \bar{\bar{u}}(f \circ \text{red})(\pi_1) d(f \circ \text{red})(\pi_2). \end{aligned}$$

It is straightforward, but tedious, to give the inverse of f . □

Example. The permutation 875963124 is {123, 132}-segmented. It has one occurrence of each of the patterns 123 and 132, namely 124 and 596. To illustrate the second proof of Proposition 9 we find the image of 875963124 under f :

$$f(875963124) = \bar{\bar{u}}f(87)df(3124) = \bar{\bar{u}}udf(7)d\bar{u}f(3)d = \bar{\bar{u}}ududd\bar{\bar{u}}udd.$$

4. ACKNOWLEDGMENTS

I thank Petter Brändén for several discussions on the subject of this paper. I also thank Einar Steingrímsson for proofreading a preliminary version of this article.

5. TABLES

card $B_{n,k}$:

$n \setminus k$	0	1	2	3	4	5	6	7
0	1							
1	1	1						
2	2	4	2					
3	5	15	15	5				
4	14	56	84	56	14			
5	42	210	420	420	210	42		
6	132	792	1980	2640	1980	792	132	
7	429	3003	9009	15015	15015	9009	3003	429

card $B_{n,k}^{[1]}$:

$n \setminus k$	0	1	2	3	4	5	6	7
0	1							
1	1	1						
2	2	3	1					
3	5	9	5	1				
4	14	28	20	7	1			
5	42	90	75	35	9	1		
6	132	297	275	154	54	11	1	
7	429	1001	1001	637	273	77	13	1

card $B_{n,k}^{[2]}$:

$n \setminus k$	0	1	2	3	4	5	6	7
0	1							
1	1	1						
2	2	4	2					
3	5	14	13	4				
4	14	48	62	36	8			
5	42	165	264	217	92	16		
6	132	572	1066	1104	670	224	32	
7	429	2002	4186	5130	3965	1912	528	64

REFERENCES

- [1] T. Chow and J. West. Forbidden subsequences and Chebyshev polynomials. *Discrete Math.*, 204(1-3):119–128, 1999.
- [2] S. Elizalde and M. Noy. Enumeration of subwords in permutations. In *Formal power series and algebraic combinatorics (Tempe, 2001)*, pages 179–189. Arizona State University, 2001.
- [3] S. Kitaev. Multi-avoidance of generalised patterns. *Discrete Math.*, 260(1-3):89–100, 2003.
- [4] S. Kitaev and T. Mansour. Simultaneous avoidance of generalized patterns. to appear in *Ars Combinatoria*.
- [5] C. Krattenthaler. Permutations with restricted patterns and dyck paths. *Advances in Applied Mathematics*, 27:510, August 2001.
- [6] T. Mansour and A. Vainshtein. Counting occurrences of 132 in a permutation. *To appear in: Adv. Appl. Math.*, 2001.
- [7] J. Riordan. *Combinatorial identities*. John Wiley & Sons Inc., New York, 1968.
- [8] R. Simion and F. W. Schmidt. Restricted permutations. *European J. Combin.*, 6(4):383–406, 1985.

MATEMATIK, CHALMERS TEKNISKA HÖGSKOLA OCH GÖTEBORGS UNIVERSITET, S-412 96 GÖTEBORG, SWEDEN
E-mail address: claesson@math.chalmers.se