

## Solutions to Chapter 5 Exercises (Part 2)

### Problem 5.5

$$f_{X,Y}(x,y) = \frac{1}{8\pi} \exp\left(-\frac{(x-1)^2 + (y+1)^2}{8}\right)$$

(a)  $Pr(X > 2, Y < 0)$

$$\begin{aligned} Pr(X > 2, Y < 0) &= \int_2^\infty \int_{-\infty}^0 \frac{1}{8\pi} \exp\left(-\frac{(x-1)^2 + (y+1)^2}{8}\right) dy dx \\ &= \int_2^\infty \frac{1}{\sqrt{8\pi}} \exp\left(-\frac{(x-1)^2}{8}\right) dx \int_{-\infty}^0 \frac{1}{\sqrt{8\pi}} \exp\left(-\frac{(y+1)^2}{8}\right) dy \\ &= Q\left(\frac{2-1}{2}\right) \Phi\left(\frac{0+1}{2}\right) = Q\left(\frac{1}{2}\right) \Phi\left(\frac{1}{2}\right) \\ &= Q(1/2)(1 - Q(1/2)) \end{aligned}$$

(b)  $Pr(0 < X < 2, |Y+1| > 2)$

$$\begin{aligned} &= Pr(\{0 < X < 2, Y > 1\} \cup \{0 < X < 2, Y < -3\}) \\ &= \int_0^2 \int_{-\infty}^{-3} f_{X,Y}(x,y) dx dy + \int_0^2 \int_1^\infty f_{X,Y}(x,y) dx dy \\ &= \int_0^2 \frac{1}{\sqrt{8\pi}} \exp\left(-\frac{(x-1)^2}{8}\right) dx \left( \int_{-\infty}^{-3} \frac{1}{\sqrt{8\pi}} \exp\left(-\frac{(y+1)^2}{8}\right) dy + \int_1^\infty \frac{1}{\sqrt{8\pi}} \exp\left(-\frac{(y+1)^2}{8}\right) dy \right) \\ &= \left( Q\left(\frac{0-1}{2}\right) - Q\left(\frac{2-1}{2}\right) \right) \left( \Phi\left(\frac{-3+1}{2}\right) + Q\left(\frac{1+1}{2}\right) \right) \\ &= \left( Q\left(-\frac{1}{2}\right) - Q\left(\frac{1}{2}\right) \right) (\Phi(-1) + Q(1)) \\ &= 2Q(1) \left( 1 - 2Q\left(\frac{1}{2}\right) \right) \end{aligned}$$

(c)  $Pr(Y > X)$

$$Pr(Y > X) = \int_{-\infty}^{\infty} \int_{-\infty}^y f_{X,Y}(x,y) dx dy$$

This integral is easier to do with a change of variables. Let us use the substitution

$$\begin{aligned} u &= x - y \\ v &= x + y \\ J \begin{pmatrix} u & v \\ x & y \end{pmatrix} &= \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \\ \left| \det \begin{bmatrix} u & v \\ x & y \end{bmatrix} \right| &= 2 \\ f_{U,V}(u,v) &= \frac{f_{X,Y}(x,y)}{2} \\ Pr(Y > X) &= Pr(U < 0) \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^0 \frac{1}{8\pi} \exp\left(-\frac{(x-1)^2 + (y+1)^2}{8}\right) \frac{1}{2} du dv \\ &= \frac{1}{16\pi} \int_{-\infty}^{\infty} \int_{-\infty}^0 \exp\left(-\frac{(u-v-2)^2 + (u+v+2)^2}{32}\right) du dv \\ &= \frac{1}{16\pi} \int_{-\infty}^{\infty} \int_{-\infty}^0 \exp\left(-\frac{(u^2 + v^2 + 4 - 4u)}{16}\right) du dv \\ &= \frac{1}{16\pi} \int_{-\infty}^{\infty} \int_{-\infty}^0 \exp\left(-\frac{(u-2)^2 + v^2}{16}\right) du dv \\ &= \frac{1}{16\pi} \int_{-\infty}^{\infty} \exp\left(-\frac{v^2}{16}\right) dv \int_{-\infty}^0 \exp\left(-\frac{(u-2)^2}{16}\right) du \\ &= \frac{1}{16\pi} \sqrt{16\pi} \sqrt{16\pi} \Phi\left(\frac{0-2}{\sqrt{8}}\right) \\ &= Q\left(\frac{1}{\sqrt{2}}\right) \end{aligned}$$

## Problem 5.29

Note that any linear transformation of jointly Gaussian random variables produces jointly Gaussian random variables. So U, V are both Gaussian random variables. And if  $\rho_{U,V} = 0$ , U and V are independent.

$$E[U] = aE[X] + bE[Y] = 0 .$$

$$E[V] = cE[X] + dE[Y] = 0 .$$

$$E[UV] = acE[X^2] + bdE[Y^2] + (ad + bc)E[XY] = ac + bd + \rho(ad + bc) .$$

Thus,  $\rho_{U,V} = 0$  if

$$ac + bd + \rho(ad + bc) = 0 .$$

So, U and V are independent, if

$$ac + bd = -\rho(ad + bc) .$$

### Problem 5.32

$$\begin{aligned} Z &= X^2 + Y^2 \\ W &= XY \\ X &= \frac{\sqrt{Z+2W} + \sqrt{Z-2W}}{2} \\ Y &= \frac{\sqrt{Z+2W} - \sqrt{Z-2W}}{2} \\ f_X(x) &= \frac{1}{\pi(1+x^2)} \\ f_Y(y) &= \frac{1}{\pi(1+y^2)} \\ J \begin{pmatrix} z & w \\ x & y \end{pmatrix} &= \begin{pmatrix} 2x & 2y \\ y & x \end{pmatrix} \\ \left| \det \left[ J \begin{pmatrix} z & w \\ x & y \end{pmatrix} \right] \right| &= 2|x^2 - y^2| \\ f_{Z,W}(z,w) &= \frac{f_X(x)f_Y(y)}{2|x^2 - y^2|} \\ &= \frac{1}{2\pi^2} \frac{1}{(1+x^2)(1+y^2)|x^2 - y^2|} \\ &= \frac{1}{2\pi^2} \frac{1}{(1+x^2 + y^2 + x^2y^2)|x^2 - y^2|} \\ &= \frac{1}{2\pi^2} \frac{1}{(1+z^2 + w^2)\sqrt{z^2 - 4w^2}} \end{aligned}$$