

Financial Risk 2-rd quarter 2012/2013 Tuesdays 10:15-12 Thursdays 13:15-15:00 in MVF31 and Pascal

Copyright © by Randy Glasbergen. www.glasbergen.com



"As an alternative to the traditional 30-year mortgage, we also offer an interest-only mortgage, balloon mortgage, reverse mortgage, upside down mortgage, inside out mortgage, loop-de-loop mortgage, and the spinning double axel mortgage with a triple lutz."



Gudrun January 2005
326 MEuro loss
72 % due to forest losses
4 times larger than second largest

## Maximum Likelihood (ML) inference (Coles p. 30-43)

Likelihood function = the function which shows how the "probability" (or likelihood) of getting the observed data depends on the parameters

$$x_1, \ldots x_n$$
 observations of i.i.d. variables  $X_1, \ldots X_n$ , density  $f(x) = f(x; \theta)$ 

$$\theta = (\theta_1, \dots \theta_d)$$
 parameters

$$L(\theta) = f(x_1; \theta) f(x_2; \theta) \dots f(x_n; \theta)$$
 likelihood function

$$\ell(\theta) = \log f(x_1; \theta) + \log f(x_2; \theta) + \ldots \log f(x_n; \theta) = \sum_{i=1}^n \log f(x_i; \theta)$$

ML estimates = the value  $\hat{\theta} = (\hat{\theta_1}, \dots \hat{\theta_d})$  which maximizes  $\ell(\theta)$ 

- MI estimates often have to be found through numerical maximization
- sometimes a maximum doesn't exist
- sometimes several local maxima (→ problem for numerical maximization)
- but typically no problems if the number of observations is "large"

### Example: ML estimation of the parameters in the PoT model

T = length of observation period

N = number of observed excesses (random variable!)

 $x_1, \ldots x_N$  observed excess sizes

$$\theta = (\lambda, \beta, \sigma)$$
 parameters

The probability of observing N excesses is  $\frac{(\lambda T)^N}{N!} \exp\{-\lambda T\}$ , plus independence and previous slide  $\rightarrow$ 

$$L(\theta) = L(\lambda, \sigma, \gamma) = \frac{(\lambda T)^N}{N!} \exp\{-\lambda T\} \prod_{i=1}^N \frac{1}{\sigma} \left(1 + \frac{\gamma}{\sigma} x_i\right)_+^{-1/\gamma - 1}$$

$$\ell(\lambda, \sigma, \gamma) = N \log(\lambda) + N \log(T) - \log(N!) - \lambda T$$

$$- N \log(\sigma) - \sum_{i=1}^N (1/\gamma + 1) \log\left(1 + \frac{\gamma}{\sigma} x_i\right)_+$$

$$\frac{\partial}{\partial \lambda} \ell(\lambda, \sigma, \gamma) = \frac{N}{\lambda} - T = 0 \quad \text{so that} \quad \hat{\lambda} = \frac{N}{T}$$

 $\hat{\sigma}, \hat{\gamma}$  obtained from numerical maximization of the second part of  $\ell(\lambda, \sigma, \gamma)$ 

## ML inference: asymptotic properties

 $\mathcal{I}(\theta) = E_{\theta}(\left(-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \ell(\theta)\right))$  expected Fisher information matrix, estimated by  $\mathcal{I}(\hat{\theta})$  or by  $I(\hat{\theta})$  where  $I(\theta) = \left(-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \ell(\theta)\right)$  is the the observed Fisher information matrix. (In the expected Fisher information matrix, the observations are replaced by the corresponding random variables when the expectations are computed)

 $\hat{\theta} = (\hat{\theta_1}, \dots \hat{\theta_d})$  asymptotically has a d-dimensional multivariate normal distribution with mean  $\theta$  and variance  $\mathcal{I}(\theta)^{-1}$ 

In particular, the variance of  $\hat{\theta}_i$  may be estimated by  $(\mathcal{I}(\hat{\theta})^{-1})_{ii}$  (= the *i*-th diagonal element of  $\mathcal{I}(\hat{\theta})^{-1}$ ), or by  $(I(\hat{\theta})^{-1})_{ii}$ . The latter is often more accurate.

 $k_{\alpha/2}$  = the  $\alpha/2$ -th quantile from the top of the standard normal distribution

$$(\hat{\theta}_i - k_{\alpha/2} \sqrt{(I(\hat{\theta})^{-1})_{i,i}}, \ \hat{\theta}_i + k_{\alpha/2} \sqrt{(I(\hat{\theta})^{-1})_{i,i}})$$
 asymptotic 100(1-  $\alpha$ ) % confidence interval for  $\theta_i$ 

#### ML inference: the delta method

$$\eta = g(\theta) = g(\theta_1, \dots \theta_d)$$
 function of the parameters

$$\hat{\eta} = g(\hat{\theta}) = g(\hat{\theta}_1, \dots \hat{\theta}_d)$$
 estimate of the function of the parameters

$$\nabla(\theta) = (\frac{\partial}{\partial \theta_1} g(\theta), \dots \frac{\partial}{\partial \theta_d} g(\theta))$$
 gradient,  $\nabla(\hat{\theta})$  estimate of gradient

 $\hat{\eta}$  asymptotically normal with mean  $\eta$  and variance  $\nabla(\theta)\mathcal{I}(\theta)^{-1}\nabla(\theta)^t$  (which e.g. can be estimated by  $\nabla(\hat{\theta})I(\hat{\theta})^{-1}\nabla(\hat{\theta})^t$ .

From this one can construct confidence intervals for  $\eta$  in the same way as the confidence intervals for  $\theta$  on the previous page.

## ML inference: Likelihood Ratio (LR) tests

 $\theta=(\theta_1,\theta_2)$  partition of  $\theta$  into two vectors  $\theta_1$  and  $\theta_2$  of dimensions d-p and p.  $\hat{\theta}_2^*$  maximizes  $l(\theta_1,\theta_2)$  over  $\theta_2$ , for  $\theta_1$  "kept fixed" (so function of  $\theta_1$ )

 $2(\ell(\hat{\theta}) - \ell(\theta_1, \hat{\theta}_2^*))$  asymptotically has a  $\chi^2$  distribution with d-p degrees of freedom if  $\theta_1$  is the true value  $\rightarrow$  LR test:

Reject  $H_0$ :  $\theta_1 = \theta_1^0$  at the significance level 100  $\alpha$ % if

### ML inference: profile likelihood confidence intervals

(often more accurate than delta method intervals, *plots from Coles* )




FIGURE 4.3. Profile likelihood for  $\xi$  in threshold excess model of daily rainfall data.





FIGURE 4.4. Profile likelihood for 100-year return level in threshold excess model of daily rainfall data.

Profile likelihood confidence intervals for the shape parameter in the Block Maxima model. The delta method probably would give similar interval in the left case, but not in the right.





The real problem!

## The problems

How much reinsurance should LFAB buy?

Should LFAB worry about windstorm losses getting worse?

How should LFAB adjust if its forest insurance portfolio grows?

and:

Can detailed modeling give better risk estimates?

Are windstorms becoming more frequent?

## 1994 PoT analysis of 1982-1993 LFAB data (the basic method, more sophisticated analysis of 1982-2005 data in later paper)

Risk	next	next 5	next 15
(MSEK)	year	years	years
10%	66	215	473
1%	366	1149	2497


$$X_i$$
 GP( $y; \sigma_t, \gamma$ )

$$\sigma_t = \exp(\alpha + \beta t)$$

$$\hat{\alpha} = 15.1$$

$$\hat{\beta} = .013 \pm .013$$

no evidence of trend in extremes



conditional probability that a loss in excees of the reinsurance level 850 MSEK exceeds x

Gudrun: 2912 MSEK, 12 years later


Windstorms of 1902 and 1969 probably comparable to Gudrun

## Choice of threshold/number of order statistics in PoT, model diagnostics


Threshold choice compromise between low bias (= good fit of model), which requires high threshold/few order statistics, and low variance, which requires low threshold/many order statistics


- mean excess plots (high variability for heavy tails)
- median excess plots
- plots of parameter estimates as function of threshold/number of order statistics
- qq- and pp-plots


automatic threshold selection procedures exist, but perhaps not all that reliable ("optimal" threshold depends on the underlying distribution which is unknown and has to be estimated).




Windstorm losses 1982-1993 excesses of 0.9 MSEK









### Some conclusions

- risk cannot be summarized into one number
- extreme value statistics provide the simplest methods (but other methods may sometimes be needed)
- didn't find clear trends
- meteorological data didn't help
- don't trust computer simulation models unless statistically validated
- companies should develop systematic techniques for thinking about "not yet seen" catastrophes
- put contractual limits to aggregate exposure

# A step in another direction: catastrophe risks

## BIG --- "happens only once"

- can't adjust and improve as experience is gained
- methods based on means, variances, central limit theory have little meaning
- difficult to keep in mind that catastrophes can (and will!) occur

a gamble --- find the odds of a gamble!

- O. Perrin, H. Rootzén, and R. Taessler, A discussion of statistical methods used to estimate extreme wind speeds. *Theor. Appl. Climatol.* **85**, 203–215 (2006)
- H. Rootzén and N. Tajvidi (1997). Extreme value statistics and wind storm losses: a case study. *Scand. Actuarial J.*, 70-94, reprinted in "Extremes and integrated risk management", Risk Books 2000.
- H. Rootzén, and N.Tajvidi (2000). Can losses caused by wind storms be predicted from meteorological observations? *Scand. Actuarial J.*, 162-175.
- H. Rootzén, and N.Tajvidi (2006). The multivariate Generalized Pareto Distribution. *Bernoulli* 12, 917-930
- E. Brodin and H. Rootzén (2009). Modelling and predicting extreme wind storm losses. *Insurance, Mathematics and Economics*