Financial Risk: Credit Risk, Lecture 1

Financial Risk, Chalmers University of Technology and University of Gothenburg, Göteborg Sweden

Spring 2019

(CTH and Univ. of Gothenburg)

Financial Risk: Credit Risk, Lecture 1

Spring 2019 1 / 26

Main goal of lectures and content of today's lecture

- The main goal of coming three lectures is to study the loss distribution for a credit portfolio
- The loss distribution is used to compute risk measures such as Value-at-Risk etc.

Today's lecture

- Short discussion of the important components of credit risk
- Study different static portfolio credit risk models.
- Discussion of the binomial loss model
- Discussion of the mixed binomial loss model

Credit risk

 $-\,$ the risk that an obligor does not honor his payments

Example of an obligor:

- A company that have borrowed money from a bank
- A company that has issued bonds.
- A household that have borrowed money from a bank, to buy a house
- A bank that has entered into a bilateral financial contract (e.g an interest rate swap) with another bank.

Example of defaults are

- A company goes bankrupt.
- As company fails to pay a coupon on time, for some of its issued bonds.
- A household fails to pay amortization or interest rate on their loan.

(CTH and Univ. of Gothenburg)

Financial Risk: Credit Risk, Lecture 1

Credit Risk

Credit risk can be decomposed into:

- arrival risk, the risk connected to whether or not a default will happen in a given time-period, for a obligor
- **timing risk**, the risk connected to the uncertainness of the exact time-point of the arrival risk (will not be studied in this course)
- recovery risk. This is the risk connected to the size of the actual loss if default occurs (will not be studied in this course, we let the recovery be fixed)
- **default dependency risk**, the risk that several obligors jointly defaults during some specific time period. This is one of the most crucial risk factors that has to be considered in a credit portfolio framework.
- The coming three lectures focuses only on default dependency risk.

Portfolio Credit Risk is important

- Portfolio credit risk models differ greatly depending on what types of portfolios, and what type of questions that should be considered. For example,
 - models with respect to risk management, such as credit Value-at-Risk (VaR) and expected shortfall (ES)
 - models with respect to valuation of portfolio credit derivatives, such as CDO's and basket default swaps
- In both cases we need to consider default dependency risk, but....
- ...in risk management modelling (e.g. VaR, ES), the timing risk is ignored, and one often talk about **static credit portfolio models**,
- ...while, when pricing credit derivatives, timing risk must be carefully modeled (not treated here)
- The coming three lectures focuses only on static credit portfolio models,

The slides for the coming three lectures are rather self-contained, but more details on certain topics can be found in the lecture notes.

The content of the lecture today and the two next lectures are **partly** based on material presented in:

- "*Quantitative Risk Management*" by McNeil A., Frey, R. and Embrechts, P. (Princeton University Press)
- "Credit Risk Modeling: Theory and Applications" by Lando, D . (Princeton University Press)
- "*Risk and portfolio analysis principles and methods*" by Hult, Lindskog, Hammerlid and Rehn. (Springer)

Static Models for homogeneous credit portfolios

- Today we will consider the following static modes for a homogeneous credit portfolio:
 - The binomial model
 - The mixed binomial model
- To understand mixed binomial models, we give a short introduction of conditional expectations
- In the next two lectures we will
 - study three different mixed binomial models.
 - discuss Value-at-Risk and Expected shortfall in a mixed binomial models.
 - Correlations etc in mixed binomial models.

Consider a homogeneous credit portfolio model with m obligors where each obligor can default up to fixed time T, and have the same constant credit loss ℓ .

• Let X_i be a random variable such that

$$X_{i} = \begin{cases} 1 & \text{if obligor } i \text{ defaults before time } T \\ 0 & \text{otherwise, i.e. if obligor } i \text{ survives up to time } T \end{cases}$$
(1)

- We assume that $X_1, X_2, ..., X_m$ are **i.i.d**, that is they are independent with identical distribution. Furthermore $\mathbb{P}[X_i = 1] = p$ so $\mathbb{P}[X_i = 0] = 1 p$.
- The total credit loss in the portfolio at time T, called L_m , is then given by

$$L_m = \sum_{i=1}^m \ell X_i = \ell \sum_{i=1}^m X_i = \ell N_m \quad \text{where } N_m = \sum_{i=1}^m X_i$$

thus, N_m is the **number** of defaults in the portfolio up to time T.

• Since ℓ is a constant, we have $\mathbb{P}[L_m = k\ell] = \mathbb{P}[N_m = k]$, so it is enough to study the distribution of N_m .

- Since X₁, X₂,... X_m are i.i.d with P [X_i = 1] = p we see that N_m = ∑_{i=1}^m X_i is binomially distributed with parameters m and p, i.e. N_m ~ Bin(m, p).
- Hence, we have

$$\mathbb{P}\left[N_m=k\right] = \binom{m}{k} p^k (1-p)^{m-k}$$

• Recalling the binomial theorem $(a + b)^m = \sum_{k=0}^m {m \choose k} a^k b^{m-k}$ we see that


$$\sum_{k=0}^{m} \mathbb{P}[N_m = k] = \sum_{k=0}^{m} \binom{m}{k} p^k (1-p)^{m-k} = (p+(1-p))^m = 1$$

proving that Bin(m, p) is a distribution.

• Furthermore, $\mathbb{E}[N_m] = mp$ since

$$\mathbb{E}[N_m] = \mathbb{E}\left[\sum_{i=1}^m X_i\right] = \sum_{i=1}^m \mathbb{E}[X_i] = mp.$$

Financial Risk: Credit Risk, Lecture 1



(CTH and Univ. of Gothenburg)

Financial Risk: Credit Risk, Lecture 1

Spring 2019 10 / 26

- The binomial distribution have very thin "tails", that is, it is extremely unlikely to have many losses (see figure).
- For example, if p = 5% and m = 50 we have that $\mathbb{P}[N_m \ge 8] = 1.2\%$ and for p = 10% and m = 50 we get $\mathbb{P}[N_m \ge 10] = 5.5\%$
- The main reason for these small numbers is due to the independence assumption for $X_1, X_2, \dots X_m$.
- To see this, recall that the variance Var(X) measures the degree of the deviation of X around its mean, i.e. $Var(X) = \mathbb{E}\left[(X \mathbb{E}[X])^2\right]$.
- Since $X_1, X_2, \ldots X_m$ are independent we have that

$$\operatorname{Var}(N_m) = \operatorname{Var}\left(\sum_{i=1}^m X_i\right) = \sum_{i=1}^m \operatorname{Var}(X_i) = mp(1-p) \tag{2}$$

where the second equality is due the independence assumption.

• Furthermore, by Chebyshev's inequality we have that for any random variable X, and any c > 0 it holds

$$\mathbb{P}\left[|X - \mathbb{E}\left[X\right]| \ge c
ight] \le \frac{\operatorname{Var}(X)}{c^2}.$$

- Example: if p = 5% and m = 50 then $Var(N_m) = 50p(1-p) = 2.375$ and $\mathbb{E}[N_m] = 50p = 2.5$.
- So with p = 5% and m = 50, the probability of say, 6 more or less losses than expected, is smaller or equal than 6.6%, since by Chebyshev

$$\mathbb{P}\left[|N_m - 2.5| \ge 6\right] \le \frac{2.375}{36} = 6.6\%.$$

• Next we show that the deviation of the fractional number of defaults in the portfolio, $\frac{N_m}{m}$, from the constant $p = \mathbb{E}\left[\frac{N_m}{m}\right]$, goes to zero as $m \to \infty$.

• So $\frac{N_m}{m}$ converges towards a constant as $m \to \infty$ (the law of large numbers).

Independent defaults and the law of large numbers

- By applying Chebyshev's inequality to $\frac{N_m}{m}$ together with Equation (2) we get $\mathbb{P}\left[\left|\frac{N_m}{m} - p\right| \ge \varepsilon\right] \le \frac{\operatorname{Var}\left(\frac{N_m}{m}\right)}{\varepsilon^2} = \frac{\frac{1}{m^2}\operatorname{Var}\left(N_m\right)}{\varepsilon^2} = \frac{mp(1-p)}{m^2\varepsilon^2} = \frac{p(1-p)}{m\varepsilon^2}$
- Thus, $\mathbb{P}\left[\left|\frac{N_m}{m} p\right| \ge \varepsilon\right] \to 0$ as $m \to \infty$ for any $\varepsilon > 0$.
- This result is called the weak law of large numbers
- For our credit portfolio it means that the fractional number of defaults in the portfolio, i.e. $\frac{N_m}{m}$, converges (in probability) to the constant p, i.e the individual default probability.
- One can also show the so called strong law of large numbers, that is

$$\mathbb{P}\left[\frac{N_m}{m} \to p \text{ when } m \to \infty\right] = 1$$

and we say that $\frac{N_m}{m}$ converges almost surely to the constant p. In these lectures we write $\frac{N_m}{m} \rightarrow p$ to indicate almost surely convergence.

Independent defaults lead to unrealistic loss scenarios

- We conclude that the independence assumption, or more generally, the i.i.d assumption for the individual default indicators $X_1, X_2, \ldots X_m$ implies that the fractional number of defaults in the portfolio $\frac{N_m}{m}$ converges to the constant *p* almost surely.
- It is an empirical fact, observed many times in the history, that defaults tend to cluster and ^{Nm}/_m have often values much bigger than p.
- Consequently, the empirical (i.e. observed) density for ^{Nm}/_m will have much more "fatter" tails compared with the binomial distribution.
- We will therefore next look at portfolio credit models that can produce more realistic loss scenarios, with densities for $\frac{N_m}{m}$ that have fat tails, which implies that $\frac{N_m}{m}$ does **not** converges to a constant with probability 1, when $m \to \infty$.

Before we continue this lecture, we need to introduce the concept of conditional expectations

- Let L^2 denote the space of all random variables X such that $\mathbb{E}\left[X^2\right] < \infty$
- Let Z be a random variable and let $L^2(Z) \subseteq L^2$ denote the space of all random variables Y such that Y = g(Z) for some function g and $Y \in L^2$
- Note that E [X] is the value μ that minimizes the quantity E [(X − μ)²]. Inspired by this, we define the conditional expectation E [X | Z] as follows:

Definition of conditional expectations

For a random variable Z, and for $X \in L^2$, the conditional expectation $\mathbb{E}[X | Z]$ is the random variable $Y \in L^2(Z)$ that minimizes $\mathbb{E}[(X - Y)^2]$.

 Intuitively, we can think of E [X | Z] as the orthogonal projection of X onto the space L²(Z), where the scalar product ⟨X, Y⟩ is defined as ⟨X, Y⟩ = E [XY].

Properties of conditional expectations

For a random variable Z it is possible to show the following properties

- **1.** If $X \in L^2$, then $\mathbb{E}\left[\mathbb{E}\left[X \mid Z\right]\right] = \mathbb{E}\left[X\right]$
- **2.** If $Y \in L^2(Z)$, then $\mathbb{E}[YX | Z] = Y\mathbb{E}[X | Z]$
- **3.** If $X \in L^2$, we define Var(X|Z) as

$$\operatorname{Var}(X|Z) = \mathbb{E}\left[X^2 \mid Z\right] - \mathbb{E}\left[X \mid Z\right]^2$$

and it holds that $Var(X) = \mathbb{E} \left[Var(X|Z)\right] + Var\left(\mathbb{E} \left[X \mid Z\right]\right)$.

Furthermore, for an event A, we can define the conditional probability $\mathbb{P}[A | Z]$ as

$$\mathbb{P}\left[A \,|\, Z\right] = \mathbb{E}\left[1_A \,|\, Z\right]$$

where 1_A is the indicator function for the event A (note that 1_A is a random variable). An example: if $X \in \{a, b\}$, let $A = \{X = a\}$, and we get that $\mathbb{P}[X = a | Z] = \mathbb{E}[1_{\{X=a\}} | Z]$.

The mixed binomial model

- The binomial model can be extended to **the mixed binomial model** which randomizes the default probability, allowing for stronger dependence.
- The mixed binomial model works as follows: Let Z be a random variable (discrete or continuous) and let $p(x) \in [0, 1]$ be a function such that the random variable p(Z) is well-defined.
- Let $X_1, X_2, ..., X_m$ be identically distributed random variables such that $X_i = 1$ if obligor *i* defaults before time T and $X_i = 0$ otherwise.
- Conditional on Z, the random variables X₁, X₂,... X_m are independent and each X_i have default probability p(Z), that is P [X_i = 1 | Z] = p(Z)
- The economic intuition behind this randomizing of the default probability p(Z) is that Z should represent some common background variable affecting all obligors in the portfolio.

• Let F(x) and \bar{p} be the distribution and mean of the random variable p(Z), that is,

$$F(x) = \mathbb{P}[p(Z) \le x] \text{ and } \mathbb{E}[p(Z)] = \overline{p}.$$
 (3)

If for example Z is a continuous random variable on ℝ with density f_Z(z) then p̄ is given by

$$\bar{p} = \mathbb{E}\left[p(Z)\right] = \int_{-\infty}^{\infty} p(z) f_Z(z) dz.$$
(4)

• Since
$$\mathbb{P}[X_i = 1 | Z] = p(Z)$$
 we get that $\mathbb{E}[X_i | Z] = p(Z)$, because $\mathbb{E}[X_i | Z] = 1 \cdot \mathbb{P}[X_i = 1 | Z] + 0 \cdot (1 - \mathbb{P}[X_i = 1 | Z]) = p(Z)$.

Note that E [X_i] = p̄ and thus p̄ = E [p(Z)] = P [X_i = 1] since
 P [X_i = 1] = E [X_i] = E [E [X_i | Z]] = E [p(Z)] = p̄

where the last equality is due to (3).

- One can show that (see in the lecture notes) $\operatorname{Var}(X_i) = \bar{p}(1-\bar{p})$ and $\operatorname{Cov}(X_i, X_j) = \mathbb{E}\left[p(Z)^2\right] - \bar{p}^2 = \operatorname{Var}(p(Z))$ (5)
- Next, letting all losses be the same and constant given by, say ℓ , then the total credit loss in the portfolio at time T, called L_m , is

$$L_m = \sum_{i=1}^m \ell X_i = \ell \sum_{i=1}^m X_i = \ell N_m \quad \text{where } N_m = \sum_{i=1}^m X_i$$

thus, N_m is the **number** of defaults in the portfolio up to time T

- Again, since $\mathbb{P}[L_m = k\ell] = \mathbb{P}[N_m = k]$, it is enough to study N_m .
- Since the random variables $X_1, X_2, \dots X_m$ now only are conditionally independent, given the outcome Z, we have

$$\mathbb{P}[N_m = k | Z] = \binom{m}{k} p(Z)^k (1 - p(Z))^{m-k}$$

• Hence,

$$\mathbb{P}[N_m = k] = \mathbb{E}\left[\mathbb{P}\left[N_m = k \mid Z\right]\right] = \mathbb{E}\left[\binom{m}{k} p(Z)^k (1 - p(Z))^k\right] \quad (6)$$

so if Z is a continuous random variable on $\mathbb R$ with density $f_Z(z)$ then

$$\mathbb{P}\left[N_m=k\right] = \int_{-\infty}^{\infty} \binom{m}{k} p(z)^k (1-p(z))^{m-k} f_Z(z) dz.$$
(7)

• Furthermore, because $X_1, X_2, \ldots X_m$ no longer are independent we have that

$$\operatorname{Var}(N_m) = \operatorname{Var}\left(\sum_{i=1}^m X_i\right) = \sum_{i=1}^m \operatorname{Var}(X_i) + \sum_{i=1}^m \sum_{j=1, j \neq i}^m \operatorname{Cov}(X_i, X_j) \quad (8)$$

and by homogeneity in the model we thus get

$$\operatorname{Var}(N_m) = m\operatorname{Var}(X_i) + m(m-1)\operatorname{Cov}(X_i, X_j). \tag{9}$$

• So inserting (5) in (9) we get that

$$\operatorname{Var}(N_m) = m\bar{p}(1-\bar{p}) + m(m-1) \left(\mathbb{E}\left[p(Z)^2\right] - \bar{p}^2 \right). \tag{10}$$

(CTH and Univ. of Gothenburg)

- Next, it is of interest to study how our portfolio will behave when $m \to \infty$, that is when the number of obligors in the portfolio goes to infinity.
- Recall that $Var(aX) = a^2 Var(X)$ so this and (10) imply that

$$\operatorname{Var}\left(\frac{N_m}{m}\right) = \frac{\operatorname{Var}(N_m)}{m^2} = \frac{\overline{p}(1-\overline{p})}{m} + \frac{(m-1)\left(\mathbb{E}\left[p(Z)^2\right] - \overline{p}^2\right)}{m}.$$

We therefore conclude that

$$\operatorname{Var}\left(\frac{N_m}{m}\right) \to \mathbb{E}\left[p(Z)^2\right] - \bar{p}^2 = \operatorname{Var}(p(Z)) \quad \text{as } m \to \infty$$
 (11)

Note that in the case when p(Z) is a constant, say p, so that p = p
 in the standard binomial loss model and

$$\mathbb{E}\left[p(Z)^2\right] - \bar{p}^2 = p^2 - p^2 = 0$$
 so $\operatorname{Var}\left(\frac{N_m}{m}\right) \to 0$ as $m \to \infty$

i.e. the fractional number of defaults in the portfolio converge to the constant $p = \bar{p}$ as portfolio size tend to infinity (law of large numbers.)

- So in the mixed binomial model, we see from (11) that the law of large numbers do not hold, i.e. Var (^{Nm}/_m) does not converge to 0.
- Consequently, the fractional number of defaults in the portfolio $\frac{N_m}{m}$ does not converge to a constant as $m \to \infty$.
- This is due to the fact that $X_1, X_2, ..., X_m$, are **not** independent. The dependence among $X_1, X_2, ..., X_m$ is created by Z.
- However, conditionally on Z, we have that the law of large numbers hold (because if we condition on Z, then $X_1, X_2, ..., X_m$ are i.i.d with default probability p(Z)), that is

given a "fixed" outcome of Z then
$$rac{N_m}{m} o p(Z)$$
 as $m o \infty$ (12)

 Since a.s convergence implies convergence in distribution (12) implies that for any x ∈ [0, 1] we have

$$\mathbb{P}\left[\frac{N_m}{m} \le x\right] \to \mathbb{P}\left[p(Z) \le x\right] \quad \text{when} \quad m \to \infty. \tag{13}$$

• Note that (13) can also be verified intuitive from (12) by making the following observation. From (12) we have that

$$\mathbb{P}\left[\left.\frac{N_m}{m} \le x \right| Z\right] \to \begin{cases} 0 & \text{if } p(Z) > x \\ 1 & \text{if } p(Z) \le x \end{cases} \quad \text{as } m \to \infty$$

that is,

$$\mathbb{P}\left[\frac{N_m}{m} \le x \mid Z\right] \to \mathbb{1}_{\{p(Z) \le x\}} \quad \text{as} \ \to \infty.$$
(14)

Next, recall that

$$\mathbb{P}\left[\frac{N_m}{m} \le x\right] = \mathbb{E}\left[\mathbb{P}\left[\frac{N_m}{m} \le x \mid Z\right]\right]$$
(15)

so (14) in (15) renders

$$\mathbb{P}\left[\frac{N_m}{m} \le x\right] \to \mathbb{E}\left[\mathbb{1}_{\{p(Z) \le x\}}\right] = \mathbb{P}\left[p(Z) \le x\right] = F(x) \quad \text{as } m \to \infty$$

where $F(x) = \mathbb{P}[p(Z) \le x]$, i.e. F(x) is the distribution function of the random variable p(Z).

Spring 2019 23 / 26

Large Portfolio Approximation (LPA)

Hence, from the above remarks we conclude the following important result:

Large Portfolio Approximation (LPA) for mixed binomial models

For large portfolios in a mixed binomial model, the distribution of the fractional number of defaults $\frac{N_m}{m}$ in the portfolio converges to the distribution of the random variable p(Z) as $m \to \infty$, that is for any $x \in [0, 1]$ we have

$$\mathbb{P}\left[\frac{N_m}{m} \le x\right] \to \mathbb{P}\left[p(Z) \le x\right] \quad \text{when} \quad m \to \infty.$$
(16)

The distribution $\mathbb{P}[p(Z) \le x]$ is called the Large Portfolio Approximation (LPA) to the distribution of $\frac{N_m}{m}$.

Examples of mixing distributions (next two lectures)

- Example 1: A mixed binomial model with p(Z) = Z where Z is a beta distribution, Z ~ Beta(a, b) and by definition of a beta distribution it holds that P[0 ≤ Z ≤ 1] = 1 so that p(Z) ∈ [0, 1].
- Example 2: Another possibility for mixing distribution p(Z) is to let p(Z) be a logit-normal distribution. This means that

$$p(Z) = \frac{1}{1 + \exp\left(-(\mu + \sigma Z)\right)}$$

where $\sigma > 0$ and Z is a standard normal. Note that $p(Z) \in [0, 1]$.

• **Example 3**: The mixed binomial model inspired by the Merton model (will be discussed coming lectures) with p(Z) given by

$$p(Z) = N\left(\frac{N^{-1}(\bar{p}) - \sqrt{\rho}Z}{\sqrt{1 - \rho}}\right)$$
(17)

where Z is a standard normal and N(x) is the distribution function of a standard normal distribution. Furthermore, $\rho \in [0, 1]$ and $\bar{p} = \mathbb{P}[X_i = 1]$. Note that $p(Z) \in [0, 1]$.

(CTH and Univ. of Gothenburg)

Financial Risk: Credit Risk, Lecture 1

 ▶
 ◄
 ■

 </t

▲□▶ ▲圖▶ ▲圖▶ ▲圖▶