

# Extra Exercises in Basic Probability for Financial Time Series

Andreas Petersson

April 12, 2016

## Easier exercises

**Ex. 1** — For  $t \in \mathbb{Z}$ , let  $X_t = Z_t + 0.5Z_{t-1}$  where  $Z_t$  is an iid sequence with mean 0 and variance  $\sigma^2$ .

- Compute the variance of the sample mean  $(X_1 + X_2 + X_3 + X_4)/4$ .
- Let  $Y_t = Z_t - 0.5Z_{t-1}$ . Compute the variance of the sample mean  $(Y_1 + Y_2 + Y_3 + Y_4)/4$  and compare to the previous part of the exercise.

**Ex. 2** — For  $t \in \mathbb{Z}$ , let  $X_t = X_0$  for all  $t \in \mathbb{Z}$  where  $X_0$  is a random variable with finite variance and zero mean. Is the process  $(X_t)_{t=1}^\infty$

- strictly stationary?
- weakly stationary?

**Ex. 3** — Let  $(X_t)_{t=1}^\infty$  be a sequence of random variables with (not necessarily identical) finite variance. For  $\alpha_1, \dots, \alpha_n \in \mathbb{R}$ , show that

$$\text{Var}\left(\sum_{i=1}^n \alpha_i X_i\right) = \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j \text{Cov}(X_i, X_j).$$

You may assume for simplicity that  $X_t$  all have zero mean.

**Ex. 4** — Let  $(X_t)_{t=1}^\infty$  be a sequence of zero-mean iid random variables with unit variance. Let  $Y_1 = X_1$  and let  $Y_t = Y_{t-1} + X_t$  for  $t \in \mathbb{N}$  with  $t > 1$ . Compute  $\gamma_Y(1, 2)$  and  $\gamma_Y(2, 3)$ . Is the process  $(Y_t)_{t=1}^\infty$  stationary?

**Ex. 5** — Let  $U \sim \mathcal{U}([0, 1])$ .

- Find the density function of  $S = U^2$ .
- Find the density function of  $T = -\log(U)/\lambda$ , where  $\lambda > 0$ .

## Harder exercises

**Ex. 6** — Let  $X = U_1 + U_2 + \dots + U_{12} - 6$  where, for  $i = 1, \dots, 12$ ,  $U_i \sim \mathcal{U}([0, 1])$ .

- Calculate the mean and variance of  $X$ .
- Use the CLT to show that  $X$  can be used to generate approximate standard normal random variables.

**Ex. 7** — Assume that the random variables  $X$  and  $Y$  are continuous with joint density  $f_{X,Y}(x, y)$  and finite means.

- a) Show that  $\mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[X]$ .
- b) Show that  $X \geq 0 \implies \mathbb{E}[X|Y] \geq 0$ .
- c) Assume that  $X$  and  $Y$  are independent and show that  $\mathbb{E}[X|Y] = \mathbb{E}[X]$ .

**Ex. 8** — Let  $B$  have the Bernoulli distribution with parameter  $p = 1/2$  and let  $X \sim \mathcal{N}(0, 1)$  be independent of  $B$ . Let  $Y = (2B - 1)X$ .

- a) Compute  $\mathbb{E}[Y]$ .
- b) Compute for  $P(Y \leq y)$  for  $y \in \mathbb{R}$ . What is the distribution of  $Y$ ? *Hint: Note that  $P(Y \leq y) = P(Y \leq y, B = 0) + P(Y \leq y, B = 1)$ .*
- c) Compute  $\text{Cov}(X, Y)$ . Are  $X$  and  $Y$  independent?

**Ex. 9** — Let  $X = \exp(Z)$  where  $Z \sim \mathcal{N}(0, 1)$  (this is called a lognormal random variable).

- a) Find the density function of  $X$
- b) Use this density to find the mean and variance of  $X$ .

**Ex. 10** — Let  $\mathcal{X}$  be the family of all (real-valued) random variables with finite variance on the probability space  $(\Omega, \mathcal{A}, P)$ . Show that  $\mathcal{X}$  is a vector space. *Hint: Use the Cauchy-Schwarz inequality  $|\mathbb{E}[XY]|^2 \leq \mathbb{E}[X^2] \mathbb{E}[Y^2]$ . You may assume that if  $X$  and  $Y$  are random variables (i.e. measurable), then so are  $\alpha X + \beta Y$  for  $\alpha, \beta \in \mathbb{R}$*