

Numerical Methods for Stochastic ODEs

Stig Larsson

<http://www.math.chalmers.se/~stig>

1. LECTURE 13

The main textbook is Klebaner [3]. For the computer exercises we refer to Higham [2]. It is very important to be able to do computer simulations with stochastic ODEs, so I do recommend that you do some of these computer exercises. In these lectures I will present the theoretical background. Course homepage: [link](#)

1.1. **Preparations.** Let X be a strong solution of the scalar Itô SDE:

$$(1) \quad \begin{aligned} dX(t) &= \mu(X(t), t) dt + \sigma(X(t), t) dB(t), \quad 0 \leq t \leq T, \\ X(0) &= X_0. \end{aligned}$$

This means, see Klebaner Definition 5.1, that X solves the integral equation

$$(2) \quad X(t) = X_0 + \int_0^t \mu(X(s), s) ds + \int_0^t \sigma(X(s), s) dB(s), \quad t \in [0, T].$$

Here B is Brownian motion and the last term is the Itô integral.

We assume that X_0 is a random variable and the coefficients are deterministic functions,

$$(3) \quad \begin{aligned} \mu &: \mathbf{R} \times [0, T] \rightarrow \mathbf{R}, \\ \sigma &: \mathbf{R} \times [0, T] \rightarrow \mathbf{R}, \end{aligned}$$

that are continuous and satisfy a global Lipschitz condition with respect to x ,

$$(4) \quad \begin{aligned} |\mu(x, t) - \mu(y, t)| &\leq L|x - y|, \\ |\sigma(x, t) - \sigma(y, t)| &\leq L|x - y|, \end{aligned} \quad \forall x, y \in \mathbf{R}, \quad t \in [0, T],$$

and a linear growth bound,

$$(5) \quad \begin{aligned} |\mu(x, t)| &\leq L(1 + |x|), \\ |\sigma(x, t)| &\leq L(1 + |x|), \end{aligned} \quad \forall x \in \mathbf{R}, \quad t \in [0, T].$$

Remark 1. The Lipschitz condition (4) is called *global* because it holds for all $x, y \in \mathbf{R}$ with the same constant L . Klebaner Theorem 5.7 assumes only a *local* Lipschitz condition, where the Lipschitz constant may depend on the size of x, y . We use a global condition in order to make the presentation simpler. Later on we will also assume a Lipschitz condition with respect to t , see (22). (There is a mistake in Theorem 5.7: the constants in (5.36) and (5.37) cannot be the same because the first one depends on N , $K = K_N$, while the second one is a global constant, independent of N .)

Remark 2. Note that (5) follows from (4) because

$$(6) \quad |\mu(x, t)| \leq |\mu(0, t)| + |\mu(x, t) - \mu(0, t)| \leq \max_{t \in [0, T]} |\mu(0, t)| + L|x - 0| \leq \tilde{L}(1 + |x|)$$

with a new constant \tilde{L} . This is because (4) is global.

¹Revised: 2008-10-28

Remark 3. Systems of SDE of the form

$$(7) \quad dX_i = \mu_i(X, t) dt + \sum_{j=1}^m \sigma_{ij}(X, t) dB_j(t), \quad i = 1, \dots, n,$$

can be written in the form (1), and analyzed in the same way, with

$$\mu : \mathbf{R}^n \times [0, T] \rightarrow \mathbf{R}^n, \quad \sigma : \mathbf{R}^n \times [0, T] \rightarrow \mathbf{R}^{n \times m},$$

and B an m -dimensional Brownian motion.

We quote Theorem 5.7 from Klebaner.

Theorem 1 (Existence and uniqueness). *If X_0 is independent of $(B(t), 0 \leq t \leq T)$ and $\mathbf{E}(|X_0|^2) < \infty$, then (1) has a unique strong solution X and*

$$(8) \quad \mathbf{E} \left(\sup_{0 \leq t \leq T} |X(t)|^2 \right) \leq C(1 + \mathbf{E}(|X_0|^2)),$$

where $C = C(L, T)$.

As a warm-up for the error analysis of numerical methods, we begin by proving a stability result. We will use this proof technique several times later. The bound (8) is proved in a similar way.

We need Gronwall's lemma.

Lemma 1 (Gronwall). *Let A, B be constants with $B \geq 0$. If*

$$\phi(t) \leq A + B \int_0^t \phi(s) ds, \quad t \in [0, T],$$

then

$$\phi(t) \leq Ae^{Bt}, \quad t \in [0, T].$$

Proof. $u(t) := A + \int_0^t \phi(s) ds$, $\phi(t) \leq u(t)$, $u' = B\phi \leq Bu$, $u(t) \leq u(0)e^{Bt} = Ae^{Bt}$. \square

We also need Doob's L_p martingale inequality, see Klebaner (7.37) p. 201.

Theorem 2 (Doob's inequality). *If Y is a martingale, then*

$$(9) \quad \mathbf{E} \left(\max_{0 \leq t \leq T} |Y(t)|^p \right) \leq \left(\frac{p}{p-1} \right)^p \mathbf{E}(|Y(T)|^p), \quad 1 < p < \infty.$$

With the L_p -norm $\|X\|_{L_p} = (\mathbf{E}(|X|^p))^{1/p} = (\int_{\Omega} |X(\omega)|^p d\mathbf{P}(\omega))^{1/p}$ this can also be written

$$\left\| \max_{0 \leq t \leq T} |Y(t)| \right\|_{L_p} \leq \frac{p}{p-1} \|Y(T)\|_{L_p}.$$

This gives a bound for the norm of the whole history in terms of the norm of the final value.

Finally we recall the isometry of the Itô integral, Klebaner (4.12),

$$(10) \quad \mathbf{E} \left(\left| \int_0^t f(s) dB(s) \right|^2 \right) = \int_0^t \mathbf{E}(|f(s)|^2) ds.$$

We are now ready to prove stability with respect to perturbation of the initial value.

Theorem 3 (Stability). *Let \hat{X} be another solution of (2) with the same μ, σ, B but a different initial value \hat{X}_0 . Then*

$$(11) \quad \mathbf{E} \left(\max_{0 \leq t \leq T} |\hat{X}(t) - X(t)|^2 \right) \leq C \mathbf{E}(|\hat{X}_0 - X_0|^2),$$

where $C = C(L, T)$.

Note that this implies uniqueness: two solutions with the same data μ, σ, B, X_0 must be equal.

Proof. We will use Gronwall's lemma with $\phi(t) = \mathbf{E}\left(\max_{0 \leq s \leq t} |\hat{X}(s) - X(s)|^2\right)$. By (2), writing $\hat{\mu} = \mu(\hat{X}(z), z)$, $\mu = \mu(X(z), z)$, etc, and using the inequality $(a+b+c)^2 \leq 3(a^2+b^2+c^2)$, we get

$$\begin{aligned} \phi(t) &= \mathbf{E}\left(\max_{0 \leq s \leq t} |\hat{X}(s) - X(s)|^2\right) \\ &= \mathbf{E}\left(\max_{0 \leq s \leq t} \left| \hat{X}_0 - X_0 + \int_0^s (\mu(\hat{X}(z), z) - \mu(X(z), z)) dz \right. \right. \\ &\quad \left. \left. + \int_0^s (\sigma(\hat{X}(z), z) - \sigma(X(z), z)) dB(z) \right|^2\right) \\ &\leq 3\left\{ \mathbf{E}\left(|\hat{X}_0 - X_0|^2\right) + \mathbf{E}\left(\max_{0 \leq s \leq t} \left| \int_0^s (\hat{\mu} - \mu) dz \right|^2\right) \right. \\ &\quad \left. + \mathbf{E}\left(\max_{0 \leq s \leq t} \left| \int_0^s (\hat{\sigma} - \sigma) dB \right|^2\right) \right\}. \end{aligned}$$

Here, by the Cauchy-Schwartz inequality and (4),

$$\begin{aligned} \mathbf{E}\left(\max_{0 \leq s \leq t} \left| \int_0^s (\hat{\mu} - \mu) dz \right|^2\right) &\leq \mathbf{E}\left(\max_{0 \leq s \leq t} \left\{ \int_0^s 1^2 dz \int_0^s |\hat{\mu} - \mu|^2 dz \right\}\right) \\ &= \mathbf{E}\left(t \int_0^t |\hat{\mu} - \mu|^2 dz\right) = t \int_0^t \mathbf{E}(|\hat{\mu} - \mu|^2) ds \\ &\leq L^2 T \int_0^t \mathbf{E}(|\hat{X}(s) - X(s)|^2) ds \\ &\leq L^2 T \int_0^t \mathbf{E}\left(\max_{0 \leq z \leq s} |\hat{X}(z) - X(z)|^2\right) ds. \end{aligned}$$

For the other term we use Doob's inequality (9) with $p = 2$ and

$$Y(t) = \int_0^t (\sigma(\hat{X}(z), z) - \sigma(X(z), z)) dB(z),$$

which is a martingale by Theorem 4.7 in Klebaner. Using also (10) and (4) we get

$$\begin{aligned} \mathbf{E}\left(\max_{0 \leq s \leq t} \left| \int_0^s (\hat{\sigma} - \sigma) dB \right|^2\right) &\leq 4\mathbf{E}\left(\left| \int_0^t (\hat{\sigma} - \sigma) dB \right|^2\right) \\ &= 4 \int_0^t \mathbf{E}(|\hat{\sigma} - \sigma|^2) ds \\ &\leq 4L^2 \int_0^t \mathbf{E}(|\hat{X}(s) - X(s)|^2) ds \\ &\leq 4L^2 \int_0^t \mathbf{E}\left(\max_{0 \leq z \leq s} |\hat{X}(z) - X(z)|^2\right) ds. \end{aligned}$$

We now have

$$\mathbf{E}\left(\max_{0 \leq s \leq t} |\hat{X}(s) - X(s)|^2\right) \leq 3\mathbf{E}(|\hat{X}_0 - X_0|^2) + C \int_0^t \mathbf{E}\left(\max_{0 \leq z \leq s} |\hat{X}(z) - X(z)|^2\right) ds$$

with $C = 3L^2(T + 4)$, or

$$\phi(t) \leq 3\mathbf{E}(|\hat{X}_0 - X_0|^2) + C \int_0^t \phi(s) ds, \quad t \in [0, T],$$

where $\phi(t) = \mathbf{E}(\max_{0 \leq s \leq t} |\hat{X}(s) - X(s)|^2)$ and Gronwall's lemma completes the proof. \square

Exercise 1. Without using Doob's inequality and with $\phi(t) = \mathbf{E}(|\hat{X}(t) - X(t)|^2)$ we get the weaker result

$$\mathbf{E}(|\hat{X}(t) - X(t)|^2) \leq C\mathbf{E}(|\hat{X}_0 - X_0|^2), \quad t \in [0, T],$$

and hence (Prove this!)

$$(12) \quad \max_{0 \leq t \leq T} \mathbf{E}(|\hat{X}(t) - X(t)|^2) \leq C\mathbf{E}(|\hat{X}_0 - X_0|^2).$$

Note that the maximum is now outside the expected value and that (11) implies (12).

Exercise 2. Prove the estimate (8).

We also need a kind of Hölder condition.

Theorem 4. *Under the assumptions of Theorem 1 we have*

$$(13) \quad \mathbf{E}(|X(t) - X(s)|^2) \leq C(1 + \mathbf{E}(|X_0|^2))|t - s|, \quad \forall t, s \in [0, T],$$

where $C = C(L, T)$.

Proof. We may assume that $s \leq t$. Working as in the previous proof, but without using Doob's inequality, we then get

$$\begin{aligned} \mathbf{E}(|X(t) - X(s)|^2) &= \mathbf{E}\left(\left|\int_s^t \mu(X(z), z) dz + \int_s^t \sigma(X(z), z) dB(z)\right|^2\right) \\ &\leq 2 \underbrace{(t-s)}_{\leq T} \int_s^t \mathbf{E}(|\mu|^2) dz + 2 \int_s^t \mathbf{E}(|\sigma|^2) dz \quad \{\text{by (5)}\} \\ &\leq 2L^2(T+1) \int_s^t \mathbf{E}((1 + |X(z)|)^2) dz \\ &\leq 4L^2(T+1) \int_s^t \{1 + \mathbf{E}(|X(z)|^2)\} dz \quad \{\text{by (8)}\} \\ &\leq C(L, T) \int_s^t \{1 + C(L, T)(1 + \mathbf{E}(|X_0|^2))\} dz \\ &\leq C(L, T)(t-s)(1 + \mathbf{E}(|X_0|^2)). \end{aligned}$$

\square

Remark 4. Note that this means, with $\|X_0\|_{L_2} = \mathbf{E}(|X_0|^2)^{1/2} \leq M$ and $C = C(M, L, T)$,

$$(14) \quad \|X(t) - X(s)\|_{L_2} = \left(\mathbf{E}(|X(t) - X(s)|^2)\right)^{1/2} \leq C|t - s|^{1/2}.$$

This is a Hölder condition with exponent $\frac{1}{2}$. By inspection of the previous proof we see that in the deterministic case, $\sigma = 0$, we have a stronger result, a Lipschitz condition,

$$(15) \quad |X(t) - X(s)| \leq C|t - s|.$$

1.2. Strong convergence of Euler's method. We introduce a mesh

$$(16) \quad 0 = t_0 < t_1 < \dots < t_n < t_{n+1} < \dots < t_N = T, \quad h_n = t_{n+1} - t_n, \quad h = \max h_n \leq 1,$$

and note that X satisfies

$$(17) \quad X(t_{n+1}) = X(t_n) + \int_{t_n}^{t_{n+1}} \mu(X(s), s) ds + \int_{t_n}^{t_{n+1}} \sigma(X(s), s) dB(s).$$

This motivates the Euler (or Euler-Maruyama) method, which defines $\{Y_n\}_{n=0}^N$ by

$$(18) \quad \begin{aligned} Y_0 &\approx X_0, \\ Y_{n+1} &= Y_n + \mu(Y_n, t_n) \int_{t_n}^{t_{n+1}} ds + \sigma(Y_n, t_n) \int_{t_n}^{t_{n+1}} dB(s). \end{aligned}$$

Since $\int_{t_n}^{t_{n+1}} ds = h_n$, $\int_{t_n}^{t_{n+1}} dB(s) = B(t_{n+1}) - B(t_n) = \Delta B_n$, this can also be written

$$(19) \quad \begin{aligned} Y_0 &\approx X_0, \\ Y_{n+1} &= Y_n + \mu(Y_n, t_n)h_n + \sigma(Y_n, t_n)\Delta B_n. \end{aligned}$$

For the practical aspects concerning the computation in a Matlab environment we refer to [2]. Here we only note that $\Delta B_n = \sqrt{h_n}\xi_n$, where the $\xi_n \in N(0, 1)$ are independent Gaussian random variables with mean zero and variance 1 which can be simulated on the computer by a random number generator.

Although we only compute the node values Y_n it is convenient for our proofs to define $Y(t)$ for $t \in [t_n, t_{n+1})$ by

$$(20) \quad Y(t) = Y_n + \mu(Y_n, t_n) \int_{t_n}^t ds + \sigma(Y_n, t_n) \int_{t_n}^t dB(s), \quad t \in [t_n, t_{n+1}).$$

Then Y is continuous on $[0, T]$, $Y(t_n) = Y_n$ for all n , and

$$(21) \quad \begin{aligned} Y(t) &= Y_0 + \int_0^t \bar{\mu}(s) ds + \int_0^t \bar{\sigma}(s) dB(s), \quad 0 \leq t \leq T, \\ \bar{\mu}(s) &= \mu(Y_n, t_n), \quad \bar{\sigma}(s) = \sigma(Y_n, t_n), \quad s \in (t_n, t_{n+1}). \end{aligned}$$

We need a Lipschitz condition with respect to t :

$$(22) \quad \begin{aligned} |\mu(x, t) - \mu(x, s)| &\leq L|t - s|, \\ |\sigma(x, t) - \sigma(x, s)| &\leq L|t - s|, \end{aligned} \quad \forall x \in \mathbf{R}, \quad t, s \in [0, T].$$

Theorem 5 (Strong convergence). *If*

$$(23) \quad \mathbf{E}(|Y_0 - X_0|^2) \leq Kh,$$

and

$$(24) \quad \mathbf{E}(|X_0|^2) \leq M,$$

then

$$(25) \quad \mathbf{E}\left(\max_{0 \leq t \leq T} |Y(t) - X(t)|^2\right) \leq Ch$$

with $C = C(L, K, M, T)$. In particular,

$$(26) \quad \mathbf{E}(|Y_n - X(t_n)|) \leq Ch^{1/2}, \quad n = 0, \dots, N.$$

Proof. We first prove that (25) implies (26). We use the Cauchy-Schwartz inequality,

$$|\mathbf{E}(fg)| \leq \sqrt{\mathbf{E}(f^2)}\sqrt{\mathbf{E}(g^2)},$$

with $f = 1$ and $g = |Y(t_n) - X(t_n)|$:

$$\mathbf{E}(|Y_n - X(t_n)|) \leq \sqrt{\mathbf{E}(|Y(t_n) - X(t_n)|^2)} \leq \sqrt{\mathbf{E}\left(\max_{0 \leq t \leq T} |Y(t) - X(t)|^2\right)} \leq Ch^{1/2}.$$

We now prove (25). Let $0 \leq s \leq t \leq T$. As in the proof of Theorem 3, using (2), (21):

$$\begin{aligned} \mathbf{E}\left(\max_{0 \leq s \leq t} |Y(s) - X(s)|^2\right) &= \left\{ \text{with } \mu(z) = \mu(X(z), z), \sigma(z) = \sigma(X(z), z) \right\} \\ &= \mathbf{E}\left(\max_{0 \leq s \leq t} \left\{ \left| Y_0 - X_0 + \int_0^s (\bar{\mu}(z) - \mu(z)) dz + \int_0^s (\bar{\sigma}(z) - \sigma(z)) dB(z) \right|^2 \right\}\right) \\ &\leq 3 \left\{ \mathbf{E}(|Y_0 - X_0|^2) + T \int_0^t \mathbf{E}(|\bar{\mu} - \mu|^2) ds + 4 \int_0^t \mathbf{E}(|\bar{\sigma} - \sigma|^2) ds \right\}. \end{aligned}$$

We split $\bar{\mu} - \mu$ into two parts and use the Lipschitz conditions (4) and (22),

$$\begin{aligned} |\bar{\mu}(s) - \mu(s)| &= |\mu(Y(t_n), t_n) - \mu(X(s), s)| \\ &\leq |\mu(Y(t_n), t_n) - \mu(X(s), t_n)| + |\mu(X(s), t_n) - \mu(X(s), s)| \\ &\leq L(|Y(t_n) - X(s)| + s - t_n) \\ &\leq L(|Y(t_n) - X(t_n)| + |X(t_n) - X(s)| + h_n) \\ &\leq L\left(\max_{0 \leq z \leq s} |Y(z) - X(z)| + |X(t_n) - X(s)| + h_n\right), \quad \text{for } s \in (t_n, t_{n+1}). \end{aligned}$$

Using also (14) and $h_n \leq 1$ we get

$$\begin{aligned} \mathbf{E}(|\bar{\mu}(s) - \mu(s)|^2) &\leq 3L^2\left(\mathbf{E}\left(\max_{0 \leq z \leq s} |Y(z) - X(z)|^2\right) + \mathbf{E}(|X(t_n) - X(s)|^2) + h_n^2\right) \\ &\leq 3L^2\left(\mathbf{E}\left(\max_{0 \leq z \leq s} |Y(z) - X(z)|^2\right) + C(s - t_n) + h_n^2\right) \\ &\leq C\left(\mathbf{E}\left(\max_{0 \leq z \leq s} |Y(z) - X(z)|^2\right) + h_n\right), \quad \text{for } s \in (t_n, t_{n+1}), \end{aligned}$$

with $C = C(M, L, T)$. The difference $\bar{\sigma} - \sigma$ is estimated in the same way.

Suppose that t belongs to the m -th mesh interval, $t \in (t_m, t_{m+1}]$, and denote $\tilde{t}_{n+1} = t_{n+1} \wedge t$ (the minimum of t and t_{n+1}). Then, since $h_n \leq h$,

$$\begin{aligned} \int_0^t \mathbf{E}(|\bar{\mu}(s) - \mu(s)|^2) ds &= \sum_{n=0}^m \int_{t_n}^{\tilde{t}_{n+1}} \mathbf{E}(|\bar{\mu}(s) - \mu(s)|^2) ds \\ &\leq C \sum_{n=0}^m \int_{t_n}^{\tilde{t}_{n+1}} \left(\mathbf{E}\left(\max_{0 \leq z \leq s} |Y(z) - X(z)|^2\right) + h_n \right) ds \\ &\leq C \int_0^t \mathbf{E}\left(\max_{0 \leq z \leq s} |Y(z) - X(z)|^2\right) ds + Ch. \end{aligned}$$

We get the same bound for the σ -term. Therefore

$$\phi(t) \leq Ch + C \int_0^t \phi(s) ds, \quad t \in [0, T],$$

with $C = C(L, K, M, T)$, $\phi(t) = \mathbf{E}\left(\max_{0 \leq s \leq t} |Y(s) - X(s)|^2\right)$. Gronwall's lemma completes the proof. \square

We have now proved *strong convergence*, or *pathwise convergence*, of order $h^{1/2}$:

$$(27) \quad \left\| \max_{0 \leq t \leq T} |Y(t) - X(t)| \right\|_{L_2} = \sqrt{\mathbf{E}\left(\max_{0 \leq t \leq T} |Y(t) - X(t)|^2\right)} \leq Ch^{1/2}.$$

Note that, due to the use of Gronwall's inequality, the constant in (27) grows exponentially with T , $C = \exp(C(L, K, M, T)T)$. By inspection of the previous proof we see that in the deterministic case, $\sigma = 0$, we have convergence of order h :

$$\max_{0 \leq t \leq T} |Y(t) - X(t)| \leq Ch.$$

This is the classical result for Euler's method for deterministic ODEs.

In the next lecture we shall study *weak convergence* and prove that

$$\left| \mathbf{E}(g(Y_n) - g(X(t_n))) \right| \leq Ch$$

for all smooth functions g . Thus the weak convergence order is h .

We shall also derive a higher order method: Milstein's method.

The presentation of some of the material in Lecture 13 was inspired by [1]. For more details see [4].

2. LECTURE 14

Recall the strong solution

$$(28) \quad X(t) = X_0 + \int_0^t \mu(X(s), s) ds + \int_0^t \sigma(X(s), s) dB(s), \quad t \in [0, T],$$

and Euler's method

$$(29) \quad Y(t) = Y_0 + \int_0^t \bar{\mu}(s) ds + \int_0^t \bar{\sigma}(s) dB(s), \quad t \in [0, T],$$

where

$$(30) \quad \bar{\mu}(s) = \mu(Y(t_n), t_n), \quad \bar{\sigma}(s) = \sigma(Y(t_n), t_n), \quad s \in (t_n, t_{n+1}),$$

are piecewise constant, "frozen", functions.

We have proved *strong convergence*, or *pathwise convergence*:

$$(31) \quad \left\| \max_{0 \leq t \leq T} |Y(t) - X(t)| \right\|_{L_2} = \sqrt{\mathbf{E} \left(\max_{0 \leq t \leq T} |Y(t) - X(t)|^2 \right)} \leq Ch^{1/2}.$$

We are often not interested in individual paths but we would like to compute the expected value of some quantity that depends on $X(t)$, i.e., we want to compute $\mathbf{E}(g(X(t)))$ for some function g .

We shall prove that

$$\left| \mathbf{E}(g(Y(t)) - g(X(t))) \right| \leq Ch$$

for all smooth functions g . This is called *weak convergence*. Thus the weak convergence order is h . We will use Kolmogorov's backward equation in the proof.

We shall also derive the Itô-Taylor expansion and use it to obtain a numerical method of higher order: Milstein's method.

All of this rests on Itô's formula.

2.1. Kolmogorov's backward equation. We recall Itô's formula for functions of x, t (see Klebaner, Theorem 4.8 or Theorem 6.1). If X satisfies

$$dX(t) = \mu(X(t), t) dt + \sigma(X(t), t) dB(t)$$

and $u = u(x, t)$ is twice differentiable in x and once in t , then (denoting the partial derivatives $u_x = \frac{\partial u}{\partial x}$, $u_{xx} = \frac{\partial^2 u}{\partial x^2}$, $u_t = \frac{\partial u}{\partial t}$)

$$(32) \quad du(X(t), t) = \left(u_t(X(t), t) + \mu(X(t), t) u_x(X(t), t) + \frac{1}{2} \sigma^2(X(t), t) u_{xx}(X(t), t) \right) dt + \sigma(X(t), t) u_x(X(t), t) dB(t).$$

This really means

$$(33) \quad u(X(t), t) - u(X(s), s) = \int_s^t (u_t + \mu u_x + \frac{1}{2} \sigma^2 u_{xx}) dz + \int_s^t \sigma u_x dB(z), \quad 0 \leq s \leq t,$$

where the integrands are evaluated at $(X(z), z)$.

Theorem 6 (Kolmogorov's backward equation, Klebaner Theorem 6.6). *Let*

$$(34) \quad dX(t) = \mu(X(t), t) dt + \sigma(X(t), t) dB(t), \quad t \in [0, T],$$

and let $u(x, t)$ be a solution of

$$(35) \quad \begin{aligned} u_t(x, t) + \mu(x, t) u_x(x, t) + \frac{1}{2} \sigma^2(x, t) u_{xx}(x, t) &= 0, \quad x \in \mathbf{R}, \quad t < T, \\ u(x, T) &= g(x). \end{aligned}$$

Then

$$(36) \quad u(x, t) = \mathbf{E}(g(X(T)) \mid X(t) = x).$$

Note: Under appropriate assumptions we can prove that (35) has a unique solution. Then (36) is the only solution.

The notation in (36) is somewhat confusing but it is the traditional one. To explain what it means, we let $Z(s) = Z(s; x, t)$ denote the unique strong solution of

$$\begin{aligned} dZ(s) &= \mu(Z(s), s) ds + \sigma(Z(s), s) dB(s), \quad t \leq s \leq T, \\ Z(t) &= x. \end{aligned}$$

In particular, the solution of (1) is $X(t) = Z(t; X_0, 0)$. Now (36) can be written

$$(37) \quad u(x, t) = \mathbf{E}(g(Z(T; x, t))).$$

Note that the variable x in (36), (37) is deterministic, while X_0 in (1) may be a random variable.

Proof. Itô's formula applied to $u(Z(s; x, t), s)$ and (35) give

$$\begin{aligned} u(Z(T; x, t), T) - u(Z(t; x, t), t) &= \int_t^T (u_t + \mu u_x + \frac{1}{2} \sigma^2 u_{xx}) ds + \int_t^T \sigma u_x dB \\ &= \int_t^T \sigma u_x dB. \end{aligned}$$

The expected value of the Itô integral is zero, so that

$$\begin{aligned} &\mathbf{E}(u(Z(T; x, t), T)) - \mathbf{E}(u(Z(t; x, t), t)) \\ &= \mathbf{E}\left(\int_t^T \sigma(Z(s; x, t), s) u_x(Z(s; x, t), s) dB(s)\right) = 0. \end{aligned}$$

Since $u(Z(T; x, t), T) = g(Z(T; x, t))$ by (35) and $\mathbf{E}(u(Z(t; x, t), t)) = \mathbf{E}(u(x, t)) = u(x, t)$, we get

$$\mathbf{E}(g(Z(T; x, t), T)) = u(x, t),$$

which is (37). \square

According to Definition 5.14 and Theorem 5.15 in Klebaner, there is a unique function $p(y, s, x, t)$, $t < s \leq T$, $x, y \in \mathbf{R}$, (a fundamental solution) such that the unique solution of (35) is given by

$$(38) \quad u(x, t) = \int_{-\infty}^{\infty} p(y, T, x, t) g(y) dy, \quad x \in \mathbf{R}, \quad t < T.$$

The function $(y, s) \mapsto p(y, s, x, t)$ (with x, t fixed) satisfies the forward equation, see Klebaner (5.5a),

$$(39) \quad \frac{\partial p}{\partial s} + \frac{\partial}{\partial y} (\mu(y, s) p) - \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(y, s) p) = 0, \quad y \in \mathbf{R}, \quad t < s \leq T.$$

The initial value at $s = t$ is the Dirac measure at x :

$$(40) \quad p(y, t, x, t) = \delta_x(y), \quad y \in \mathbf{R}.$$

In the simplified case when $\mu = 0$, $\sigma^2 = 1$, (39), (40) become

$$(41) \quad \begin{aligned} p_s - \frac{1}{2} p_{yy} &= 0, & y \in \mathbf{R}, \quad s > t, \\ p(y, t, x, t) &= \delta_x(y), & y \in \mathbf{R}, \end{aligned}$$

which is the forward heat equation with solution

$$p(y, s, x, t) = \frac{1}{\sqrt{2\pi(s-t)}} e^{-\frac{(x-y)^2}{2(s-t)}}, \quad s > t, \quad x, y \in \mathbf{R},$$

the Gauss kernel. This is proved by taking the Fourier transform of (41) with respect to y . Then (38) becomes

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi(T-t)}} e^{-\frac{(x-y)^2}{2(T-t)}} g(y) dy.$$

Exercise 3. Note that the partial differential equations in (39) and (35) may be written as $Lp = 0$ and $L^*u = 0$, respectively, where $L = \frac{\partial}{\partial t} + \frac{\partial}{\partial x}(\mu(x, t)\cdot) - \frac{1}{2}\frac{\partial^2}{\partial x^2}(\sigma^2(x, t)\cdot)$ and $L^* = -\frac{\partial}{\partial t} - \mu(x, t)\frac{\partial}{\partial x} - \frac{1}{2}\sigma^2(x, t)\frac{\partial^2}{\partial x^2}$. Show that the operators L and L^* are formally adjoint in the sense that

$$(42) \quad \int_0^T \int_{\mathbf{R}} (L\phi)\psi dx dt = \int_0^T \int_{\mathbf{R}} \phi(L^*\psi) dx dt \quad \forall \phi, \psi \in C_0^\infty(\mathbf{R} \times (0, T)).$$

Exercise 4. Show that (38) follows from (35) and (39), (40). Hint: multiply (35) in the form $0 = u_s + \mu u_y + \frac{1}{2}\sigma^2 u_{yy}$ by $p(y, s, x, t)$, integrate $\int_t^T \int_{\mathbf{R}} \cdots dy ds$, integrate by parts and use (39), (40).

2.2. Weak convergence of Euler's method. In the following theorem we make rather strong assumptions on the coefficients and the initial value in order to make the proof simpler. These can be relaxed.

Theorem 7 (Weak convergence). *Assume that μ, σ, g are sufficiently smooth and decay sufficiently fast as $|x| \rightarrow \infty$. Let X and Y be solutions of (28) and (29) with $Y_0 = X_0$ a deterministic variable. Then*

$$(43) \quad |\mathbf{E}(g(Y(T))) - \mathbf{E}(g(X(T)))| \leq Ch.$$

Proof. Let $u = u(x, t)$ be the solution of the Kolmogorov backward equation (35). Then by (36), (37) we have

$$(44) \quad u(x, t) = \mathbf{E}(g(Z(T; x, t))),$$

and, in particular, since X_0 is deterministic,

$$(45) \quad u(X_0, 0) = \mathbf{E}(g(Z(T; X_0, 0))) = \mathbf{E}(g(X(T))),$$

Since Y satisfies (28), Itô's formula (33) gives

$$\begin{aligned} & u(Y(T), T) - u(Y(0), 0) \\ &= \int_0^T \left(u_t(Y(t), t) + \bar{\mu}(t)u_x(Y(t), t) + \frac{1}{2}\bar{\sigma}^2(t)u_{xx}(Y(t), t) \right) dt \\ & \quad + \int_0^T \bar{\sigma}(t)u_x(Y(t), t) dB(t). \end{aligned}$$

From (35) we get

$$u_t(Y(t), t) = -\mu(Y(t), t)u_x(Y(t), t) - \frac{1}{2}\sigma^2(Y(t), t)u_{xx}(Y(t), t),$$

so that

$$u(Y(T), T) - u(Y(0), 0) = \int_0^T \left((\bar{\mu} - \mu)u_x + \frac{1}{2}(\bar{\sigma}^2 - \sigma^2)u_{xx} \right) dt + \int_0^T \bar{\sigma}u_x dB(t),$$

where the integrands are evaluated at $(Y(t), t)$. Take expectation here, use $u(Y(T), T) = g(Y(T))$ from (35), use $Y_0 = X_0$ and (45) to get

$$\mathbf{E}(u(Y(0), 0)) = \mathbf{E}(u(Y_0, 0)) = \mathbf{E}(u(X_0, 0)) = \mathbf{E}(g(X(T))),$$

and use $\mathbf{E}(\int_0^T \bar{\sigma} u_x dB) = 0$. We get

$$\begin{aligned} \mathbf{E}(g(Y(T))) - \mathbf{E}(g(X(T))) &= \int_0^T \left\{ \mathbf{E}((\bar{\mu} - \mu)u_x) + \frac{1}{2}\mathbf{E}((\bar{\sigma}^2 - \sigma^2)u_{xx}) \right\} dt \\ &= \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \left\{ \mathbf{E}((\bar{\mu} - \mu)u_x) + \frac{1}{2}\mathbf{E}((\bar{\sigma}^2 - \sigma^2)u_{xx}) \right\} dt. \end{aligned}$$

We shall show

$$(46) \quad \sup_{t \in (t_n, t_{n+1})} \left| \mathbf{E}((\bar{\mu}(t) - \mu(Y(t), t))u_x(Y(t), t)) \right| \leq Ch_n,$$

$$(47) \quad \sup_{t \in (t_n, t_{n+1})} \frac{1}{2} \left| \mathbf{E}((\bar{\sigma}^2(t) - \sigma^2(Y(t), t))u_{xx}(Y(t), t)) \right| \leq Ch_n,$$

which implies the desired result.

To prove (46) we define

$$v(x, t) = (\mu(Y_n, t_n) - \mu(x, t))u_x(x, t),$$

and note that (46) means (recall that $\bar{\mu}(t) = \mu(Y(t_n), t_n)$ for $t \in (t_n, t_{n+1})$)

$$\left| \mathbf{E}(v(Y(t), t)) \right| \leq Ch_n, \quad t \in (t_n, t_{n+1}).$$

Since Y satisfies (28), Itô's formula (33) gives

$$\begin{aligned} v(Y(t), t) - v(Y(t_n), t_n) &= \int_{t_n}^t \left(v_t(Y(t), t) + \bar{\mu}(t)v_x(Y(t), t) + \frac{1}{2}\bar{\sigma}^2(t)v_{xx}(Y(t), t) \right) dt \\ &\quad + \int_{t_n}^t \bar{\sigma}(t)v_x(Y(t), t) dB(t). \end{aligned}$$

Here $v(Y(t_n), t_n) = 0$ and after taking expectation we get

$$\begin{aligned} \left| \mathbf{E}(v(Y(t), t)) \right| &= \left| \int_{t_n}^t \mathbf{E} \left(v_t(Y(t), t) + \bar{\mu}(t)v_x(Y(t), t) + \frac{1}{2}\bar{\sigma}^2(t)v_{xx}(Y(t), t) \right) dt \right| \\ &\leq h_n \sup_{t \in (t_n, t_{n+1})} \left(|v_t(Y(t), t)| + |\bar{\mu}(t)| |v_x(Y(t), t)| + \frac{1}{2}|\bar{\sigma}^2(t)| |v_{xx}(Y(t), t)| \right), \end{aligned}$$

which proves (46) provided that

$$\begin{aligned} \max_{\mathbf{R} \times [0, T]} |\mu(x, t)| &\leq C, \\ \max_{\mathbf{R} \times [0, T]} |\sigma(x, t)| &\leq C, \\ \max_{\mathbf{R} \times [0, T]} (|v_t(x, t)| + |v_x(x, t)| + |v_{xx}(x, t)|) &\leq C, \end{aligned}$$

where the last one in its turn follows from the bounds

$$\begin{aligned} \max_{\mathbf{R} \times [0, T]} (|\mu_t(x, t)| + |\mu_x(x, t)| + |\mu_{xx}(x, t)|) &\leq C, \\ \max_{\mathbf{R} \times [0, T]} (|u_{xt}(x, t)| + |u_x(x, t)| + |u_{xx}(x, t)| + |u_{xxx}(x, t)|) &\leq C. \end{aligned}$$

Such bounds can be proved provided that μ, σ, g are sufficiently smooth and decay sufficiently fast as $|x| \rightarrow \infty$. This proves (46), and (47) is proved in the same way. \square

This proof was inspired by [5].

If X_0 is a random variable, then we must first show

$$u(X_0, 0) = \mathbf{E}(g(X(T)) \mid X_0),$$

so that (by the law of double expectation)

$$\mathbf{E}(u(X_0, 0)) = \mathbf{E}(\mathbf{E}(g(X(T)) \mid X_0)) = \mathbf{E}(g(X(T)))$$

instead of (45). The proof can then be continued in the same way. We do not present the details.

2.3. Sampling error. In practice we compute several sample paths $\{Y(t, \omega_j)\}_{j=1}^M$, $\omega_j \in \Omega$, and approximate the expected value $\mathbf{E}(g(X(T)))$ by the average $\frac{1}{M} \sum_{j=1}^M g(Y(T, \omega_j))$. The total error is then the sum of the *discretization error* and the *statistical error (sampling error)*:

$$\begin{aligned} & \left| \frac{1}{M} \sum_{j=1}^M g(Y(T, \omega_j)) - \mathbf{E}(g(X(T))) \right| \\ & \leq \left| \mathbf{E}(g(Y(T))) - \mathbf{E}(g(X(T))) \right| + \left| \frac{1}{M} \sum_{j=1}^M (g(Y(T, \omega_j)) - \mathbf{E}(g(Y(T)))) \right|. \end{aligned}$$

We have shown that the first part is $\leq Ch$, and by using the *central limit theorem* it can be shown that the sampling error is $\leq \frac{C}{\sqrt{M}}$. See [5].

2.4. A numerical method of higher order: Milstein's method. We derive the first steps of the *Itô-Taylor expansion*. We consider for simplicity the autonomous equation

$$dX(t) = \mu(X(t)) dt + \sigma(X(t)) dB(t),$$

where $\mu = \mu(x), \sigma = \sigma(x)$ do not depend on t . Then

$$(48) \quad X(t) = X(t_0) + \int_{t_0}^t \mu(X(s)) ds + \int_{t_0}^t \sigma(X(s)) dB(s).$$

Itô's formula is, see Klebaner (4.53),

$$\begin{aligned} (49) \quad f(X(t)) &= f(X(t_0)) + \int_{t_0}^t \left(\mu(X(s)) f'(X(s)) + \frac{1}{2} \sigma^2(X(s)) f''(X(s)) \right) ds \\ &+ \int_{t_0}^t \sigma(X(s)) f'(X(s)) dB(s) \\ &= f(X(t_0)) + \int_{t_0}^t L_0 f(X(s)) ds + \int_{t_0}^t L_1 f(X(s)) dB(s), \end{aligned}$$

where the differential operators L_0 and L_1 are defined by

$$\begin{aligned} L_0 f &= \mu f' + \frac{1}{2} \sigma^2 f'', \\ L_1 f &= \sigma f'. \end{aligned}$$

In particular, with $f(x) = x$ we get $L_0 f = \mu$, $L_1 f = \sigma$ and we retrieve (48). Next we take $f = \mu$ and $f = \sigma$ in (49) and insert the result into (48). We get

$$\begin{aligned} X(t) &= X(t_0) + \int_{t_0}^t \left(\mu(X(t_0)) + \int_{t_0}^s L_0 \mu(X(z)) dz + \int_{t_0}^s L_1 \mu(X(z)) dB(z) \right) ds \\ &\quad + \int_{t_0}^t \left(\sigma(X(t_0)) + \int_{t_0}^s L_0 \sigma(X(z)) dz + \int_{t_0}^s L_1 \sigma(X(z)) dB(z) \right) dB(s) \\ &= X(t_0) + \mu(X(t_0)) \int_{t_0}^t ds + \sigma(X(t_0)) \int_{t_0}^t dB(s) + R_1(t, t_0), \end{aligned}$$

where the remainder is

$$\begin{aligned} R_1(t, t_0) &= \int_{t_0}^t \int_{t_0}^s L_0 \mu(X(z)) dz ds \\ &\quad + \int_{t_0}^t \int_{t_0}^s L_1 \mu(X(z)) dB(z) ds \\ &\quad + \int_{t_0}^t \int_{t_0}^s L_0 \sigma(X(z)) dz dB(s) \\ &\quad + \int_{t_0}^t \int_{t_0}^s L_1 \sigma(X(z)) dB(z) dB(s). \end{aligned}$$

This is the motivation for Euler's method.

Next use Itô's formula with $f = L_1 \sigma = \sigma \sigma'$ and insert into the last term in the remainder to get

$$\begin{aligned} X(t) &= X(t_0) + \mu(X(t_0)) \int_{t_0}^t ds + \sigma(X(t_0)) \int_{t_0}^t dB(s) \\ &\quad + \sigma(X(t_0)) \sigma'(X(t_0)) \int_{t_0}^t \int_{t_0}^s dB(z) dB(s) + R_2(t, t_0), \end{aligned}$$

where the new remainder $R_2(t, t_0)$ consists of five complicated terms.

This is the motivation for Milstein's method. We have here

$$\int_{t_0}^t ds = t - t_0, \quad \int_{t_0}^t dB(s) = B(t) - B(t_0),$$

and the iterated Itô integral, see Klebaner Example 4.12 on p. 107,

$$\int_{t_0}^t \int_{t_0}^s dB(z) dB(s) = \int_{t_0}^t \left(B(s) - B(t_0) \right) dB(s) = \frac{1}{2} \left((B(t) - B(t_0))^2 - (t - t_0) \right).$$

The Milstein method defines $\{Y_n\}_{n=0}^N$ by $Y_0 = X_0$ and

$$Y_{n+1} = Y_n + \mu(Y_n) h_n + \sigma(Y_n) \Delta B_n + \frac{1}{2} \sigma(Y_n) \sigma'(Y_n) ((\Delta B_n)^2 - h_n),$$

where

$$h_n = t_{n+1} - t_n, \quad \Delta B_n = B(t_{n+1}) - B(t_n).$$

It can be shown that it converges with strong order h :

$$\mathbf{E}(|Y_n - X(t_n)|) \leq Ch.$$

This idea can be extended to derive methods of higher order but these are not very useful in practice because the coefficients are many times iterated integrals, for example, $\int_{t_0}^t \cdots \int_{t_0}^{z_2} dB(z_1) \cdots dB(z_n) ds_1 \cdots ds_k$, which are difficult to evaluate.

More material on iterated Itô formulas and Itô-Taylor expansions can be found in [4].

REFERENCES

- [1] L. C. Evans, “An introduction to stochastic differential equations”, lecture notes, Department of Mathematics, University of California, Berkeley. <http://www.math.berkeley.edu/~evans/SDE.course.pdf>
- [2] D. J. Higham “An Algorithmic Introduction to Numerical Simulation of SDE”, SIAM Review 43 (2001), 525–522. [link](#)
- [3] F. C. Klebaner, “Introduction to Stochastic Calculus with Applications”, second edition, Imperial College Press, 2005.
- [4] P. E. Kloeden and E. Platen “Numerical Solution of Stochastic Differential Equations”, Springer 1995.
- [5] K. S. Moon, J. Goodman, A. Szepessy, R. Tempone, and G. Zouraris, “Stochastic and Partial Differential Equations with Adapted Numerics”, Chapter 5, lecture notes, NADA, Royal Institute of Technology, Stockholm. <http://www.math.kth.se/~szepessy/sdepde.pdf>
<http://www.math.kth.se/~szepessy/sdepde.ps>