

NOTES ON TIME DEPENDENT PROBLEMS IN 2D

1. THE MODEL PROBLEM

We first consider the following time dependent model problem,

$$(1) \quad \begin{aligned} \dot{u} - \nabla \cdot (a \nabla u) &= f, & x &= (x_1, x_2) \in \Omega, & 0 < t < T, \\ u(x, t) &= 0, & x &= (x_1, x_2) \in \partial\Omega, & 0 < t < T, \\ u(x, 0) &= u_0(x), & x &= (x_1, x_2) \in \Omega, \end{aligned}$$

where $u = u(x, t) = u(x_1, x_2, t)$ is the unknown function that we wish to compute, with time derivative, $\frac{\partial u}{\partial t}$, denoted by \dot{u} . We assume that $\Omega \subset \mathbb{R}^2$ has a polygonal boundary. The functions $a = a(x, t)$ and $f = f(x, t)$ are *data* to the problem. We also need to specify *boundary data*: in this model problem we have *homogeneous Dirichlet boundary conditions* ($u = 0$) on the entire boundary $\partial\Omega$, for all times, $0 < t < T$, and *initial data*: $u_0(x)$, which specifies the solution, for $x \in \Omega$, at time $t = 0$.

2. THE NUMERICAL METHOD

We shall construct a numerical method by *first discretizing in space* (using finite elements) to obtain a finite dimensional system of linear, ordinary differential equations, which we finally solve numerically using, e.g., the backward Euler method.

2.1. Space Discretization.

2.1.1. *Variational Formulation.* Multiply the differential equation in (1) by a *test function* $v = v(x_1, x_2)$ such that $v = 0$ on $\partial\Omega$ and integrate over Ω :

$$\iint_{\Omega} \dot{u}v \, dx_1 dx_2 - \iint_{\Omega} \nabla \cdot (a \nabla u)v \, dx_1 dx_2 = \iint_{\Omega} fv \, dx_1 dx_2, \quad 0 < t < T.$$

We now integrate by parts (see the notes on *Robin Boundary Conditions in 2D* for details):

$$\iint_{\Omega} \dot{u}v \, dx_1 dx_2 - \int_{\partial\Omega} (n \cdot (a \nabla u))v \, ds + \iint_{\Omega} a \nabla u \cdot \nabla v \, dx_1 dx_2 = \iint_{\Omega} fv \, dx_1 dx_2, \quad 0 < t < T.$$

Since

$$v = 0 \text{ on } \partial\Omega,$$

we obtain

$$\iint_{\Omega} \dot{u}v \, dx_1 dx_2 + \iint_{\Omega} a \nabla u \cdot \nabla v \, dx_1 dx_2 = \iint_{\Omega} fv \, dx_1 dx_2, \quad 0 < t < T.$$

We thus state the following *variational formulation* of (1):

Find $u(x_1, x_2, t)$ such that, for every *fixed* t : $u(x_1, x_2, t) \in V_0$, and

$$(2) \quad \iint_{\Omega} \dot{u}v \, dx_1 dx_2 + \iint_{\Omega} a \nabla u \cdot \nabla v \, dx_1 dx_2 = \iint_{\Omega} fv \, dx_1 dx_2, \quad 0 < t < T, \quad \forall v \in V_0,$$

where V_0 denotes the vector space of functions $v = v(x_1, x_2)$ such that $v = 0$ on $\partial\Omega$, that are sufficiently regular for the integrals in (2) to exist.

2.1.2. Discretization in space. In order to discretize (2) in space, we introduce the vector space V_{h0} of *continuous, piecewise linear* functions, $v(x_1, x_2)$, on a *triangulation*, $\mathcal{T}_h = \{K_i\}_{i=1}^{ntri}$, of Ω , with the corresponding set of *internal nodes*, $\mathcal{N}_{h0} = \{N_i\}_{i=1}^{nintnodes}$, such that $v = 0$ on $\partial\Omega$, and state the following (*space*) *discrete* counterpart of (2):

Find $U(x_1, x_2, t)$ such that, for every *fixed* t : $U(x_1, x_2, t) \in V_{h0}$, and

$$(3) \quad \iint_{\Omega} \dot{U}v \, dx_1 dx_2 + \iint_{\Omega} a \nabla U \cdot \nabla v \, dx_1 dx_2 = \iint_{\Omega} fv \, dx_1 dx_2, \quad 0 < t < T, \quad \forall v \in V_{h0}.$$

2.1.3. Ansatz. We now seek a solution, $U(x_1, x_2, t)$, to (3), expressed (for every *fixed* t) in the basis of *tent functions* $\{\varphi_i\}_{i=1}^{nintnodes} \subset V_{h0}$. (Note that only “tents” with “poles” at the internal nodes belong to the basis, since all functions in V_{h0} are zero on the boundary $\partial\Omega$.) In other words, we make the *Ansatz*

$$(4) \quad U(x_1, x_2, t) = \sum_{j=1}^{nintnodes} \xi_j(t) \varphi_j(x_1, x_2),$$

and seek to determine the (time dependent) coefficient vector

$$\xi(t) = \begin{bmatrix} \xi_1(t) \\ \xi_2(t) \\ \vdots \\ \xi_{nintnodes}(t) \end{bmatrix} = \begin{bmatrix} U(N_1, t) \\ U(N_2, t) \\ \vdots \\ U(N_{nintnodes}, t) \end{bmatrix},$$

of nodal values of $U(x_1, x_2, t)$, in such a way that (3) is satisfied.

Consider *very carefully* the structure of $U(x_1, x_2, t)$ in (4): For every *fixed* time, t , we note that $U(x_1, x_2, t)$, as a function of $x = (x_1, x_2)$, is a continuous, piecewise linear function with weights given by $\xi(t)$.

2.1.4. Construction of space discrete system of ODE. We substitute (4) into (3),

$$(5) \quad \sum_{j=1}^{nintnodes} \dot{\xi}_j(t) \left(\iint_{\Omega} \varphi_j v \, dx_1 dx_2 \right) + \sum_{j=1}^{nintnodes} \xi_j(t) \left(\iint_{\Omega} a \nabla \varphi_j \cdot \nabla v \, dx_1 dx_2 \right) = \iint_{\Omega} fv \, dx_1 dx_2, \quad 0 < t < T, \quad \forall v \in V_{h0}.$$

Since $\{\varphi_i\}_{i=1}^{nintnodes} \subset V_{h0}$ is a *basis* of V_{h0} , (5) is equivalent to

$$A(t) = \begin{bmatrix} a_{1,1}(t) & \cdots & a_{1,\text{nin}}(t) \\ \vdots & \ddots & \vdots \\ a_{\text{nin},1}(t) & \cdots & a_{\text{nin},\text{nin}}(t) \end{bmatrix} \text{ is the (possibly time dependent) } \textit{stiffness matrix}, \text{ and}$$

$$b(t) = \begin{bmatrix} b_1(t) \\ \vdots \\ b_{\text{nin}}(t) \end{bmatrix} \text{ is the (possibly time dependent) } \textit{load vector}.$$

Exercise 1. Show that, for the time dependent reaction-diffusion problem with Robin boundary conditions,

$$\begin{aligned} u - \nabla \cdot (a \nabla u) + cu &= f, & x &= (x_1, x_2) \in \Omega, & 0 < t < T, \\ -n \cdot (a \nabla u) &= \gamma(u - g_D) + g_N, & x &= (x_1, x_2) \in \partial\Omega, & 0 < t < T, \\ u(x, 0) &= u_0(x), & x &= (x_1, x_2) \in \Omega, \end{aligned}$$

the system (7) generalizes to,

$$(8) \quad M \dot{\xi}(t) + (A(t) + M_c(t) + R(t)) \xi(t) = b(t) + rv(t), \quad 0 < t < T,$$

where $M_c(t)$ is the *mass matrix* coming from the reactive term, $c(x_1, x_2, t)u(x_1, x_2, t)$, and $R(t)$, $rv(t)$ are the contributions from the Robin boundary conditions to the system matrix and right-hand side, respectively. (Compare with the notes on *Robin Boundary Conditions in 2D*). Note that (8) is an n_{nodes} -dimensional system of linear, ordinary differential equations, since in this case we also include the nodes on the boundary $\partial\Omega$.

2.2. Time Discretization. In order to discretize (7) in time, we let $0 = t_0 < t_1 < t_2 < \cdots < t_L = T$ be discrete time levels with corresponding time steps $k_n = t_n - t_{n-1}$, $n = 1, \dots, L$. Further, we let ξ^n denote an *approximation* of $\xi(t_n)$, $n = 1, \dots, L$.

There are different possible choices of *initial data*, $\xi^0 = \xi(0)$, to (7): the simplest is to let

$$\xi^0 = \begin{bmatrix} \xi_1(0) \\ \xi_2(0) \\ \vdots \\ \xi_{n_{\text{intnodes}}}(0) \end{bmatrix} = \begin{bmatrix} u_0(N_1) \\ u_0(N_2) \\ \vdots \\ u_0(N_{n_{\text{intnodes}}}) \end{bmatrix},$$

which corresponds to letting $U(x_1, x_2, 0) = \sum_{j=1}^{n_{\text{intnodes}}} \xi_j(0) \varphi_j(x_1, x_2)$ be the *nodal interpolant* of $u_0(x_1, x_2) = u(x_1, x_2, 0)$. (An alternative would be to choose $U(x_1, x_2, 0)$ as the $L_2(\Omega)$ -projection of u_0 , but then we would need to *compute* ξ^0 .)

We now *integrate* (7) (element-wise) over one time interval $[t_{n-1}, t_n]$,

$$\int_{t_{n-1}}^{t_n} M \dot{\xi}(t) dt + \int_{t_{n-1}}^{t_n} A(t) \xi(t) dt = \int_{t_{n-1}}^{t_n} b(t) dt.$$

Since M is a constant matrix, we get,

$$(9) \quad M(\xi(t_n) - \xi(t_{n-1})) + \int_{t_{n-1}}^{t_n} A(t) \xi(t) dt = \int_{t_{n-1}}^{t_n} b(t) dt.$$

Given an approximation, ξ^{n-1} , of $\xi(t_{n-1})$, approximating the integrals in (9) using *right end-point quadrature* gives the *backward Euler method* defining ξ^n by,

$$M(\xi^n - \xi^{n-1}) + A(t_n)\xi^n k_n = b(t_n)k_n,$$

i.e.,

$$M \frac{\xi^n - \xi^{n-1}}{k_n} + A(t_n)\xi^n = b(t_n).$$

The *backward Euler method* for solving (7) thus becomes: Given $\xi^0 = \xi(0)$, for $n = 1, \dots, L$, solve the linear system of equations,

$$(M + k_n A_n)\xi^n = M\xi^{n-1} + k_n b_n,$$

where we have introduced the notation

$$A_n = A(t_n), \quad b_n = b(t_n).$$

Exercise 2. Show that the backward Euler method for solving (8) reads: Given $\xi^0 = \xi(0)$, for $n = 1, \dots, L$, solve the linear system of equations:

$$(M + k_n(A(t_n) + M_c(t_n) + R(t_n)))\xi^n = M\xi^{n-1} + k_n(b(t_n) + rv(t_n)).$$