

# *Adaptive Variational Multiscale Methods*

**Axel Målqvist**

`axel.malqvist@it.uu.se`

**Division of Scientific Computing  
Uppsala University**

# Outline and Papers

---

## Outline

- Model problem, elliptic linear pde
- Variational multiscale method, symmetric version
- Derivation of proposed method, examples
- A posteriori error estimation
- Adaptivity
- Application to oil reservoir problem
- Convection dominated problem
- Future work

## Papers


- M.G. Larson and A. Målqvist, *Adaptive Variational Multiscale Methods Based on A Posteriori Error Estimation: Energy Norm Estimates for Elliptic Problems*, CMAME 2007
- M.G. Larson and A. Målqvist, *A Mixed Adaptive Variational Multiscale Method with Applications in Oil Reservoir Simulation* M3AS 2009 (accepted)

## Model Problem

**Poisson equation:** Find  $u$  such that

$$\begin{aligned} -\nabla \cdot a \nabla u &= f \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

where  $a(x) \geq a_0 > 0$  bounded,  $f \in L^2(\Omega)$ , and  $\Omega \in \mathbf{R}^d$ ,  $d = 2, 3$ .




**Weak form:** Find  $u \in \mathcal{V} = H_0^1(\Omega)$  such that

$$a(u, v) = (a \nabla u, \nabla v) = (f, v) = l(v) \quad \text{for all } v \in \mathcal{V}.$$

# Variational Multiscale Method: Hughes et. al. 95, 98

We split the space  $\mathcal{V}_c \oplus \mathcal{V}_f = \mathcal{V}$



- $\mathcal{V}_c$  is a finite dimensional approximation of  $\mathcal{V}$ . (finite element space)
- $\mathcal{V}_f$  can be chosen in different ways
  - Hierarchical basis
  - $L^2(\Omega)$ -orthogonal to  $\mathcal{V}_c$
  - Wavelet modified hierarchical basis

# Symmetric Variational Multiscale Method

Starting from the model problem: find  $u \in \mathcal{V}$  such that

$$a(u, v) = l(v) \quad \text{for all } v \in \mathcal{V}$$

and setting

$$u = u_c + u_f \quad v = v_c + v_f$$

we get: find  $u_c + u_f \in \mathcal{V}_c \oplus \mathcal{V}_f$  such that

$$a(u_c + u_f, v_c + v_f) = l(v_c + v_f) \quad \text{for all } v_c + v_f \in \mathcal{V}_c \oplus \mathcal{V}_f$$

Note that  $u_f \in \mathcal{V}_f$  satisfies the equation

$$a(u_f, v_f) = l(v_f) - a(u_c, v_f) \quad \text{for all } v_f \in \mathcal{V}_f$$

## Fine Scale Equations

Given the fine scale equation

$$a(u_f, v_f) = l(v_f) - a(u_c, v_f) \quad \text{for all } v_f \in \mathcal{V}_f$$

we let  $u_f = u_{f,l} + u_{f,c} \in \mathcal{V}_f$  with

$$a(u_{f,l}, v_f) = l(v_f) \quad \text{for all } v_f \in \mathcal{V}_f$$

$$a(u_{f,c}, v_f) = -a(u_c, v_f) \quad \text{for all } v_f \in \mathcal{V}_f$$

Let  $\mathcal{T} : \mathcal{V}_c \rightarrow \mathcal{V}_f$  denote the solution operator  $u_{f,c} = \mathcal{T}u_c$ . We get

$$u = u_c + \mathcal{T}u_c + u_{f,l}$$

$$a(u_c + \mathcal{T}u_c + u_{f,l}, v_c + v_f) = l(v_c + v_f)$$

for all  $v_c \in \mathcal{V}_c$  and  $v_f \in \mathcal{V}_f$ .

## Coarse Scale Equations

Since  $u_{f,l}$  is directly determined we get the following problem for  $u_c$ : find  $u_c \in \mathcal{V}_c$  such that

$$a(u_c + \mathcal{T}u_c, v_c + \mathcal{T}v_c) = l(v_c + \mathcal{T}v_c) - a(u_{f,l}, v_c + \mathcal{T}v_c)$$

for all  $v_c \in \mathcal{V}_c$ .

- Here we chose  $v_f = \mathcal{T}v_c$  to get a symmetric formulation
- Note that  $a((I + \mathcal{T})v_c, v_f) = 0$  and  $l(v_f) - a(u_{f,l}, v_f) = 0$  i.e.  $I + \mathcal{T}$  decouples the problem. Any choice of  $v_f \in \mathcal{V}_f$  is ok.
- In standard VMS  $v_f = 0$  in this step and thus when approximating the local effects using numerical or analytical tools the resulting method usually gives non-symmetric matrix.

## Approximation of Fine Scale Solutions

- Let  $\tilde{\mathcal{T}}$  be a computable approximation of  $\mathcal{T}$
- Let  $U_{f,l}$  be a computable approximation of  $u_{f,l}$

We get the method: find  $U_c \in \mathcal{V}_c$  such that

$$a(U_c + \tilde{\mathcal{T}}U_c, v_c + \tilde{\mathcal{T}}v_c) = l(v_c + \tilde{\mathcal{T}}v_c) - a(U_{f,l}, v_c + \tilde{\mathcal{T}}v_c)$$

for all  $v_c \in \mathcal{V}_c$ . On matrix form this leads to,

$$\tilde{K}U_c = \tilde{b}$$

Given  $U_c$ ,  $U_{f,l}$ , and  $\tilde{\mathcal{T}}$ ,  $U_f$  can be computed.

Compare with [Hou et. al. 97] or [Arbogast 04] where basis functions are modified using local computations.



## Construction of $\tilde{\mathcal{T}}$

Recall that  $u_c = \sum_i u_{c,i} N_{c,i}$  with  $\{N_{c,i}\}$  as a basis in  $\mathcal{V}_c$  and let

$$a(\mathcal{T} N_{c,i}, v_f) = -a(N_{c,i}, v_f) \quad \text{for all } v_f \in \mathcal{V}_f$$

By linearity


$$\mathcal{T} u_c = \sum_i u_{c,i} \mathcal{T} N_{c,i}$$

and thus we are led to computing  $\mathcal{T} N_{c,i}$  for each coarse basis function  $N_{c,i}$ .

We define  $\tilde{\mathcal{T}}$  by solving these problems approximately by

- Restricting to a localized patch problem  $\text{supp}(N_{c,i}) \subset \omega_i$
- Discretizing using a fine subgrid on  $\omega_i$

# Refinement and Layers



We let  $H$  be coarse scale mesh size and  $h$  be fine scale mesh size. Further we let  $L$  denote the number of layers of coarse elements in the patch. Typically homogeneous Dirichlet boundary condition are used.

## Construction of $U_{f,l}$

Recall that  $u_{f,l} \in \mathcal{V}_f$  solves

$$a(u_{f,l}, v_f) = l(v_f) \quad \text{for all } v_f \in \mathcal{V}_f$$


Using a partition of unity  $\varphi_i$  we can split the right hand side as follows  $l(v_f) = \sum_i l(\varphi_i v_f)$  to get,

$$u_{f,l} = \sum_i u_{f,l,i}$$
$$a(u_{f,l,i}, v_f) = l(\varphi_i v_f)$$

Again we find an approximation by restricting to patches and discretizing the subgrid.

# Parallel Structure

One local problem for each coarse dof, minimal communication.






## Simple Observation About Decay in $\mathcal{V}_f$

Consider,

$$-\Delta u = \varphi_i \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$


where  $\varphi_i$  has local support in center of  $\Omega$ . The weak form reads:  
find  $u \in \mathcal{W}$  s.t.,  $(\nabla u, \nabla v) = (\varphi_i, v)$  for all  $v \in \mathcal{W}$ .



To the left  $\mathcal{W} = \mathcal{V}_c \oplus \mathcal{V}_f$ , middle  $\mathcal{W} = \mathcal{V}_f$  using hierarchical split,  
and right  $\mathcal{W} = \mathcal{V}_f$  using  $L^2$ -orthogonal split.

# Simple Observation About Decay in $\mathcal{V}_f$

Decay of flux integrated over the boundary.






We see exponential decay with respect distance measured in nof coarse elements. This effect gives rapid convergence as the patch size increases.

# Application to a Mixed Problem

Poisson equation on mixed form:

$$\begin{cases} \frac{1}{a} \boldsymbol{\sigma} - \nabla u = 0 & \text{in } \Omega \\ -\nabla \cdot \boldsymbol{\sigma} = f & \text{in } \Omega \\ \boldsymbol{n} \cdot \boldsymbol{\sigma} = 0 & \text{on } \Gamma \end{cases}$$

where the permeability  $a$  is constant, random, or taken from the SPE data set (upperness in log-scale),



## Splitting Based on RT-elements

We use lowest order RT basis functions together with piecewise constants.

- Let  $\pi_c$  is the RT-interpolant onto  $\mathcal{V}_c$  and  $P_c$  be the  $L^2$ -projection onto  $W_c$
- We define  $\sigma = \pi_c \sigma + (I - \pi_c) \sigma$  and thus  $\sigma_f = (I - \pi_c) \sigma \in \mathcal{V}_f$   $\sigma_c = \pi_c \sigma \in \mathcal{V}_c$ .
- Further we define  $u = P_c u_c + (1 - P_c) u = u_c + u_f \in \mathcal{W}_c \oplus \mathcal{W}_f$ .
- Thus we are using an  $L^2$ -orthogonal splitting in the scalar variable.

Hierarchical split for lagrangian elements leads to nodal exactness in the coarse solution while here we get exactness of average values on coarse elements.



## Some Terms Disappear

Find  $\boldsymbol{\sigma}_c \in \mathcal{V}_c$ ,  $\boldsymbol{\sigma}_f \in \mathcal{V}_f$ ,  $u_c \in \mathcal{W}_c$ , and  $u_f \in \mathcal{W}_f$  such that,

$$\left\{ \begin{array}{l} (\frac{1}{a}(\boldsymbol{\sigma}_c + \boldsymbol{\sigma}_f), \mathbf{v}_c + \mathbf{v}_f) + (u_c + u_f, \nabla \cdot (\mathbf{v}_c + \mathbf{v}_f)) = 0 \\ \quad -(\nabla \cdot (\boldsymbol{\sigma}_c + \boldsymbol{\sigma}_f), w_c + w_f) = (f, w_c + w_f) \\ (\frac{1}{a}\boldsymbol{\sigma}_f, \mathbf{v}_f) + (u_f, \nabla \cdot \mathbf{v}_f) = -(\frac{1}{a}\boldsymbol{\sigma}_c, \mathbf{v}_f) - (u_c, \nabla \cdot \mathbf{v}_f) \\ \quad -(\nabla \cdot \boldsymbol{\sigma}_f, w_f) = (f, w_f) + (\nabla \cdot \boldsymbol{\sigma}_c, w_f) \end{array} \right.$$

for all  $\mathbf{v}_c \in \mathcal{V}_c$ ,  $\mathbf{v}_f \in \mathcal{V}_f$ ,  $w_c \in \mathcal{W}_c$ , and  $w_f \in \mathcal{W}_f$ .

Since for coarse elements  $K$

$$(w_f, \nabla \cdot \mathbf{v}_c) = \sum_K \nabla \cdot \mathbf{v}_c \int_K w_f dx = 0,$$

$$(w_c, \nabla \cdot \mathbf{v}_f) = \sum_K w_c \int_K \nabla \cdot \mathbf{v}_f dx = \sum_K w_c \int_{\partial K} \mathbf{n} \cdot \mathbf{v}_f ds = 0.$$


## Approximate Fine Scales

$$\begin{cases} (\frac{1}{a}\boldsymbol{\sigma}_f, \mathbf{v}_f) + (u_f, \nabla \cdot \mathbf{v}_f) = -(\frac{1}{a}\boldsymbol{\sigma}_c, \mathbf{v}_f) \\ -(\nabla \cdot \boldsymbol{\sigma}_f, w_f) = (f, w_f) \end{cases}$$

- We apply the abstract framework
- Divide the fine scale problem into contributions from the coarse scale part  $\boldsymbol{\sigma}_c$  and right hand side  $f$
- Let  $\boldsymbol{\sigma}_c = \sum_i \sigma_{c,i} \phi_i$  where  $\phi_i$  are the Raviart-Thomas basis functions. Solve the local problem driven by the basis functions (one problem for each basis function)
- Localize by restricting the problem to a patch and using homogeneous Neumann conditions
- Discretize using a suitable subgrid

## Example of Local Solutions $\xi_i = \mathcal{T} \phi_i$


$$\begin{cases} (\frac{1}{a} \xi_i, \mathbf{v}_f) + (\beta_i, \nabla \cdot \mathbf{v}_f) = -(\frac{1}{a} \phi_i, \mathbf{v}_f) \\ -(\nabla \cdot \xi_i, w_f) = 0. \end{cases}$$



We use 3 layer patches and plot absolute value of the flux  $|\xi_i|$ .

## Example of Convergence

- Reference mesh has  $32 \times 32$  elements
- The coarse mesh has  $8 \times 8$  elements.
- We let  $f = 1$  lower left corner and  $f = -1$  in upper right, otherwise  $f = 0$ .



Error compared to reference solution.

## A Posteriori Error Estimate (mixed version, Neumann)

The following energy norm bound holds

$$\|\boldsymbol{\sigma} - \boldsymbol{\Sigma}\|_a^2 = \left\| \frac{1}{\sqrt{a}} (\boldsymbol{\sigma} - \boldsymbol{\Sigma}) \right\|^2 \leq C_a \sum_i (R_{\omega_i}^2 + R_{\partial\omega_i}^2)$$

where

$$R_{\omega_i}^2 = \left\| \frac{1}{a} (\boldsymbol{\Sigma}_c^i \boldsymbol{\phi}_i + \boldsymbol{\Sigma}_{f,i}) - \nabla U_{f,i}^* \right\|_{\omega_i}^2 + \left\| \frac{h}{a} (f\psi_i + \nabla \cdot (\boldsymbol{\Sigma}_c^i \boldsymbol{\phi}_i + \boldsymbol{\Sigma}_{f,i})) \right\|_{\omega_i}^2 \\ + \sum_{K \in \omega_i} \|h^{-1/2} [U_{f,i}^*]\|_{\partial K}^2$$

$$R_{\partial\omega_i}^2 = \|h^{-1/2} U_{f,i}^*\|_{\partial\omega_i \setminus \Gamma}^2$$

$U^*$  is a post processed version (Lovadina and Stenberg 06)

of  $U$ ,  $C_a \sim \|\sqrt{a}\|_{L^\infty(\omega_i)}$ .

## Adaptive Strategy


We have the error bound

$$\|\sigma - \Sigma\|_a^2 \leq C_a \sum_i (R_{\omega_i}^2 + R_{\partial\omega_i}^2)$$


1. Let  $h = H/2$  and  $L = 1$  for all  $i$ .
2. Compute the solution  $\{U, \Sigma\}$ .
3. Calculate residuals for each coarse RT basis functions.
4. Mark large entries.
5. For marked entries  $R_{\omega_i}^2$  let  $h := h/2$ .
6. For marked entries  $R_{\partial\omega_i}^2$  let  $L := L + 1$ .
7. Return to 1 or stop if estimators are small enough.

# Adaptivity with Permeability from SPE Data Set

Layer 1 and 50 in the SPE comparative sol. proj. (log scale).




Plot of the sol. (pressure),  $q = 1$  upper right  $q = -1$  lower left.



## Refinements and layers SPE50

We use  $55 \times 15$  coarse elements and a reference mesh with  $440 \times 120$  elements.




We start the adaptive algorithm with **one refinement** and **one layer** in all local problems. After three iterations in the algorithm marking 30%.



# Convergence of Adaptive Algorithm

We compare error in energy norm with reference solution.




- Critical areas are found
- A majority of the patches uses one layer and one refinement.

# Convection Dominated Problem

$$-\epsilon \Delta u + \nabla \cdot (bu) = f \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

where  $\epsilon = 0.01$ ,  $f = 1$  lower left corner,  $b = [b_x, b_x]$ ,  $b_x$  oscillates between 0.01 and 1, 48 periods. Let  $H = 1/24$ ,  $h = H/4$  and study relative error of mean compared to reference solution.



Hierarchical split with continuous linear finite elements.

## Summary and Future Work

---

The AVMS provides:

- Systematic technique for construction of a computable approximation of the fine scale part of the solution using decoupled localized subgrid problems.
- A posteriori error estimation framework (also for goal functionals)
- Adaptive algorithms for automatic tuning of critical discretization parameters

Future work:

- Scale up numerics, parallel code, 3D.
- A priori error analysis, capture decay.
- Multiscale approach to the coupled transport-pressure equation. (Time dependent problems Nordbotten 09)
- Tests on more realistic data, compare with other methods.