

### MAN 240 (2003) : Inlämningsuppgift 3

In the following, to translate from old Biggs to new Biggs, replace chapters 8,9,10,11,12 by chapters 15,16,17,18 and 19.

1. Get your hands on an up-to-date map of Europe. To have something to compare with, see for example

<http://www.wunderground.com/global/Region/EU/Temperature.html>

Make a graph  $G$  whose nodes are the countries of Europe, with an edge between every pair of countries which share a border.

(a) Order the countries alphabetically according to their Swedish names, and then color  $G$  using the greedy algorithm. How many colors are used ?

(b) Do the same in English. How many colors used this time ?

(c) What is  $\chi(G)$  ? Explain.

(Note : Don't forget to include teeny weeny countries like Lichtenstein, Andorra, San Marino, Vatican City and Monaco !!).

2. Exercises 8.8.5 and 8.8.6.

3 (a) How many pairwise non-isomorphic trees on 8 vertices are there ? Draw them all.

(b) According to Cayley's theorem, there are 16 labelled trees on 4 vertices. Draw them all !

4. Let  $G$  be any graph. Prove that either  $G$  or its' complement must be connected.

5. Let  $G$  be a directed graph without a directed cycle. Prove that  $G$  is a network, i.e.: has a source and a sink.

6. Let  $X$  be any compact 2-D surface in  $\mathbf{R}^3$ . Explain why  $\chi(X) = 2 - 2g$ , where  $g$  is the number of 'holes' in  $X$ .

7. Let  $P$  denote the Petersen graph.

(a) Give an explicit isomorphism between the usual pentagonal representa-

tion of  $P$  and the hexagonal one in exercise 8.8.3.

- (b) Indicate cycles of lengths 5,6,8 and 9 in  $P$ .
- (c) Show that it is not possible to edge-color  $P$  with 3 colors.
- (d) Solve exercise 10.7.1.
- (e) Hence, or otherwise, deduce that  $P$  has no Hamilton cycle.

**8.** Refer to the network in Fig. 12.3.

- (a) Take away all the arrows and find a minimal weight spanning tree in the resulting undirected graph.
- (b) Find a shortest path from  $s$  to  $t$ .
- (c) Find a maximal flow from  $s$  to  $t$ .

**9 (a)** Give an example to show that the claim of exercise 8.8.22 is false.

(b) On the other hand show that, for any graph  $G$  on  $n$  vertices,

$$\chi(G) + \chi(\overline{G}) \leq n + 1.$$

**10.** Investigate which  $n$ -tuples  $(d_1, \dots, d_n)$  of positive integers can be the degrees of the vertices in a simple (resp. multi-) graph on  $n$  vertices. In this connection, solve exercise 8.8.17.

**11.** The following is a famous theorem :

*Turan's theorem* : Let  $n, p$  be positive integers. Let  $t, r$  be such that

$$n = t(p - 1) + r \quad \text{and} \quad 1 \leq r \leq p - 1.$$

Put

$$M(n, p) := \frac{p-2}{2(p-1)}n^2 - \frac{r(p-1-r)}{2(p-1)}.$$

Then any simple graph on  $n$  vertices with more than  $M(n, p)$  edges contains a copy of  $K_p$ .

(a) Give an example of a graph with  $n$  vertices and  $M(n, p)$  edges which contains no  $K_p$ .

(b) Show that the argument we used to prove Mantel's theorem (the special case  $p = 3$ ) gives, instead of exactly  $M(n, p)$ , the slightly weaker upper

bound

$$|E(G)| \leq \frac{p-2}{2(p-1)}n^2.$$

**\*(c)** If you're feeling lucky, prove the theorem !!

**12.** Given 12 coins, one of which is flawed (so that it is either slightly lighter or heavier than the others), describe a strategy for finding the flawed coin which requires at most three weighings.

(Note : I don't remember if the strategy also reveals whether the flawed coin is light or heavy).

**13.** Let  $G$  be a graph and  $n$  a positive integer. Define  $\chi_G(n)$  to be the number of ways to vertex-color  $G$  using at most  $n$  colors. More precisely,  $\chi_G(n)$  is the number of (not necessarily surjective !) functions

$$f : V(G) \rightarrow \{1, \dots, n\}$$

such that  $f(x) \neq f(y)$  whenever vertices  $x$  and  $y$  are adjacent in  $G$ .

Your task is to find formulas for  $\chi_G(n)$  for the following graphs :

- (a)  $G$  is a tree on  $k$  vertices.
- (b)  $G$  is the complete graph on  $k$  vertices.
- (c)  $G$  is the cycle of length  $k$ .
- (d)  $G$  is the graph to the left in Fig. 8.5.

(Note : In parts (a), (b) and (c), your formula will be a function of both  $k$  and  $n$ ).