

A Finite Element Crash Course

M. Asadzadeh F. Bengzon

October 1, 2004

1 Continuous Piecewise Linears

Consider a partition,

$$\mathcal{T}_h : 0 = x_0 < x_1 < x_2 < \dots < x_N = 1, \quad (1.1)$$

of the interval $0 \leq x \leq 1$ into N subintervals of equal length h , see Figure 1 below.




Figure 1: Illustration of a partition \mathcal{T}_h .

Let us use this partition to define a function space, V_h , the space of all continuous piecewise linear functions, vanishing at the left end-point $x = 0$ of the interval.

A set of basis functions $\{\varphi_i\}_1^N$ for V_h is defined by

$$\varphi_i(x_j) = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}$$

Because of their shape, these functions are commonly called *hat functions*.




Figure 2: Basis functions, or *hat function*, $\varphi_i(x)$ for V_h .

Note that there is no hat function at $x = 0$ and only a half hat at $x = 1$.




Figure 3: A piecewise linear sine function.

2 Weak and Variational Formulations

Consider the two-point boundary value problem

$$\begin{aligned} -u''(x) &= f(x), & 0 < x < 1, \\ u(0) &= 0, & u'(1) = 0. \end{aligned} \tag{2.1}$$

A *variational formulation* of (2.16) is obtained by multiplying the *residual* $R(u) = -u''(x) - f(x)$ by any smooth function $v(x)$ satisfying $v(0) = 0$, and integrating over the interval $0 \leq x \leq 1$. Because $R(u) = 0$, we have trivially

$$\int_0^1 R(u)v(x) dx = \int_0^1 -u''(x)v(x) - f(x)v(x) dx = 0, \tag{2.2}$$

that is,

$$-\int_0^1 u''(x)v(x) dx = \int_0^1 f(x)v(x) dx, \quad (2.3)$$

for all functions $v(x)$. Consequently, the variational formulation says that the residual $R(u)$ should be orthogonal to $v(x)$. Usually, $v(x)$ is referred to as a *test function*.

Further, using the regularity (i.e., smoothness) of $v(x)$ we can integrate by parts, to get

$$\begin{aligned} -\int_0^1 u''(x)v(x) dx &= -\left[u'(x)v(x)\right]_0^1 + \int_0^1 u'(x)v'(x) dx \\ &= -u'(1)v(1) + u'(0)v(0) + \int_0^1 u'(x)v'(x) dx. \end{aligned} \quad (2.4)$$

Since $u'(1) = 0$, and $v(0) = 0$, we have the *weak form*¹ of (2.16)

$$\int_0^1 u'(x)v'(x) dx = \int_0^1 f(x)v(x) dx, \quad (2.5)$$

for all $v(x)$ such that $v(0) = 0$.

2.1 A Finite Element Method

If $U(x)$ is a solution approximation to (2.16) the residual $R(U)$ can no longer be zero for *every* test function $v(x)$. Instead, we replace the condition that the residual should be identically zero by the condition that the residual should be orthogonal to all functions within a suitable so-called *test space* V_h . We choose the space of all continuous piecewise linear functions, which at $x = 0$, as our test space.

A *Finite Element Method* for (2.16) reads: find $U(x) \in V_h$, such that

$$\int_0^1 U'(x)v'(x) dx = \int_0^1 f(x)v(x) dx, \quad (2.6)$$

for all $v(x) \in V_h$.

¹Note that the weak form only involves one derivative of u whereas (2.16) involves two.

Note that since we also choose $U(x) \in V_h$ we have the *ansatz*,

$$U(x) = \xi_1\varphi_1(x) + \xi_2\varphi_2(x) + \dots + \xi_N\varphi_N(x). \quad (2.7)$$

Here, ξ_i , $i = 1, 2, \dots, N$ are unknown constants that must be determined. We do this by evaluating (2.6) for N different choices of test functions $v(x)$. As a result, we get a system of equations from which the constants ξ_i can be computed.

Hence, substituting (2.7) into (2.6) and choosing the basis functions as test functions², i.e., $v(x) = \varphi_j(x)$ for $j = 1, 2, \dots, N$, we get a system of equations,

$$\left\{ \begin{array}{l} \int_0^1 (\xi_1\varphi_1' + \xi_2\varphi_2' + \dots + \xi_N\varphi_N')\varphi_1' dx = \int_0^1 f\varphi_1 dx \\ \int_0^1 (\xi_1\varphi_1' + \xi_2\varphi_2' + \dots + \xi_N\varphi_N')\varphi_2' dx = \int_0^1 f\varphi_2 dx \\ \vdots \\ \int_0^1 (\xi_1\varphi_1' + \xi_2\varphi_2' + \dots + \xi_N\varphi_N')\varphi_N' dx = \int_0^1 f\varphi_N dx \end{array} \right. . \quad (2.8)$$

Using matrix notation, we write this as

$$\mathbf{A}\boldsymbol{\xi} = \mathbf{b}, \quad (2.9)$$

where the entries of the matrix \mathbf{A} and the vector \mathbf{b} are given by

$$a_{ij} = \int_0^1 \varphi_i'(x) \varphi_j'(x) dx, \quad b_j = \int_0^1 f(x)\varphi_j(x) dx, \quad i, j = 1, 2, \dots, N. \quad (2.10)$$

Assembly Process. To compute the integrals of (2.10) we note that

$$\varphi_i(x) = \frac{1}{h} \begin{cases} x - x_{i-1}, & \text{if } x_{i-1} \leq x \leq x_i, \\ x_{i+1} - x, & \text{if } x_i \leq x \leq x_{i+1}, \\ 0, & \text{otherwise.} \end{cases} \quad (2.11)$$

²It is easy to verify that testing (2.6) against all $v \in V_h$ reduces to testing against the basis functions φ_j . This is a consequence of the fact that these bases are linearly independent. In other words, if (2.6) is satisfied for each φ_j separately, then it must also be satisfied for an arbitrary combination of them.




Figure 4: Overlapping and non-overlapping hat functions.

For $|i - j| > 1$, we have $a_{ij} = 0$, since the bases $\varphi_i(x)$ and $\varphi_j(x)$ lack any overlapping support, that is, one of them is always zero. However, if $i = j$, then

$$a_{ii} = \int_0^1 \varphi_i(x)\varphi_i(x) dx = \int_{x_{i-1}}^{x_i} \frac{1}{h^2} dx + \int_{x_i}^{x_{i+1}} \frac{(-1)(-1)}{h^2} dx = \frac{2}{h}. \quad (2.12)$$

where we have used that $x_i - x_{i-1} = x_{i+1} - x_i = h$. Further, if $j = i + 1$, then

$$a_{i,i+1} = \int_0^1 \varphi_i(x)\varphi_{i+1}(x) dx = \int_{x_i}^{x_{i+1}} \frac{(-1)}{h} \frac{1}{h} dx = -\frac{1}{h}. \quad (2.13)$$

Changing i to $i - 1$, we also get $a_{i-1,i} = -1/h$. Hence, by symmetry, we have

$$\mathbf{A} = \frac{1}{h} \begin{bmatrix} 2 & -1 & & & & \\ -1 & 2 & -1 & & & \\ & -1 & 2 & -1 & & \\ & & \ddots & \ddots & \ddots & \\ & & & -1 & 2 & -1 \\ & & & & -1 & 1 \end{bmatrix}. \quad (2.14)$$

Note the last entry which is just $1/h$, because the basis function $\varphi_N(x)$ is half.

Generally, the entries b_i of the vector \mathbf{b} must be evaluated by quadrature, since they involve the function $f(x)$. However, if $f(x) = 1$, then we can

compute b_i by noting that it is the area under $\varphi_i(x)$. We thus get

$$b_i = \int_0^1 f(x)\varphi_i(x) dx = \int_{x_{i-1}}^{x_{i+1}} \varphi_i(x) dx = h. \quad (2.15)$$

Solving (2.9) using $f(x) = 1$ and $N = 30$, gives $U(x)$ as shown below.




Figure 5: Finite element solution $U(x)$.

2.2 Three Equivalent Problems

Let us turn to consider the more general boundary value problem

$$-(au')' = f, \quad 0 < x < 1, \quad u(0) = u(1) = 0. \quad (2.16)$$

where $a(x) > 0$ is a positive, piecewise continuous, and bounded function for $0 \leq x \leq 1$.

Let us also introduce the function space V , defined by

$$V = \{v : \|v\| + \|v'\| < \infty, \quad v(0) = v(1) = 0\}. \quad (2.17)$$

As before, the variational formulation of the problem (2.16) is obtained by multiplying the equation by a function $v \in V$ and integrating over the interval $0 \leq x \leq 1$, i.e.,

$$-\int_0^1 (au')'v dx = \int_0^1 fv dx. \quad (2.18)$$

By partial integration, we get

$$-[au'v]_0^1 + \int_0^1 au'v' dx = \int_0^1 fv dx. \quad (2.19)$$

but, since $v(0) = v(1) = 0$, we have

$$\int_0^1 au'v' dx = \int_0^1 fv dx. \quad (2.20)$$

So, that the *variational formulation* reads: find $u \in V$, such that

$$\int_0^1 au'v' dx = \int_0^1 fv dx, \quad (2.21)$$

for all $v \in V$.

Theorem 1. *A solution u to (2.16) also satisfies the variational formulation (2.21).*

Proof. Consider (2.21) written as $-\int_0^1 (au')'v dx = \int_0^1 fv dx$, which can also be written as

$$\int_0^1 (-(au')' - f)v dx = 0, \quad (2.22)$$

for all $v \in V$. We claim that this also implies $-(au')' = f$. To see this, suppose that our claim is not true. Then, there exists a point ξ within the interval $0 \leq x \leq 1$, such that $-(au')' - f > 0$ (or, < 0) at this point. In this case, let us take a positive (or, negative) function v and insert it into (2.22). We get

$$\int_0^1 (-(au')' - f)v dx > 0, \quad (2.23)$$

which contradicts (2.22). Thus, our claim is true and the proof is complete.

Equivalent Minimization Problem. Our boundary value problem (2.16) can also be formulated as a minimization problem, namely, find $u \in V$, such that

$$F(u) = \min_{w \in V} F(w). \quad (2.24)$$

Here, $F(w)$ is the total energy of $w(x)$ given by

$$F(w) = \frac{1}{2} \int_0^1 a(w')^2 dx - \int_0^1 fw dx. \quad (2.25)$$

Theorem 2. *The minimization problem (2.24) above is equivalent with the variational formulation (2.21).*

Proof. (\Leftarrow) For $w \in V$, let $v = w - u$, then $v \in V$ and

$$\begin{aligned} F(w) &= F(u + v) = \frac{1}{2} \int_0^1 a(u' + v')^2 dx - \int_0^1 f(u + v) dx \\ &= \underbrace{\frac{1}{2} \int_0^1 a u'^2 dx}_A + \underbrace{\int_0^1 a u' v' dx}_B + \frac{1}{2} \int_0^1 a v'^2 dx - \underbrace{\int_0^1 f u dx}_C - \underbrace{\int_0^1 f v dx}_D, \end{aligned} \quad (2.26)$$

but $B + D = 0$ by virtue of (2.21). Further, by the definition of F we have $A + C = F(u)$. Thus,

$$F(w) = F(u) + \frac{1}{2} \int_0^1 a(v')^2 dx, \quad (2.27)$$

and since $a(x) > 0$ we must have $F(w) > F(u)$.

(\Rightarrow) Let us now assume that $F(u) \leq F(w)$ and set $g(\varepsilon) = F(u + \varepsilon v)$. We know then that g has a minimum at $\varepsilon = 0$, that is to say, $g'(0) = 0$. However,

$$\begin{aligned} g(\varepsilon) &= F(u + \varepsilon v) = \frac{1}{2} \int_0^1 a(u' + \varepsilon v')^2 dx - \int_0^1 f(u + \varepsilon v) dx \\ &= \frac{1}{2} \int_0^1 a(u')^2 + 2a\varepsilon u' v' + a\varepsilon^2 (v')^2 dx - \int_0^1 f u dx - \varepsilon \int_0^1 f v dx, \end{aligned} \quad (2.28)$$

so $g'(\varepsilon)$ is given by

$$g'(\varepsilon) = \frac{1}{2} \int_0^1 2a\varepsilon (v')^2 + 2a u' v' dx - \int_0^1 f v dx. \quad (2.29)$$

Evaluating this derivative at $\varepsilon = 0$ implies that

$$\int_0^1 a u' v' dx - \int_0^1 f v dx = 0, \quad (2.30)$$

which is precisely the variational formulation (2.21).

2.3 A Basic Error Estimate

Numerical techniques, such as the Finite Element Method, are worthless unless they are supplemented by some form of error analysis. *A priori* error analyses are performed theoretically and before any computations has been done. Here, we show a basic error estimate.

Theorem 3. *Let $u \in V$ be a solution to*

$$-u'' = f, \quad 0 < x < 1, \quad u(0) = 0, \quad u'(1) = 0. \quad (2.31)$$

Also, let $U \in V_h$ be a solution approximation satisfying

$$\int_0^1 U'v' dx = \int_0^1 fv dx, \quad (2.32)$$

for all $v \in V_h$. Here, $V_h \subset V = \{v : \|v\| + \|v'\| < \infty, v(0) = v(1) = 0\}$ is the usual space of piecewise linears vanishing at $x = 0$ and $x = 1$. It holds that the error $e = u - U$ obeys

$$\|(u - U)'\| \leq \|(u - v)'\|, \quad (2.33)$$

for all $v \in V_h$.

Proof: Pick a function $v \in V_h$. We then have

$$\begin{aligned} \|(u - U)'\|^2 &= \int_0^1 (u' - U')^2 dx \\ &= \int_0^1 (u' - U')(u' - v' + v' - U') dx \\ &= \int_0^1 (u' - U')(u' - v') dx + \int_0^1 (u' - U')(v' - U') dx. \end{aligned} \quad (2.34)$$

Further, it also holds that

$$\int_0^1 u'v' dx = \int_0^1 fv dx, \quad (2.35)$$

for all $v \in V$, and that

$$\int_0^1 U'v' dx = \int_0^1 fv dx, \quad (2.36)$$

for all $v \in V_h \subset V$. Subtracting these two relations we have

$$\int_0^1 (u' - U')v' dx = 0, \quad (2.37)$$

for all $v \in V_h$. As a consequence, the last integral of the estimate vanish, i.e.,

$$\|(u - U)'\|^2 = \int_0^1 (u' - U')(u' - v') dx \leq \|(u - U)'\| \|(u - v)'\|, \quad (2.38)$$

where we used the Cauchy-Schwarz inequality. Hence,

$$\|(u - U)'\| \leq \|(u - v)'\|, \quad (2.39)$$

and the proof is complete.