

# Lecture 8: The Simplex method

## An algebraic derivation of the pricing step

$$\begin{aligned}
 z^* &= \infimum \mathbf{c}^T \mathbf{x} &= \infimum \mathbf{c}_B^T \mathbf{x}_B + \mathbf{c}_N^T \mathbf{x}_N \\
 &\text{subject to } \mathbf{A}\mathbf{x} = \mathbf{b}, &\text{subject to } \mathbf{B}\mathbf{x}_B + \mathbf{N}\mathbf{x}_N = \mathbf{b}, \\
 &\mathbf{x} \geq \mathbf{0}^n &\mathbf{x}_B \geq \mathbf{0}^m; \mathbf{x}_N \geq \mathbf{0}^{n-m} \\
 &= \mathbf{c}_B^T \mathbf{B}^{-1} \mathbf{b} + \infimum [\mathbf{c}_N^T - \mathbf{c}_B^T \mathbf{B}^{-1} \mathbf{N}] \mathbf{x}_N \\
 &\text{subject to } \mathbf{B}^{-1} \mathbf{b} - \mathbf{B}^{-1} \mathbf{N} \mathbf{x}_N \geq \mathbf{0}^m, \\
 &\mathbf{x}_N \geq \mathbf{0}^{n-m}
 \end{aligned}$$

- $\mathbf{x}_N = \mathbf{0}^{n-m}$  feasible. Let  $\tilde{\mathbf{c}}_N := \mathbf{c}_N^T - \mathbf{c}_B^T \mathbf{B}^{-1} \mathbf{N}$
- If *reduced cost*  $\tilde{\mathbf{c}}_N \geq \mathbf{0}^{n-m}$  then  $\mathbf{x}_N = \mathbf{0}^{n-m}$  is optimal

- If  $\tilde{\mathbf{c}}_N \not\geq \mathbf{0}^{n-m}$  then  $\exists j \in N$  with  $\tilde{c}_j < 0$ . Then the current point  $\mathbf{x}_N = \mathbf{0}^{n-m}$  may be non-optimal
- Generate a feasible descent direction
- Choose one that leads to a neighboring extreme point
- Swap one variable in  $B$  for one in  $N$
- Increase one variable in  $N$  from zero
- Choose  $j^*$  to be among  $\arg \min_{j \in N} \tilde{c}_j$
- We have then decided on the search direction

## The basis change

- What is this direction?
- In  $\mathbf{x}_N$ -space:  $\mathbf{p}_N = \mathbf{e}_{j^*}$  (unit vector)
- In  $\mathbf{x}_B$ -space:  $\mathbf{x}_B = \mathbf{B}^{-1}\mathbf{b} - \mathbf{B}^{-1}\mathbf{N}\mathbf{x}_N \implies \mathbf{p}_B = -\mathbf{B}^{-1}\mathbf{N}\mathbf{p}_N = -\mathbf{B}^{-1}\mathbf{N}_{j^*}$
- So, search direction in  $\mathbb{R}^n$ :

$$\mathbf{p} = \begin{pmatrix} \mathbf{p}_B \\ \mathbf{p}_N \end{pmatrix} = \begin{pmatrix} -\mathbf{B}^{-1}\mathbf{N}_{j^*} \\ \mathbf{e}_{j^*} \end{pmatrix}$$

- Descent? Yes, because  $\mathbf{c}^T\mathbf{p} = \tilde{c}_{j^*} < 0!$

- Feasible? Must check that  $\mathbf{A}\mathbf{p} = \mathbf{0}^m$  and that  $p_i \geq 0$  if  $x_i = 0$ ,  $i \in B$ . The first true by construction:

- (a)  $\mathbf{A}\mathbf{p} = \mathbf{B}\mathbf{p}_B + \mathbf{N}\mathbf{p}_N = -\mathbf{B}\mathbf{B}^{-1}\mathbf{N}_{j^*} + \mathbf{N}\mathbf{e}_{j^*} = \mathbf{0}^m$

- (b) Suppose that  $\mathbf{x}_B > \mathbf{0}^m$ . Then, at least a small step in  $\mathbf{p}$  keeps  $\mathbf{x}_B \geq \mathbf{0}^m$ .

But if there is an  $i^*$  with  $(\mathbf{x}_B)_{i^*} = 0$  and  $(\mathbf{p}_B)_{i^*} < 0$  then it is not a feasible direction

- Must then perform a basis change without moving! A *degenerate basis change*: swap  $x_{j^*}$  for  $x_{i^*}$  in the basis
- Otherwise (and normally), we utilize the unit direction

- Line search? Linear objective; move the maximum step!
- Maximum step: If  $\mathbf{p}_B \geq \mathbf{0}^m$  there is no finite maximum step! We have found an extreme direction  $\mathbf{p}$  along which  $\mathbf{c}^T \mathbf{x}$  tends to  $-\infty$ ! *Unbounded solution*
- Otherwise: Some basic variables will reach zero eventually. Choose as the outgoing variable a variable  $i \in B$  with minimum in

$$x_{j^*} := \underset{i \in B}{\text{minimum}} \left\{ \frac{(\mathbf{B}^{-1} \mathbf{b})_i}{(\mathbf{B}^{-1} \mathbf{N}_{j^*})_i} \mid (\mathbf{B}^{-1} \mathbf{N}_{j^*})_i > 0 \right\}$$

- Done. In the basis, replace  $i^*$  by  $j^*$ ; goto pricing step
- If  $x_{j^*} = 0$  then the above corresponds to a “degenerate basis change”

## Computational notes—how do we do all of this?

- Given basis matrix  $B$ , solve

$$Bx_B = b$$

- Gives us BFS:  $x_B = B^{-1}b$

- Pricing step: (a) Solve

$$B^T y = c_B \implies y^T = c_B^T B^{-1}$$

- (b) Calculate  $\tilde{c}_N^T = c_N^T - y^T N$ , the reduced cost vector
- Choose the incoming variable,  $x_{j^*}$

- Outgoing variable: Solve

$$\mathbf{B}\mathbf{p}_B = -\mathbf{N}_{j^*}$$

- Quotient rule for  $(\mathbf{B}^{-1}\mathbf{b})_i/(-\mathbf{p}_B)_i$  gives outgoing variable,  $x_{i^*}$ , and value of the new basic variable,  $x_{j^*}$
- Note: Three similar linear systems in  $\mathbf{B}$ ! LU factorization + three triangular substitutions
- Factorizations can be updated after basis change rather than done from scratch
- LP solvers like Cplex and XPRESS-MP have excellent numerical solvers for linear systems
- Linear systems the bulk of the work in solving an LP

## Convergence

- *If all of the basic feasible solutions are non-degenerate, then the Simplex algorithm terminates after a finite number of iterations*
- *Proof: (Rough argument) Non-degeneracy implies that the step length is  $> 0$ ; hence, we cannot return to an old BFS once we have left it. There are finitely many BFSs*

- Degeneracy: Can actually lead to cycling—the same sequence of BFSs is returned to indefinitely!
- Remedy: Change the incoming/outgoing criteria!  
Bland's rule: Sort variables according to some index ordering. Take the first possible index in the list.  
Incoming variable first in the list with the right sign of the reduced cost; outgoing variable the first in the list among the minima in the quotient rule

## Initial BFS: Phase I of the Simplex method

- If a starting BFS cannot be found, do the following:
- Suppose  $\mathbf{b} \geq \mathbf{0}^m$ . Introduce *artificial variables*  $a_i$  in every row (or rows without a unit column)
- Solve the following Phase-I problem:

$$\begin{aligned}
 &\text{minimize} && w = (\mathbf{1}^m)^\top \mathbf{a} \\
 &\text{subject to} && \mathbf{Ax} + \mathbf{I}^m \mathbf{a} = \mathbf{b}, \\
 & && \mathbf{x} \geq \mathbf{0}^n, \\
 & && \mathbf{a} \geq \mathbf{0}^m
 \end{aligned}$$

- Possible cases: (a)  $w^* = 0$ , meaning that  $\mathbf{a}^* = \mathbf{0}^m$  must hold. There is then a BFS in the *original* problem

- Start Phase-II, to solve the original problem, starting from this BFS
- (b)  $w^* > 0$ . The optimal basis then has some  $a_i^* > 0$ ; due to the objective function construction, there exists no BFS in the original problem. The problem is then infeasible!
- What to do then? Modelling errors? Can be detected from the optimal solution. In fact, some LP problems are pure feasibility problems