Milton & Arnold, Chapter 1, OH no 1

Definition 1.2.1: Sample space and sample point

A sample space for an experiment is a set S with the property that each physical outcome of the experiment corresponds to exactly one element of S, An element of S is called a sample point.

# Definition 1.2.2: Event

Any subset A of a sample space S is called an event. The emptyset  $\emptyset$  is called the impossible event; the subset S is called the certain event.

#### Träddiagram

Se exempel 1.2.1 och 1.2.2, sida 6

#### $\underline{\text{Venndiagram}}$

Se t.ex figur 2.1, 2.2 och 2.3, sida 27–29

Definition 1.2.3: Mutually exclusive events

Two events  $A_1$  and  $A_2$  are mutually exclusive if and only if  $A_1\cap A_2=\emptyset$ . Events  $A_1,A_2,A_3,\ldots$  are mutually exclusive if and only if  $A_i\cap A_j=\emptyset$  for  $i\neq j$ .

Se exempel 1.2.4, sida 9

Del A

Milton & Arnold, Chapter 1, OH no 2

Relative Frequency Approximation

 $P[A] \approx \frac{f}{n} = \frac{\text{number of times event } A \text{ occurred}}{\text{number of times experiment was run}}$ 

Se exempel 1.1.2, sida 4

Classical Formula

$$P[A] = \frac{n(A)}{n(S)} = \frac{\text{number of ways } A \text{ can occur}}{\text{number of ways the experiment can proceed}}$$

Se exempel 1.1.3, sida 4-5

Del A

Milton & Arnold, Chapter 1, OH no 3

## Definition 1.3.1: Permutation

A permutation is an arrangement of objects in a definite order.

#### Multiplication principle

Consider an experiment taking place in k stages. Let  $n_i$  denote the number of ways in which stage i can occur for  $i=1,2,\ldots,k$ . Alltogether the experiment can occur in

$$\prod^k n_i = n_1 \cdot n_2 \cdot \cdot \cdot n_k$$

ways.

# Theorem 1.3.1: Counting permutations

The number of permutations of n distinct objects used r at a time denoted by  ${}_nP_r,$  is

$${}_{n}P_{r} = \frac{n!}{(n-r)!}$$

Del A

Milton & Arnold, Chapter 1, OH no 4

#### Definition 1.3.2: Combination

A combination is an selection of objects without regard to order.

## Theorem 1.3.2: Counting combinations

The number of combinations of n distinct objects selected r at a time, denoted by  ${}_{n}C_{r}$  or  $\binom{n}{r}$ , is given by

$$_{n}C_{r} = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

Notera sambandet  ${}_{n}P_{r} = {}_{n}C_{r} \cdot r!$ 

# $\underline{A\,nkomstkontroll}$

Se exempel 1.3.6, sida 15