

## Definition 1.2.1: Sample space and sample point

A sample space for an experiment is a set  $S$  with the property that each physical outcome of the experiment corresponds to exactly one element of  $S$ . An element of  $S$  is called a sample point.

## Definition 1.2.2: Event

Any subset  $A$  of a sample space  $S$  is called an event. The emptyset  $\emptyset$  is called the *impossible* event; the subset  $S$  is called the *certain* event.

Träddiagram

Se exempel 1.2.1, sida 6

Venn diagram

Se figur 2.1, sida 27

## Definition 1.2.3: Mutually exclusive events

Two events  $A_1$  and  $A_2$  are mutually exclusive if and only if  $A_1 \cap A_2 = \emptyset$ . Events  $A_1, A_2, A_3, \dots$  are mutually exclusive if and only if  $A_i \cap A_j = \emptyset$  for  $i \neq j$ .

## Relative Frequency Approximation

$$P[A] \approx \frac{f}{n} = \frac{\text{number of times event } A \text{ occurred}}{\text{number of times experiment was run}}$$

Se exempel 1.1.2, sida 4

## Classical Formula

$$P[A] = \frac{n(A)}{n(S)} = \frac{\text{number of ways } A \text{ can occur}}{\text{number of ways the experiment can proceed}}$$

Se exempel 1.1.3, sida 4-5

## Definition 1.3.1: Permutation

A permutation is an arrangement of objects in a definite order.

## Multiplication principle

Consider an experiment taking place in  $k$  stages. Let  $n_i$  denote the number of ways in which stage  $i$  can occur for  $i = 1, 2, \dots, k$ . Altogether the experiment can occur in

$$\prod_{i=1}^k n_i = n_1 \cdot n_2 \cdots n_k$$

ways.

## Theorem 1.3.1: Counting permutations

The number of permutations of  $n$  distinct objects used  $r$  at a time, denoted by  ${}_nP_r$ , is

$${}_nP_r = \frac{n!}{(n-r)!}$$

## Definition 1.3.2: Combination

A combination is an selection of objects without regard to order.

## Theorem 1.3.2: Counting combinations

The number of combinations of  $n$  distinct objects selected  $r$  at a time, denoted by  ${}_nC_r$  or  $\binom{n}{r}$ , is given by

$${}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

Notera sambandet  ${}_nP_r = {}_nC_r \cdot r!$

Ankomstkontroll

Se exempel 1.3.6, sida 15