

## NOTES ON TIME DEPENDENT PROBLEMS

## 1. THE MODEL PROBLEM

We consider the following time dependent model problem,

$$(1) \quad \begin{aligned} \dot{u} - (au')' &= f, & x_{\min} < x < x_{\max}, & 0 < t < T, \\ u(x_{\min}, t) &= 0, & & 0 < t < T, \\ u(x_{\max}, t) &= 0, & & 0 < t < T, \\ u(x, 0) &= u_0(x), & x_{\min} < x < x_{\max}, & \end{aligned}$$

where  $u = u(x, t)$  is the unknown function that we wish to compute, with time derivative,  $\frac{\partial u}{\partial t}$ , denoted by  $\dot{u}$ , and  $x$ -derivative,  $\frac{\partial u}{\partial x}$ , denoted by  $u'$ . The functions  $a = a(x, t)$  and  $f = f(x, t)$  are *data* to the problem. We also need to specify *boundary data*: in (1) we have *homogeneous Dirichlet boundary conditions* at both end-points,  $x = x_{\min}, x_{\max}$ , for all times,  $0 < t < T$ , and *initial data*:  $u_0(x)$ , which specifies the solution, for  $x_{\min} < x < x_{\max}$ , at time  $t = 0$ .

## 2. THE NUMERICAL METHOD

We construct a numerical method by *first discretizing in space* (using finite elements) to obtain a finite dimensional system of linear, ordinary differential equations. We then *discretize in time* and solve the system of ODE numerically (using the backward Euler method).

## 2.1. Space Discretization.

2.1.1. *Variational Formulation.* Multiply the differential equation in (1) by a *test function*  $v(x) \in H_0^1([x_{\min}, x_{\max}]) := \left\{ v(x) : \int_{x_{\min}}^{x_{\max}} v'(x)^2 dx < \infty, v(x_{\min}) = v(x_{\max}) = 0 \right\}$ , and integrate over  $[x_{\min}, x_{\max}]$ :

$$\int_{x_{\min}}^{x_{\max}} \dot{u}v \, dx - \int_{x_{\min}}^{x_{\max}} (au')'v \, dx = \int_{x_{\min}}^{x_{\max}} fv \, dx, \quad 0 < t < T.$$

We now integrate by parts:

$$\int_{x_{\min}}^{x_{\max}} \dot{u}v \, dx - [(au')v]_{x=x_{\min}}^{x=x_{\max}} + \int_{x_{\min}}^{x_{\max}} au'v' \, dx = \int_{x_{\min}}^{x_{\max}} fv \, dx, \quad 0 < t < T.$$

Since

$$v(x_{\min}) = v(x_{\max}) = 0,$$

we obtain

$$\int_{x_{\min}}^{x_{\max}} \dot{u}v \, dx + \int_{x_{\min}}^{x_{\max}} au'v' \, dx = \int_{x_{\min}}^{x_{\max}} fv \, dx, \quad 0 < t < T.$$

We now state the following *variational formulation* of (1):

Find  $u(x, t)$  such that, for every fixed  $t$ :  $u(x, t) \in H_0^1([x_{\min}, x_{\max}])$ , and

*Date*: September 16, 2002.

$$(2) \quad \int_{x_{\min}}^{x_{\max}} \dot{u}v \, dx + \int_{x_{\min}}^{x_{\max}} au'v' \, dx = \int_{x_{\min}}^{x_{\max}} fv \, dx, \quad 0 < t < T, \quad \forall v \in H_0^1([x_{\min}, x_{\max}]).$$

2.1.2. *Discretization in Space.* In order to discretize (2) in space, we introduce the vector space,  $\mathring{V}_h$ , of *continuous, piecewise linear* functions,  $v(x)$ , on a partition,  $x_{\min} = x_0 < x_1 < \dots < x_N < x_{N+1} = x_{\max}$ , of  $[x_{\min}, x_{\max}]$ , such that  $v(x_{\min}) = v(x_{\max}) = 0$ , and state the following (*space discrete*) counterpart of (2):

Find  $U(x, t)$  such that, for every fixed  $t$ :  $U(x, t) \in \mathring{V}_h$ , and

$$(3) \quad \int_{x_0}^{x_{N+1}} \dot{U}v \, dx + \int_{x_0}^{x_{N+1}} aU'v' \, dx = \int_{x_0}^{x_{N+1}} fv \, dx, \quad 0 < t < T, \quad \forall v \in \mathring{V}_h.$$

2.1.3. *Ansatz.* We now seek a solution,  $U(x, t)$ , to (3), expressed (for every fixed  $t$ ) in the basis of *hat functions*  $\{\varphi_i\}_{i=1}^N \subset \mathring{V}_h$ . (Note that  $\varphi_0$  and  $\varphi_{N+1}$  do *not* belong to the basis, since all functions in  $\mathring{V}_h$  are zero at the end-points.) In other words, we make the *Ansatz*

$$(4) \quad U(x, t) = \sum_{j=1}^N \xi_j(t) \varphi_j(x),$$

and seek to determine the (time dependent) coefficient vector

$$\xi(t) = \begin{bmatrix} \xi_1(t) \\ \xi_2(t) \\ \vdots \\ \xi_N(t) \end{bmatrix} = \begin{bmatrix} U(x_1, t) \\ U(x_2, t) \\ \vdots \\ U(x_N, t) \end{bmatrix},$$

of nodal values of  $U(x, t)$ , in such a way that (3) is satisfied.

Consider *very carefully* the structure of  $U(x, t)$  in (4): For every fixed time,  $t$ , we note that  $U(x, t)$ , as a function of  $x$ , is a continuous, piecewise linear function with weights given by  $\xi(t)$ .

2.1.4. *Construction of Space Discrete System of ODE.* We substitute (4) into (3):

$$(5) \quad \sum_{j=1}^N \dot{\xi}_j(t) \left( \int_{x_0}^{x_{N+1}} \varphi_j v \, dx \right) + \sum_{j=1}^N \xi_j(t) \left( \int_{x_0}^{x_{N+1}} a \varphi_j' v' \, dx \right) = \int_{x_0}^{x_{N+1}} fv \, dx, \\ 0 < t < T, \quad \forall v \in \mathring{V}_h.$$

Since  $\{\varphi_i\}_{i=1}^N \subset \mathring{V}_h$  is a *basis* for  $\mathring{V}_h$ , (5) is equivalent to

$$(6) \quad \sum_{j=1}^N \dot{\xi}_j(t) \left( \int_{x_0}^{x_{N+1}} \varphi_j \varphi_i \, dx \right) + \sum_{j=1}^N \xi_j(t) \left( \int_{x_0}^{x_{N+1}} a \varphi_j' \varphi_i' \, dx \right) = \int_{x_0}^{x_{N+1}} f \varphi_i \, dx, \\ 0 < t < T, \quad i = 1, \dots, N,$$



$$b(t) = \begin{bmatrix} b_1(t) \\ \vdots \\ b_N(t) \end{bmatrix} \text{ is the (possibly time dependent) } \textit{load vector}.$$

**2.2. Time Discretization.** To discretize (7) in time, let  $0 = t_0 < t_1 < t_2 < \dots < t_L = T$  be discrete time levels with corresponding time steps  $k_n = t_n - t_{n-1}$ ,  $n = 1, \dots, L$ . Further, let  $\xi^n$  denote the *approximation* of  $\xi(t_n)$ ,  $n = 1, \dots, L$ .

There are different possible choices of *initial data*,  $\xi^0 = \xi(0)$ , to (7): the simplest is

$$\xi^0 = \begin{bmatrix} \xi_1(0) \\ \xi_2(0) \\ \vdots \\ \xi_N(0) \end{bmatrix} = \begin{bmatrix} u_0(x_1) \\ u_0(x_2) \\ \vdots \\ u_0(x_N) \end{bmatrix},$$

which corresponds to letting  $U(x, 0) = \sum_{j=1}^N \xi_j(0) \varphi_j(x)$  be the *nodal interpolant* of  $u_0(x) = u(x, 0)$ . (An alternative would be to choose  $U(x, 0)$  as the  $L_2([x_{\min}, x_{\max}])$ -projection of  $u_0$ , but then we would need to *compute*  $\xi^0$ .)

We now *integrate* (7) (element-wise) over one time interval  $[t_{n-1}, t_n]$ :

$$\int_{t_{n-1}}^{t_n} M \dot{\xi}(t) dt + \int_{t_{n-1}}^{t_n} A(t) \xi(t) dt = \int_{t_{n-1}}^{t_n} b(t) dt.$$

Since  $M$  is a constant matrix, we get:

$$(8) \quad M(\xi(t_n) - \xi(t_{n-1})) + \int_{t_{n-1}}^{t_n} A(t) \xi(t) dt = \int_{t_{n-1}}^{t_n} b(t) dt.$$

Given an approximation,  $\xi^{n-1}$ , of  $\xi(t_{n-1})$ , approximating the integrals in (8) using *right end-point quadrature* gives the *backward Euler method* defining  $\xi^n$  by

$$M(\xi^n - \xi^{n-1}) + A(t_n) \xi^n k_n = b(t_n) k_n,$$

i.e.,

$$(9) \quad M \frac{\xi^n - \xi^{n-1}}{k_n} + A(t_n) \xi^n = b(t_n).$$

For solving (7) using the *backward Euler method* we can now state the following *algorithm*:

Given  $\xi^0 = \xi(0)$ . For  $n = 1, \dots, L$ : Solve the linear system of equations

$$(10) \quad (M + k_n A_n) \xi^n = M \xi^{n-1} + k_n b_n.$$

In (10) we have introduced the notation

$$A_n = A(t_n), \quad b_n = b(t_n).$$

*Remark.* Observe the similarity between (7) and (9): We may alternatively view the backward Euler method as approximating the derivative by a difference quotient, and evaluating the other terms at the right end-point of the time interval.