

# Numerical homogenization of multiscale problems

Axel Målqvist<sup>1</sup>

New trends in asymptotic methods for multiscale PDEs




Karlstad University

2019-10-23

---



<sup>1</sup>Chalmers University of Technology and University of Gothenburg

# Multiscale materials



# Multiscale problems

We consider applications such as



- ▷ composite materials      ▷ flow in a porous medium

that require numerical solution of partial differential equations with rough data (module of elasticity, conductivity, or permeability).

Major challenge: Features on multiple non-separated scales.

# Outline

- ① **Elliptic problem: Homogenization and FEM**
- ② Introduction to LOD
- ③ High contrast data
- ④ Applications
- ⑤ Conclusions



# Elliptic model problem

The Poisson equation

$$-\nabla \cdot \mathbf{A} \nabla u = f \quad \text{in } \Omega \qquad u = 0 \quad \text{on } \partial\Omega$$

with data  $0 < \alpha \leq A \leq \beta < \infty$  and  $f \in L^2(\Omega)$ .

---



# Homogenization of an elliptic model problem

## The Poisson equation

$$-\nabla \cdot A \nabla u = f \quad \text{in } \Omega \quad u = 0 \quad \text{on } \partial\Omega$$

with data  $0 < \alpha \leq A \leq \beta < \infty$  and  $f \in L^2(\Omega)$ .

---

**Homogenization theory** Let  $A = A(x/\epsilon)$  be  $\epsilon$ -periodic, and consider

$$-\nabla \cdot (A(x/\epsilon) \nabla u_\epsilon(x, x/\epsilon)) = f(x).$$

It can be shown that as  $\epsilon \rightarrow 0$ ,  $u_\epsilon \rightarrow v$  solves

$$-\nabla \cdot (A^* \nabla v(x)) = f(x),$$

- In 1D,  $A^* = \frac{1}{\langle 1/A \rangle}$ , i.e. the harmonic average.
- Otherwise  $A^*$  is computed by solving a cell problem.

# Homogenization of an elliptic model problem

## The Poisson equation

$$-\nabla \cdot \mathbf{A} \nabla u = f \quad \text{in } \Omega \quad u = 0 \quad \text{on } \partial\Omega$$

with data  $0 < \alpha \leq A \leq \beta < \infty$  and  $f \in L^2(\Omega)$ .

---

### Homogenization theory

- Considers  $\epsilon \rightarrow 0$ , i.e. large scale separation.
- Periodic data leads to a reference cell problem.
- Single scale rather than a continuum of scales in data.
- Numerical approaches inspired by homogenization theory (e.g. HMM and MsFEM).

Multiscale finite element method, (Hou & Wu), 1996.

Heterogeneous multiscale method, (Engquist & E), 2003.

# Finite element method

The Poisson equation (weak form):  $u \in V = H_0^1(\Omega)$  such that

$$a(u, v) := \int_{\Omega} (\textcolor{brown}{A} \nabla u) \cdot \nabla v \, dx = \int_{\Omega} f \cdot v \, dx \text{ for all } v \in V$$

with data  $0 < \alpha \leq A \leq \beta < \infty$  and  $f \in L^2(\Omega)$ .

# Finite element method

The Poisson equation (FE approximation):  $u_h \in V_h \subset V$  such that

$$a(u_h, v) := \int_{\Omega} (\textcolor{brown}{A} \nabla u_h) \cdot \nabla v \, dx = \int_{\Omega} f v \, dx \text{ for all } v \in V_h$$

with data  $0 < \alpha \leq A \leq \beta < \infty$  and  $f \in L^2(\Omega)$ .

---

Numerical error (piecewise linear continuous FE approximation)

- For solution  $u \in H^2(\Omega)$  we have

$$\|u - u_h\| := \|A^{1/2} \nabla(u - u_h)\|_{L^2(\Omega)} \leq C\beta^{1/2} h \|D^2 u\|_{L^2(\Omega)} \sim C(\alpha, \beta, \textcolor{brown}{A}') h.$$

- The mesh size  $h$  has to resolve the variations in  $A$ , e.g.  $h < \epsilon$  if  $A$  is  $\epsilon$ -periodic.

# Finite element method




The Poisson equation (FE approximation):  $u_h \in V_h \subset V$  such that

$$a(u_h, v) := \int_{\Omega} (\mathbf{A} \nabla u_h) \cdot \nabla v \, dx = \int_{\Omega} f v \, dx \text{ for all } v \in V_h$$

with data  $0 < \alpha \leq A \leq \beta < \infty$  and  $f \in L^2(\Omega)$ .

---

**Example** (periodic coefficient):  $A(x) = (2 - \cos(2\pi x/\epsilon))^{-1}$ ,  $\epsilon = 0.02$ ,  $f = 1$



# Multiscale approach

## Objectives:

- Find a subspace of  $V_H^{\text{ms}} \subset V_h$  for which the Galerkin approximation fulfills

$$\|u_h - u_H^{\text{ms}}\| \leq C(\alpha, \beta) H \approx C(\alpha, \beta, \mathbf{A}') h,$$

but with  $\dim(V_H^{\text{ms}}) \ll \dim(V_h)$ .

- Show that a basis for  $V_H^{\text{ms}}$  can be constructed by local parallel computations.
- Demonstrate efficiency for applications where  $V_H^{\text{ms}}$  is reused (eigenvalue, time dependent, semi-linear, systems).


Variational multiscale method, (Hughes), 1995

# Outline

- ① Elliptic problem: Homogenization and FEM
- ② **Introduction to LOD**
- ③ High contrast data
- ④ Applications
- ⑤ Conclusions

# Multiscale decomposition

- (coarse) FE mesh  $\mathcal{T}$  with parameter  $H > h$
- P1-FE space  $V_H := \{v \in V \mid \forall T \in \mathcal{T}, v|_T \in P_1(T)\}$
- $\mathfrak{I}_{\mathcal{T}} : V \rightarrow V_H$  some interpolation operator




## Decomposition

$$V = V_H \oplus V^f \quad \text{with } V^f := \text{kernel } \mathfrak{I}_{\mathcal{T}} = \{v \in V \mid \mathfrak{I}_{\mathcal{T}} v = 0\}$$

---

Example:



rough coefficient


# Multiscale decomposition

- (coarse) FE mesh  $\mathcal{T}$  with parameter  $H > h$
- P1-FE space  $V_H := \{v \in V \mid \forall T \in \mathcal{T}, v|_T \in P_1(T)\}$
- $\mathfrak{I}_{\mathcal{T}} : V \rightarrow V_H$  some interpolation operator

## Decomposition

$$V = V_H \oplus V^f \quad \text{with } V^f := \text{kernel } \mathfrak{I}_{\mathcal{T}} = \{v \in V \mid \mathfrak{I}_{\mathcal{T}} v = 0\}$$

Example:



# Orthogonalization

- For each  $v \in V$  define finescale projection  $Qv \in V^f$  by


$$a(Qv, w) = a(v, w) \quad \text{for all } w \in V^f$$

## $a$ -Orthogonal Decomposition

$$V = V_H^{\text{ms}} \oplus V^f \quad \text{with } V_H^{\text{ms}} := (V_H - QV_H)$$

---

Example:



# Ideal multiscale representation

Given the space  $V_H^{\text{ms}}$  we construct a Galerkin approximation:

## Ideal method

Find  $u_H^{\text{ms}} \in V_H^{\text{ms}}$  such that

$$a(u_H^{\text{ms}}, v) = (f, v), \quad \forall v \in V_H^{\text{ms}}.$$

We have that  $u - u_H^{\text{ms}} = u_f \in V^f$  since  $u_H^{\text{ms}}$  is the  $a$ -orthogonal projection of  $u$  onto  $V_H^{\text{ms}}$ . Therefore

$$\|u_f\|^2 = a(u, u_f) = (f, u_f) = (f, u_f - \mathfrak{I}_{\mathcal{T}} u_f) \leq \frac{C_{\mathfrak{I}_{\mathcal{T}}}}{\alpha^{1/2}} \|Hf\|_{L^2(\Omega)} \|u_f\|.$$

For  $V_H^{\text{ms}}$  to be useful we need a discrete local basis.


# Modified nodal basis

- $\mathcal{N}$  denotes set of interior vertices of  $\mathcal{T}$
- $\phi_x \in V_H$  denotes classical nodal basis function ( $x \in \mathcal{N}$ )
- $Q\phi_x \in V^f$  denotes the finescale correction of  $\phi_x$  ( $x \in \mathcal{N}$ )

Ideal multiscale FE space

$$V_H^{\text{ms}} = \text{span} \{ \phi_x - Q\phi_x \mid x \in \mathcal{N} \}$$

Example




# Modified nodal basis

- $\mathcal{N}$  denotes set of interior vertices of  $\mathcal{T}$
- $\phi_x \in V_H$  denotes classical nodal basis function ( $x \in \mathcal{N}$ )
- $Q\phi_x \in V^f$  denotes the finescale correction of  $\phi_x$  ( $x \in \mathcal{N}$ )

Ideal multiscale FE space



$$V_H^{\text{ms}} = \text{span} \{ \phi_x - Q\phi_x \mid x \in \mathcal{N} \}$$

Example



# Localization

- Define nodal patches of  $\ell$ -th order  $\omega_{T,\ell}$  about  $T \in \mathcal{T}$

 $\omega_{T,1}$  $\omega_{T,2}$ 

- Correctors  $Q_\ell^T \phi_x \in V^f(\omega_{T,\ell}) := \{v \in V^f \mid v|_{\Omega \setminus \omega_{T,\ell}} = 0\}$  solve


$$a(Q_\ell^T \phi_x, w) = \int_T A \nabla \phi_x \cdot \nabla w \, dx \quad \text{for all } w \in V^f(\omega_{T,\ell})$$

Localized multiscale FE spaces

$$V_{H,\ell}^{\text{ms}} = \text{span}\{\phi_x - \sum_{T \in \mathcal{T}} Q_\ell^T \phi_x \mid x \in \mathcal{N}\}$$

# Fine scale discretization

- Finescale mesh



- Reference FE space

$$V_h := \{v \in V \mid \forall T \in \mathcal{T}(\Omega), v|_T \in P_1(T)\}$$

- Reference FE solution  $u_h \in V_h$  solves

$$a(u_h, v) = (f, v) \quad \text{for all } v \in V_h$$

- Fully discrete correctors  $Q_{\ell,h}^T \phi_x \in V_h^f(\omega_{T,\ell}) := V^f(\omega_{T,\ell}) \cap V_h$ :

$$a(Q_{\ell,h}^T \phi_x, w) = (A \nabla \phi_x, \nabla w)_T \quad \text{for all } w \in V_h^f(\omega_{T,\ell})$$

# Localized Orthogonal Decomposition (LOD)

Fully discrete multiscale FE spaces

$$V_{H,\ell}^{\text{ms},h} = \text{span}\{\phi_x - \sum_{T \in \mathcal{T}} Q_{\ell,h}^T \phi_x \mid x \in \mathcal{N}\}$$

Fully discrete multiscale approximation  $u_{H,\ell}^{\text{ms},h} \in V_{H,\ell}^{\text{ms},h}$

$$a(u_{H,\ell}^{\text{ms},h}, v) = (f, v) \quad \text{for all } v \in V_{H,\ell}^{\text{ms},h}$$

Remarks:

- $\dim V_{H,\ell}^{\text{ms},h} = |\mathcal{N}| = \dim V_H$
- The basis functions have local support, with overlap depending on  $\ell$ , and are independent.

# A priori error analysis

## Lemma (Truncation error)

$$\|Q_h v_H - Q_{\ell,h} v_H\| \leq C_1 \gamma^\ell \|Q_h v_H\|, \quad \forall v_h \in V_H$$

$C_1 < \infty$  and  $\gamma < 1$  depends on  $\beta/\alpha$  but not  $A'$ .

By choosing  $\ell = C_2 \log(H^{-1})$  with appropriate  $C_2$  we guarantee that the truncation leads to a higher order perturbation:



## Theorem (A priori error bound)

$$\|u_h - u_{H,\ell}^{\text{ms},h}\| \leq C(\alpha, \beta) H,$$


with  $C$  independent of  $A'$ .

M. & Peterseim, Localization of elliptic multiscale problems, Math. Comp., 2014.

# Numerical experiment: Poisson's equation





$$H = 2^{-1}, 2^{-2}, \dots, 2^{-7}$$
$$h = 2^{-9}, \ell = \log(1/H)$$




$\|u_h - u_{H,\ell}^{\text{ms},h}\|$  vs. #dof

# Numerical experiment: Poisson's equation



$$H = 2^{-1}, 2^{-2}, \dots, 2^{-7}$$


$$h = 2^{-9}, \ell = \log(1/H)$$



$\|u_h - u_{H,\ell}^{\text{ms},h}\|$  vs. #dof





# 3D implementation in python

- Trilinear shape functions on cubes.
- Petrov-Galerkin formulation reduces communication, Elfversson et.al. Numer. Math. 2016.
- Storage of all basis function is not needed. The full solution can be recomputed (at a lower cost) once  $\mathfrak{I}_{\mathcal{T}} u_{H,\ell}^{\text{ms},h}$  is computed.



Corrector function  $Q^T \phi_x$ , implementation by Fredrik Hellman.




# Numerical experiment: Poisson's equation 3D




$$H = 2^{-2}, 2^{-3}, 2^{-4}, 2^{-5}$$
$$h = 2^{-6}, \ell = \log(1/H)$$

$\|u_h - u_{H,\ell}^{ms,h}\|$  vs. #dof

# Numerical experiment: Poisson's equation 3D



$$H = 2^{-2}, 2^{-3}, 2^{-4}, 2^{-5}$$
$$h = 2^{-6}, \ell = \log(1/H)$$



$\|u_h - u_{H,\ell}^{ms,h}\|$  vs. #dof

# Outline


- ① Elliptic problem: Homogenization and FEM
- ② Introduction to LOD
- ③ **High contrast data**
- ④ Applications
- ⑤ Conclusions

# High contrast data

Poisson equation:

$$-\nabla \cdot A \nabla u = f \quad \text{in } \Omega \quad u = 0 \quad \text{on } \partial\Omega.$$


$A = 1$  in  $\Omega_1$  (black),  $A = \alpha$  in  $\Omega_\alpha$ ,  $\alpha \ll 1$ , and  $f = \chi_{[1/4,3/4]^2}$ .






- High contrast data with channels leads to non-local behaviour.
- The decay rate of the basis functions determines the accuracy of LOD.
- The choice of interpolant  $\mathfrak{I}_T v = \sum_{x \in N} \bar{v}_{\omega_x} \phi_x$  affects the decay.

# Numerical example: High contrast

High contrast data Three examples:  $H = 2^{-4}$ ,  $h = 2^{-10}$ ,



We let  $\alpha = 10^{-1}, \dots, 10^{-6}$  and plot  $\|u_h - u_{H,k}^{\text{ms},h}\|$  vs.  $k$ , with  $\mathfrak{I}_{\mathcal{T}}^{SZ}$ ,



# Scott-Zhang type interpolation

## Nodal variables:

Let  $x \in \mathcal{N}$  be nodes of  $\mathcal{T}$  and  $\sigma_x \subset \Omega$  associated domains. We define a  $L^2(\sigma_x)$ -dual basis  $\psi_x \in V_H$  fulfilling,

$$\int_{\sigma_x} \psi_x \phi_y = \delta_{xy}.$$

Let the nodal variable  $N_x(v) = \int_{\sigma_x} \psi_x v$  and,


$$\mathfrak{I}_{\mathcal{T}}^\sigma v = \sum_{x \in \mathcal{N}} N_x(v) \phi_x.$$

- $\sigma_x$  does not need to be full elements  $T$  or vertex patches  $U_1(x)$ .
- The stability of  $|N_x(v)| \leq \|\psi\|_{L^2(\sigma_x)} \|v\|_{L^2(\sigma_x)}$  depends on the size and shape of  $\sigma_x$  and its distance to  $x$ .

# Geometry dependent interpolation

## Selection of $\sigma_x$ :

By letting  $\sigma_x \subset \Omega_1$  (frequently enough) we guarantee decay,  
i.e. nodes in high conductivity channels are needed.



Let  $U_1(x)$  be the vertex patch at node  $x$ .

- Type I node: for  $x \in \Omega_1$  let  $\sigma_x \subset U_1(x) \cap \Omega_1$ , connected.
- Type II node: for  $x \in \Omega_\alpha$  let  $\sigma_x = U_\delta(x)$ ,  $0 < \delta \leq 1$ .

# Weighted Poincaré inequality and decay

The following weighted Poincaré inequality holds:

$$\|A^{1/2}v_f\|_{L^2(\mathcal{T})} \leq CH \|A^{1/2}\nabla v_f\|_{L^2(U_1(\mathcal{T}))}, \quad \forall v_f \in V^f = \ker(\mathfrak{J}_{\mathcal{T}}^{\sigma}).$$

This is used to prove contrast independent decay.

## Theorem

With  $\delta < 1/2$  we have,

$$\|A^{1/2}\nabla Q^T v_H\|_{\Omega \setminus U_k(\mathcal{T})} \leq C\gamma^k \|A^{1/2}\nabla Q^T v_H\|_{L^2(\Omega)},$$

where  $C$  and  $\gamma$  are independent of  $\beta/\alpha$ .


Hellman & M. Contrast independent localization of multiscale problems,  
MMS 2017

# Numerical example: High contrast

High contrast data Three examples:  $H = 2^{-4}$ ,  $h = 2^{-10}$ ,



We let  $\alpha = 10^{-1}, \dots, 10^{-6}$  and plot  $\|u_h - u_{H,k}^{\text{ms},h}\|$  vs.  $k$  with  $\mathfrak{I}_{\mathcal{T}}^{\text{SZ}}$ ,




# Numerical example: High contrast

High contrast data Three examples:  $H = 2^{-4}$ ,  $h = 2^{-10}$ ,




We let  $\alpha = 10^{-1}, \dots, 10^{-6}$  and plot  $\|u_h - u_{H,k}^{\text{ms},h}\|$  vs.  $k$  with  $\mathfrak{I}_T^\sigma$ ,



# Interface model

Thin structures so far need to be resolved.



As  $\epsilon \rightarrow 0$  we have convergence (with rate) to an interface problem.

$$-\nabla \cdot A \nabla u = f, \quad \text{in } \Omega$$

$$u = 0, \quad \text{on } \partial\Omega$$

$$[u] = 0, \quad \text{on } \Gamma$$

$$-\nabla_\Gamma \cdot A_\Gamma \nabla_\Gamma u = f_\Gamma - [n \cdot A \nabla u], \quad \text{on } \Gamma.$$

# Weak form and LOD

On weak form we have: find  $u \in V = H_0^1(\Omega) \cap H^1(\Gamma)$  such that

$$a(u, v) := \int_{\Omega} A \nabla u \cdot \nabla v \, dx + \int_{\Gamma} A_{\Gamma} \nabla_{\Gamma} u \cdot \nabla_{\Gamma} v \, ds = \int_{\Omega} fv \, dx + \int_{\Gamma} f_{\Gamma} v \, ds,$$

for all  $v \in V = H_0^1(\Omega) \cap H^1(\Gamma)$ . We note that  $a(\cdot, \cdot)$  is a scalar product on  $V$ .

## Localized Orthogonal Decomposition

Given an interpolant  $\mathfrak{I}_{\mathcal{T}} : V \rightarrow V_H$  we can formulate the LOD method: find  $u_H^{\text{ms}} \in V_H^{\text{ms}}$  such that


$$a(u_H^{\text{ms}}, v) = \int_{\Omega} fv \, dx + \int_{\Gamma} f_{\Gamma} v \, ds, \quad \text{for all } v \in V_H^{\text{ms}}.$$

# Edge based Scott-Zhang interpolation

Integration on edges/faces. We have the interpolation bound:

$$\|v - \mathfrak{I}_T v\|_{L^2(\Omega)} + \|v - \mathfrak{I}_T v\|_{L^2(\Gamma)} \leq CH \left( \|v\|_{H^1(\Omega)}^2 + \|v\|_{H^1(\Gamma)}^2 \right)^{1/2}.$$

Decay follows if the nodal variables are restricted to the interfaces.






Element based vs. edge based in log scale with interface is at  $x = 0.5$ .

Hellman, M., & Wang, Numer. upscaling in fractured domains, arXiv, 2019

# Numerical example

Let  $A_\Gamma = 2$  and  $A$  be random between 0.1 and 0.9 with forcing in the bulk and on the interfaces.

Coarser mesh has 237 dof and finest mesh has 219345 dof,  
 $k = \log(H^{-1})$ .



Error in energy norm vs number of uniform refinements.

# Outline


- ① Elliptic problem: Homogenization and FEM
- ② Introduction to LOD
- ③ High contrast data
- ④ **Applications**
- ⑤ Conclusions

# Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem

$$-\nabla \cdot \mathbf{A} \nabla u = \lambda u \quad \text{in } \Omega \quad u = 0 \quad \text{on } \partial\Omega$$

with data  $0 < \alpha \leq A \leq \beta < \infty$




# Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem (variational form): find  $u \in V := H_0^1(\Omega)$  and  $\lambda \in \mathbb{R}$  such that

$$a(u, v) := \int_{\Omega} (\mathbf{A} \nabla u) \cdot \nabla v \, dx = \lambda \int_{\Omega} u \cdot v \, dx \text{ for all } v \in V$$

with data  $0 < \alpha \leq A \leq \beta < \infty$




# Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem (FE approximation):  
 $u_h \in V_h \subset V$  and  $\lambda_h \in \mathbb{R}$  such that

$$a(u_h, v) := \int_{\Omega} (\mathbf{A} \nabla u_h) \cdot \nabla v \, dx = \lambda_h \int_{\Omega} u_h \cdot v \, dx \text{ for all } v \in V_h$$

with data  $0 < \alpha \leq A \leq \beta < \infty$



# Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem (FE approximation):  
 $u_h \in V_h \subset V$  and  $\lambda_h \in \mathbb{R}$  such that

$$a(u_h, v) := \int_{\Omega} (\mathbf{A} \nabla u_h) \cdot \nabla v \, dx = \lambda_h \int_{\Omega} u_h \cdot v \, dx \text{ for all } v \in V_h$$

with data  $0 < \alpha \leq A \leq \beta < \infty$

---

Numerical error (piecewise linear continuous FE approximation)

- For an eigenpair  $(u^{(k)}, \lambda^{(k)})$  with  $u^{(k)} \in H^2(\Omega)$  it holds

$$\lambda^{(k)} \leq \lambda_h^{(k)} \leq \lambda^{(k)} + C(\alpha, \beta, \mathbf{A}', k) h^2,$$

$$\|u^{(k)} - u_h^{(k)}\| := \|\mathbf{A}^{1/2} \nabla (u^{(k)} - u_h^{(k)})\|_{L^2(\Omega)} \leq C(\alpha, \beta, \mathbf{A}', k) h.$$

- The mesh size  $h$  has to resolve the variations in  $A$ , e.g.  $h < \epsilon$  if  $A$  is periodic.

# LOD approximation

**LOD:** Find  $u_{H,\ell}^{\text{ms},h} \in V_{H,\ell}^{\text{ms},h}$ ,  $\lambda_{H,\ell}^{\text{ms},h} \in \mathbb{R}$

$$a(u_{H,\ell}^{\text{ms},h}, v) = \lambda_{H,\ell}^{\text{ms},h} (u_{H,\ell}^{\text{ms},h}, v) \quad \text{for all } v \in V_{H,\ell}^{\text{ms},h}$$

## Theorem


$$\lambda_h^{(k)} \leq \lambda_{H,\ell}^{\text{ms},h,(k)} \leq \lambda_h^{(k)} + CH^4,$$

$$\|u_h^{(k)} - u_{H,\ell}^{\text{ms},h,(k)}\| \leq CH^2,$$

with  $C$  independent of  $A'$  and the regularity of the eigenfunctions and  $(\lambda_h, u_h)$  is the reference solution.

M. & Peterseim, Computation of eigenvalues by nume. upscaling, 2015.

# Numerical example: eigenvalues



$k$	$\lambda_h^{(k)}$	$e^{(k)}(1/2\sqrt{2})$	$e^{(k)}(1/4\sqrt{2})$	$e^{(k)}(1/8\sqrt{2})$	$e^{(k)}(1/16\sqrt{2})$
1	21.4144522	5.472755371	0.237181706	0.010328293	0.000781683
2	40.9134676	-	0.649080539	0.032761482	0.002447049
3	44.1561133	-	1.687388874	0.097540102	0.004131422
4	60.8278691	-	1.648439518	0.028076168	0.002079812
5	65.6962136	-	2.071005692	0.247424446	0.006569640
6	70.1273082	-	4.265936007	0.232458016	0.016551520
7	82.2960238	-	3.632888104	0.355050163	0.013987920
8	92.8677605	-	6.850048057	0.377881216	0.049841235
9	99.6061234	-	10.305084010	0.469770376	0.026027378
10	109.1543283	-	-	0.476741452	0.005606426
11	129.3741945	-	-	0.505888044	0.062382302
12	138.2164330	-	-	0.554736550	0.039487317
13	141.5464639	-	-	0.540480876	0.043935515
14	145.7469718	-	-	0.765411709	0.034249528
15	152.6283573	-	-	0.712383825	0.024716759
16	155.2965039	-	-	0.761104705	0.026228034
17	158.2610708	-	-	0.749058367	0.091826207
18	164.1452194	-	-	0.840736127	0.118353184
19	171.1756923	-	-	0.946719951	0.111314058
20	179.3917590	-	-	0.928617606	0.119627862

Table: Errors  $e^{(k)}(H) =: \frac{\lambda_H^{\text{ms},(k)} - \lambda_h^{(k)}}{\lambda_h^{(k)}}$  and  $h = 2^{-7} \sqrt{2}$ .

# Parabolic equations

The parabolic problem: Find  $u \in V$  such that

$$(\dot{u}, v) + (A \nabla u, \nabla v) = (f(t), v), \quad \forall v \in V, \quad t > 0$$

and  $u(0) = u_0 \in L^2(\Omega)$ . We assume  $A$  to be independent of  $t$ .

FE Backward Euler: Find  $u_h^n \in V_h$  such that

$$(\bar{\partial}_t u_h^n, v) + a(u_h^n, v) = (f^n, v), \quad \forall v \in V_h,$$

and  $u_h^0 \in V_h$  some approximation of  $u_0$ .

LOD: Find  $(u_H^{\text{ms}})^n \in V_{H,\ell}^{\text{ms},h}$  such that

$$(\bar{\partial}_t (u_H^{\text{ms}})^n, v) + a((u_H^{\text{ms}})^n, v) = (f^n, v), \quad \forall v \in V_{H,\ell}^{\text{ms},h},$$

and  $(u_H^{\text{ms}})^0 \in V_{H,\ell}^{\text{ms},h}$  some approximation of  $u_0$ .

# Parabolic equations



## Theorem

$$\|u_h^n - (u_H^{\text{ms}})^n\|_{L^2(\Omega)} \leq C(1 + \log(\frac{t_n}{\tau}))H^2(t_n^{-1}\|u_h^0\|_{L^2(\Omega)} + \|f\|_{W^{1,\infty}(L^2(\Omega))})$$

with  $C$  independent of  $A'$ .

- The analysis uses classic a priori error estimation techniques and the elliptic results.
- The term  $t_n^{-1}$  appears also in  $u - u_h$  bounds if  $u_0 \in L^2(\Omega)$ . The log term can be avoided if  $f(t) \in H_0^1(\Omega)$ .
- The case  $f = f(u)$  can also be treated, under certain growth conditions on  $f'(u)$  and  $f''(u)$ .
- The case  $A = A(t)$  or  $A = A(u)$  is not covered and would require updates of  $V_{H,\ell}^{\text{ms},h}$ .


# Numerical experiment: The heat equation



$$H = 2^{-1}, 2^{-2}, \dots, 2^{-6}$$

$$h = 2^{-7}, \ell = \log(1/H), u_0 = 1,$$

$$T = 1, \tau = 0.001$$



$\|u_h^n - (u_H^{ms})^n\|$  vs. #dof

M.& Persson, *Multiscale techniques for parabolic problems*, Numer. Math. 2018.

# More applications

## Stationary/eigenvalue problems

- Semilinear, (Henning, M., Peterseim), 2014.
- Gross-Pitaevskii, (Henning, M., Peterseim), 2014.
- Helmholtz, (Gallistl & Peterseim), 2015.
- Reduced basis, (Abdulle & Henning), 2015.
- Elasticity, (Henning & Persson), 2016.
- High contrast, (Peterseim & Scheichl), 2016.
- Iterative solvers, (Kornhuber & Yserentant), 2017.
- Network models, (Ketttil et. al.), 2019.

## Time-dependent problems

- Thermoelasticity, (M. & Persson), 2017.
- Wave equation, (Abdulle & Henning), 2017.
- Two phase flow, (Hellman & M.), 2019

# Outline

- ① Elliptic problem: Homogenization and FEM
- ② Introduction to LOD
- ③ High contrast data
- ④ Applications
- ⑤ **Conclusions**

# Conclusion and outlook

- By LOD we compute an effective stiffness matrix on a coarse scale. (numerical homogenization)
- Rapidly varying diffusion of low contrast is well understood.
- New ideas for high contrast problems (which is a great challenge for any method).
- Great reduction in computational cost when the basis is reused (load cases, eigenvalues, time dependent, non-linear, control).
- Future challenges: random diffusion, interfaces, multiscale in time, and fully non-linear problems.

Thank you for your attention!