

Computation of eigenvalues using multiscale techniques

Axel Målqvist¹

Daniel Peterseim²


2014-09-10

¹Chalmers University of Technology and University of Gothenburg


²Rheinische Friedrich-Wilhelms Universität Bonn

Multiscale problems

Applications such as



▷ composite materials



▷ flow in a porous medium

require numerical solution of partial differential equations with rough data (module of elasticity, conductivity, or permeability).

Major challenge: Features on **multiple non-separated scales**.

Multiscale methods

Let A be rapidly varying data and consider a differential equation and its corresponding numerical approximation,

$$\mathcal{L}(A)u = f \qquad \mathcal{L}_h(A)u_h = f_h.$$

Classical finite element methods typically give

$$\|u - u_h\| \leq C(A, A')h^\gamma.$$

Multiscale methods seek an upscaled representation

$$\mathcal{L}_H(A)u_H = f_H$$

fulfilling $\|u_h - u_H\| \leq C(A)H^\gamma$ with C independent of A' .

How well is the spectrum of $\mathcal{L}$ preserved?


- 1 **Model problem**
- 2 Upscaling technique and error analysis
- 3 Numerical experiments
- 4 Applications to non-linear eigenvalue problems
- 5 Conclusions

Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem

$$-\nabla \cdot \mathbf{A} \nabla u = \lambda u \quad \text{in } \Omega \quad u = 0 \quad \text{on } \partial\Omega$$

with data $0 < \alpha \leq \mathbf{A} \leq \beta < \infty$




Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem (variational form): find $u \in V := H_0^1(\Omega)$ and $\lambda \in \mathbb{R}$ such that

$$a(u, v) := \int_{\Omega} (\mathbf{A} \nabla u) \cdot \nabla v \, dx = \lambda \int_{\Omega} u \cdot v \, dx \quad \text{for all } v \in V$$

with data $0 < \alpha \leq A \leq \beta < \infty$




Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem (FE approximation):
 $u_h \in V_h \subset V$ and $\lambda_h \in \mathbb{R}$ such that

$$a(u_h, v) := \int_{\Omega} (\mathbf{A} \nabla u_h) \cdot \nabla v \, dx = \lambda_h \int_{\Omega} u_h \cdot v \, dx \quad \text{for all } v \in V_h$$

with data $0 < \alpha \leq A \leq \beta < \infty$



Model multiscale eigenvalue problem

Prototypical self-adjoint eigenvalue problem (FE approximation):
 $u_h \in V_h \subset V$ and $\lambda_h \in \mathbb{R}$ such that

$$a(u_h, v) := \int_{\Omega} (\mathbf{A} \nabla u_h) \cdot \nabla v \, dx = \lambda_h \int_{\Omega} u_h \cdot v \, dx \quad \text{for all } v \in V_h$$

with data $0 < \alpha \leq A \leq \beta < \infty$

Numerical error (piecewise linear continuous FE approximation)

- For an eigenpair $(u^{(k)}, \lambda^{(k)})$ with $u^{(k)} \in H^2(\Omega)$ it holds

$$\lambda^{(k)} \leq \lambda_h^{(k)} \leq \lambda^{(k)} + C(A, A', k) h^2,$$
$$\|u^{(k)} - u_h^{(k)}\| := \|A^{1/2} \nabla(u^{(k)} - u_h^{(k)})\|_{L^2(\Omega)} \leq C(A, A', k) h.$$

- The mesh size h has to resolve the variations in A , e.g. $h < \epsilon$ if A is periodic.

Objectives

Investigate how the Localized Orth. Decomposition (LOD) in



A. Målqvist and D. Peterseim.

Localization of Elliptic Multiscale Problems.

Mathematics of Computation 2014

preserves the (low part of the) spectrum of $-\nabla \cdot A \nabla$.

Without assumptions on scales (A') or regularity (u):

$$\lambda_h \leq \lambda_H^{\text{ms}} \leq \lambda_h + CH^4,$$

$$|||u_h - u_H^{\text{ms}}||| \leq CH^2.$$



A. Målqvist and D. Peterseim.


Computation of eigenvalues by numerical upscaling.

accepted for publication in Numerische Mathematik

- 1 Model problem
- 2 **Upscaling technique and error analysis**
- 3 Numerical experiments
- 4 Applications to non-linear eigenvalue problems
- 5 Conclusions

Multiscale decomposition


- (coarse) FE mesh $\mathcal{T}$ with parameter H
- P1-FE space $V_H := \{v \in V \mid \forall T \in \mathcal{T}, v|_T \in P_1(T)\}$
- $\mathfrak{I}_{\mathcal{T}} : V \rightarrow V_H$ a Clément interpolation operator



Decomposition

$$V = V_H \oplus V^f \quad \text{with } V^f := \text{kernel } \mathfrak{I}_{\mathcal{T}} = \{v \in V \mid \mathfrak{I}_{\mathcal{T}} v = 0\}$$

Example:



rough coefficient


Multiscale decomposition

- (coarse) FE mesh $\mathcal{T}$ with parameter H
- P1-FE space $V_H := \{v \in V \mid \forall T \in \mathcal{T}, v|_T \in P_1(T)\}$
- $\mathfrak{I}_{\mathcal{T}} : V \rightarrow V_H$ a Clément interpolation operator

Decomposition

$$V = V_H \oplus V^f \quad \text{with } V^f := \text{kernel } \mathfrak{I}_{\mathcal{T}} = \{v \in V \mid \mathfrak{I}_{\mathcal{T}} v = 0\}$$

Example:



Orthogonalization


- For each $v \in V_H$ define finescale projection $\mathfrak{F}v \in V^f$ by

$$a(\mathfrak{F}v, w) = a(v, w) \quad \text{for all } w \in V^f$$

Orthogonal Decomposition

$$V = V_H^{\text{ms}} \oplus V^f \quad \text{with } V_H^{\text{ms}} := (V_H - \mathfrak{F}V_H)$$

Example:



Ideal multiscale representation

Given the space V_H^{ms} we construct a Galerkin approximation:

Ideal method

Find $u_H^{\text{ms}} \in V_H^{\text{ms}}$, $\lambda_H^{\text{ms}} \in \mathbb{R}$ such that

$$a(u_H^{\text{ms}}, v) = \lambda_H^{\text{ms}}(u_H^{\text{ms}}, v), \quad \forall v \in V_H^{\text{ms}}.$$

- We note that $\dim(V_H^{\text{ms}}) = \dim(V_H)$.
- For V_H^{ms} to be useful we need a discrete localized basis.
- But first of all we need to show that λ_H^{ms} is a good approximation of λ .

A priori error bound (ideal case)

For the k :th eigenvalue it holds

Theorem

$$\lambda^{(k)} \leq \lambda_H^{ms,(k)} \leq \lambda^{(k)} + CH^4,$$

C independent of A' and only H^1 -regularity of the eigenfunctions.

Sketch of proof for the *lowest* eigenvalue:

- Let $u^{(1)} := u = u_c + u_f$ with $u_c \in V_H^{ms}$ and $u_f \in V^f$, such that $\|u\|_{L^2(\Omega)} = 1$. Then

$$\begin{aligned} \lambda_H^{ms,(1)} &\leq \frac{a(u_c, u_c)}{(u_c, u_c)} \leq \frac{a(u, u)}{(u_c, u_c)} = \frac{a(u, u)}{(u - u_f, u - u_f)} \\ &= \frac{\lambda^{(1)}}{(u, u) - 2(u, u_f) + (u_f, u_f)} \leq \frac{\lambda^{(1)}}{1 - 2(u, u_f)}. \end{aligned}$$

A priori error bound (ideal case)

For the k :th eigenvalue it holds

Theorem

$$\lambda^{(k)} \leq \lambda_H^{ms,(k)} \leq \lambda^{(k)} + CH^4,$$

C independent of A' and only H^1 -regularity of the eigenfunctions.

Sketch of proof for the *lowest* eigenvalue:

- Since $\mathfrak{I}_{\mathcal{T}} u_f = 0$, $(\mathfrak{I}_{\mathcal{T}} u, u_f) = 0$ (weighted Clement, CV99), $a(u_c, u_f) = 0$, and $|||u|||^2 = \lambda^{(1)}$, we have

$$\begin{aligned} (u, u_f) &= (u - \mathfrak{I}_{\mathcal{T}} u, u_f - \mathfrak{I}_{\mathcal{T}} u_f) \leq CH^2 |||u||| \cdot |||u_f||| \leq C' H^2 |||u_f|||, \\ |||u_f|||^2 &= a(u, u_f) = \lambda^{(1)} (u - \mathfrak{I}_{\mathcal{T}} u, u_f - \mathfrak{I}_{\mathcal{T}} u_f) \leq CH^2 |||u_f|||. \end{aligned}$$

- We conclude $\lambda_H^{ms,(1)} \leq \frac{\lambda^{(1)}}{1-CH^4} \leq \lambda^{(1)} + 2CH^4$.

A priori error bound (ideal case)

For the k :th eigenfunction it holds

Theorem

$$|||u^{(k)} - u_H^{ms,(k)}||| \leq CH^2,$$

C independent of A' and only H^1 -regularity of the eigenfunctions.

- Similar arguments using $\Im_{\mathcal{T}} u_f = 0$ and $(\Im_{\mathcal{T}} u, u_f) = 0$.
- Only $H^1(\Omega)$ regularity is assumed.

Can we find a localized discrete basis that approximates V_H^{ms} ?


Modified nodal basis

- $\mathcal{N}$ denotes set of interior vertices of $\mathcal{T}$
- $\phi_x \in V_H$ denotes classical nodal basis function ($x \in \mathcal{N}$)
- $\mathfrak{F}\phi_x \in V^f$ denotes finescale correction of ϕ_x ($x \in \mathcal{N}$)

Ideal multiscale FE space

$$V_H^{\text{ms}} = \text{span} \{ \phi_x - \mathfrak{F}\phi_x \mid x \in \mathcal{N} \}$$

Example




Modified nodal basis

- $\mathcal{N}$ denotes set of interior vertices of $\mathcal{T}$
- $\phi_x \in V_H$ denotes classical nodal basis function ($x \in \mathcal{N}$)
- $\mathfrak{F}\phi_x \in V^f$ denotes finescale correction of ϕ_x ($x \in \mathcal{N}$)


Ideal multiscale FE space

$$V_H^{\text{ms}} = \text{span} \{ \phi_x - \mathfrak{F}\phi_x \mid x \in \mathcal{N} \}$$

Example




Modified nodal basis



Localization

- Define nodal patches of ℓ -th order $\omega_{x,\ell}$ about $x \in \mathcal{N}$



- Localized corrections $\mathfrak{F}\phi_{x,\ell} \in V^f(\omega_{x,\ell}) := \{v \in V^f \mid v|_{\Omega \setminus \omega_{x,\ell}} = 0\}$ solve


$$a(\mathfrak{F}\phi_{x,\ell}, w) = a(\phi_x, w) \quad \text{for all } w \in V^f(\omega_{x,\ell})$$

Localized multiscale FE spaces

$$V_{H,\ell}^{\text{ms}} = \text{span}\{\phi_x - \mathfrak{F}\phi_{x,\ell} \mid x \in \mathcal{N}\}$$


Fine scale discretization

- Finescale mesh



$\mathcal{T}$

mesh refinement



$\mathcal{T}_h$ with $h \leq H$

- Reference FE space

$$V_h := \{v \in V \mid \forall T \in \mathcal{T}(\Omega), v|_T \in P_1(T)\}$$

- Reference FE solution $u_h \in V_h$ and $\lambda_h \in \mathbb{R}$ solves

$$a(u_h, v) = \lambda_h(u_h, v) \quad \text{for all } v \in V_h$$

- Fully discrete corrections $\mathfrak{F}\phi_{x,\ell}^h \in V_h^f(\omega_{x,\ell}) := V^f(\omega_{x,\ell}) \cap V_h$:

$$a(\mathfrak{F}\phi_{x,\ell}^h, w) = a(\phi_x, w) \quad \text{for all } w \in V_h^f(\omega_{x,\ell})$$

Localized Orthogonal Decomposition (LOD)

Fully discrete multiscale FE spaces

$$V_{H,\ell}^{\text{ms},h} = \text{span}\{\phi_x - \mathfrak{F}\phi_{x,\ell}^h \mid x \in \mathcal{N}\}$$

Fully discrete multiscale approximation $u_{H,\ell}^{\text{ms},h} \in V_{H,\ell}^{\text{ms},h}$, $\lambda_{H,\ell}^{\text{ms},h} \in \mathbb{R}$

$$a(u_{H,\ell}^{\text{ms},h}, v) = \lambda_{H,\ell}^{\text{ms},h}(u_{H,\ell}^{\text{ms},h}, v) \quad \text{for all } v \in V_{H,\ell}^{\text{ms},h}$$

Remarks:

- $\dim V_{H,\ell}^{\text{ms},h} = |\mathcal{N}| = \dim V_H$
- The basis functions have local support, with overlap depending on ℓ , and are independent.

A priori error analysis (discrete case)

Lemma (Truncation error)

$$||| \mathfrak{F}\phi_x^h - \mathfrak{F}\phi_{x,\ell}^h ||| \leq C_1 \gamma^\ell ||| \mathfrak{F}\phi_x^h |||.$$

$C_1 < \infty$ and $\gamma < 1$ depends on β/α , not A' .

By choosing $\ell = C_2 \log(H^{-1})$ with appropriate C_2 we guarantee that the truncation leads to a higher order perturbation:

Theorem

$$\begin{aligned} \lambda_h^{(k)} &\leq \lambda_{H,\ell}^{ms,h,(k)} \leq \lambda_h^{(k)} + CH^4, \\ ||| u_h^{(k)} - u_{H,\ell}^{ms,h,(k)} ||| &\leq CH^2, \end{aligned}$$

with C independent of A' and the regularity of the eigenfunctions.

A priori error analysis (discrete case)

The result can be improved using a postprocessing technique:



J. Xu and A. Zhou.

A two-grid discretization scheme for eigenvalue problems.
Mathematics of Computation 2001.

Find $u_h^p \in V_h$ s.t.

$$a(u_h^p, v) = \lambda_{H,\ell}^{\text{ms},h}(u_{H,\ell}^{\text{ms},h}, v), \quad v \in V_h,$$

and letting $\lambda_h^p = a(u_h^p, u_h^p)/(u_h^p, u_h^p)$.

Theorem

$$\begin{aligned} \lambda_h^{(k)} &\leq \lambda_h^{p,(k)} \leq \lambda_h^{(k)} + CH^6, \\ ||| u_h^{(k)} - u_h^{p,(k)} ||| &\leq CH^4. \end{aligned}$$

Outline

- 1 Model problem
- 2 Upscaling technique and error analysis
- 3 **Numerical experiments**
- 4 Applications to a non-linear eigenvalue problems
- 5 Conclusions


Eigenvalue Problem



k	$\lambda_h^{(k)}$	$e^{(k)}(1/2 \sqrt{2})$	$e^{(k)}(1/4 \sqrt{2})$	$e^{(k)}(1/8 \sqrt{2})$	$e^{(k)}(1/16 \sqrt{2})$
1	9.6436869	0.003494567	0.000034466	0.000000546	0.000000010
2	15.1989274	0.009621397	0.000079887	0.000000845	0.000000010
3	19.7421815	0.023813222	0.000213097	0.000002073	0.000000023
4	29.5281571	0.096910157	0.000724615	0.000006574	0.000000076
5	31.9265496	0.094454625	0.000874659	0.000009627	0.000000138
6	41.4922250	-	0.002395227	0.000019934	0.000000254
7	44.9604884	-	0.002443271	0.000019683	0.000000223
8	49.3631826	-	0.003651870	0.000028869	0.000000308
9	49.3655623	-	0.004266472	0.000032835	0.000000355
10	56.7389993	-	0.006863742	0.000055219	0.000000618
11	65.4085991	-	0.011534878	0.000082414	0.000000856
12	71.0947630	-	0.012596114	0.000090083	0.000001002
13	71.6064671	-	0.014249938	0.000098736	0.000001006
14	79.0043994	-	0.021801461	0.000164436	0.000001605
15	89.3706421	-	0.033550079	0.000211985	0.000002296
16	92.3648207	-	0.040060692	0.000239441	0.000002295
17	97.4459210	-	0.037438984	0.000284936	0.000002724
18	98.7545147	-	0.044544409	0.000269854	0.000002559
19	98.7545639	-	0.047835987	0.000276139	0.000002539
20	101.6755971	-	0.038203654	0.000297356	0.000002909

Table : Errors $e^{(k)}(H) =: \frac{\lambda_H^{\text{ms},(k)} - \lambda_h^{(k)}}{\lambda_h^{(k)}}$ and $h = 2^{-7} \sqrt{2}$.

Eigenvalue Problem




k	$\lambda_h^{(k)}$	$e^{(k)}(1/2 \sqrt{2})$	$e^{(k)}(1/4 \sqrt{2})$	$e^{(k)}(1/8 \sqrt{2})$	$e^{(k)}(1/16 \sqrt{2})$
1	21.4144522	5.472755371	0.237181706	0.010328293	0.000781683
2	40.9134676	-	0.649080539	0.032761482	0.002447049
3	44.1561133	-	1.687388874	0.097540102	0.004131422
4	60.8278691	-	1.648439518	0.028076168	0.002079812
5	65.6962136	-	2.071005692	0.247424446	0.006569640
6	70.1273082	-	4.265936007	0.232458016	0.016551520
7	82.2960238	-	3.632888104	0.355050163	0.013987920
8	92.8677605	-	6.850048057	0.377881216	0.049841235
9	99.6061234	-	10.305084010	0.469770376	0.026027378
10	109.1543283	-	-	0.476741452	0.005606426
11	129.3741945	-	-	0.505888044	0.062382302
12	138.2164330	-	-	0.554736550	0.039487317
13	141.5464639	-	-	0.540480876	0.043935515
14	145.7469718	-	-	0.765411709	0.034249528
15	152.6283573	-	-	0.712383825	0.024716759
16	155.2965039	-	-	0.761104705	0.026228034
17	158.2610708	-	-	0.749058367	0.091826207
18	164.1452194	-	-	0.840736127	0.118353184
19	171.1756923	-	-	0.946719951	0.111314058
20	179.3917590	-	-	0.928617606	0.119627862

Table : Errors $e^{(k)}(H) =: \frac{\lambda_H^{\text{ms},(k)} - \lambda_h^{(k)}}{\lambda_h^{(k)}}$ and $h = 2^{-7} \sqrt{2}$.

Eigenvalue problem, non-nested meshes

In many applications $V_H \not\subset V_h$, e.g., a domain with inclusions,




where A varies between 100 (gray) and 1 (white).

Using the nodal interpolant $I_h : C(\Omega) \rightarrow V_h$ we can construct a space $\tilde{V}_H = \{I_h \phi_x, x \in \mathcal{N}\} \subset V_h$ with $V_H = \text{span}(\phi_x) \not\subset V_h$.

Then $\mathfrak{I}_{\mathcal{T}} : V \rightarrow \tilde{V}_H$ is used to get V^f and V_H^{ms} .


Eigenvalue Problem, non-nested meshes



ℓ	$\lambda_h^{(\ell)}$	$e^{(\ell)}(1/2\sqrt{2})$	$e^{(\ell)}(1/4\sqrt{2})$	$e^{(\ell)}(1/8\sqrt{2})$	$e^{(\ell)}(1/16\sqrt{2})$
1	25.6109462	0.025518831	0.000572341	0.000017083	0.000000700
2	58.9623566	-	0.005235813	0.000090490	0.000002710
3	67.5344854	-	0.006997582	0.000154850	0.000006488
4	98.2808694	-	0.023497502	0.000358178	0.000011675
5	121.2290664	-	0.052366141	0.000563438	0.000016994
6	125.2014779	-	0.066627585	0.000747688	0.000019934
7	156.0597873	-	0.145676350	0.001579177	0.000034329
8	168.2376096	-	0.095360287	0.001320185	0.000043781
9	197.4467434	-	0.343991317	0.002888471	0.000049479
10	209.4657306	-	-	0.003223901	0.000056318
11	222.4472476	-	-	0.003431462	0.000080284
12	245.5656759	-	-	0.005906282	0.000102243
13	253.7074603	-	-	0.006215809	0.000121646
14	288.0756442	-	-	0.013859535	0.000180899
15	298.8903269	-	-	0.010587124	0.000138404
16	311.4410556	-	-	0.012159268	0.000161510
17	324.6865434	-	-	0.012143676	0.000176624
18	336.7931865	-	-	0.016554437	0.000233067
19	379.5697606	-	-	0.023254268	0.000325324
20	386.9938901	-	-	0.028772395	0.000383532

Table : Errors $e^{(k)}(H) =: \frac{\lambda_H^{\text{ms},(k)} - \lambda_h^{(k)}}{\lambda_h^{(k)}}$ and $h \approx 2^{-9} - 2^{-7}$.

Eigenvalue Problem, non-nested, postpr.



ℓ	$\lambda_h^{(\ell)}$	$e^{(\ell)}(1/2\sqrt{2})$	$e^{(\ell)}(1/4\sqrt{2})$	$e^{(\ell)}(1/8\sqrt{2})$	$e^{(\ell)}(1/16\sqrt{2})$
1	25.6109462	0.001559704	0.000003765	0.000000008	3.5e-10
2	58.9623566	-	0.000191532	0.000000213	1.9e-08
3	67.5344854	-	0.000284980	0.000000474	0.000000001
4	98.2808694	-	0.002239689	0.000002253	0.000000004
5	121.2290664	-	0.007461217	0.000005065	0.000000008
6	125.2014779	-	0.011284614	0.000006826	0.000000008
7	156.0597873	-	0.042466017	0.000023867	0.000000024
8	168.2376096	-	0.025093182	0.000027547	0.000000042
9	197.4467434	-	0.186960343	0.000072471	0.000000051
10	209.4657306	-	-	0.000105777	0.000000079
11	222.4472476	-	-	0.000131569	0.000000129
12	245.5656759	-	-	0.000286351	0.000000213
13	253.7074603	-	-	0.000268463	0.000000255
14	288.0756442	-	-	0.000915102	0.000000473
15	298.8903269	-	-	0.000762135	0.000000403
16	311.4410556	-	-	0.000873769	0.000000504
17	324.6865434	-	-	0.000955392	0.000000642
18	336.7931865	-	-	0.001335246	0.000000977
19	379.5697606	-	-	0.002896202	0.000001886
20	386.9938901	-	-	0.007202657	0.000001908

Table : Errors $e^{(k)}(H) =: \frac{\lambda_H^{\text{ms},(k)} - \lambda_h^{(k)}}{\lambda_h^{(k)}}$ and $h \approx 2^{-9} - 2^{-7}$.

- 1 Model problem
- 2 Upscaling technique and error analysis
- 3 Numerical experiments
- 4 **Applications to non-linear eigenvalue problems**
- 5 Conclusions

The quadratic eigenvalue problem

Consider the quadratic eigenvalue problem (QEP): find $u \in V$, $\|u\|_{L^2(\Omega)} = 1$, and $\lambda \in \mathbb{C}$ such that

$$(A \nabla u, \nabla v) + \lambda c(u, v) + \lambda^2(u, v) = 0, \quad \forall v \in V.$$

This equation appears in structural mechanics and describes damped vibrations. Discretization gives an algebraic QEP,

$$Kx + \lambda Cx + \lambda^2 Mx = 0,$$

where K is stiffness, C is damping, and M is mass matrix.

If C is symmetric, real, and positive: the eigenvalues are complex conjugate with negative real part.



F. Tisseur and K. Meerbergen.

The quadratic eigenvalue problem SIAM Review 2001.

The quadratic eigenvalue problem

Consider the quadratic eigenvalue problem (QEP): find $u \in V$, $\|u\|_{L^2(\Omega)} = 1$, and $\lambda \in \mathbb{C}$ such that

$$(A \nabla u, \nabla v) + \lambda c(u, v) + \lambda^2(u, v) = 0, \quad \forall v \in V.$$

This equation appears in structural mechanics and describes damped vibrations. Discretization gives an algebraic QEP,

$$Kx + \lambda Cx + \lambda^2 Mx = 0,$$

where K is stiffness, C is damping, and M is mass matrix.

- Rayleigh damping $C = \alpha_0 M + \alpha_1 K$ leads to unchanged eigenmodes.
- Systems of Rayleigh damped components, α_0, α_1 are functions.

QEP: Linearization and approximation

Linearization: ($y = \lambda x$)

$$\begin{bmatrix} K & 0 \\ 0 & M \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \lambda \begin{bmatrix} -C & -M \\ M & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

On weak form: find (u_1, u_2) and $\lambda \in \mathbb{C}$ such that,

$$a(u, v) = \lambda b(u, v),$$

$$a(u, v) = (A \nabla u_1, \nabla v_1) + (u_2, v_2),$$



$$b(u, v) = -c(u_1, v_1) - (u_2, v_1) + (u_1, v_2).$$

Multiscale formulation:

Find $u_H^{\text{ms}} \in V_H^{\text{ms}} \times V_H^{\text{ms}}$ and $\lambda_H^{\text{ms}} \in \mathbb{C}$ such that,


$$a(u_H^{\text{ms}}, v) = \lambda_H^{\text{ms}} b(u_H^{\text{ms}}, v), \quad \forall v \in V_H^{\text{ms}} \times V_H^{\text{ms}}.$$

Numerical experiment



$$H = 2^{-1}, 2^{-2}, \dots, 2^{-5 \cdot 10^1}$$



$$h = 2^{-6}, k = \infty$$




$$|\lambda_h - \lambda_H^{\text{ms}}| / |\lambda_h| \text{ vs. } \# \text{dof}$$

$$A = 1, c(u, v) = \int_{\Omega} (1 + \sin(10x)) u v \, dx$$

Numerical experiment





$H = 2^{-1}, 2^{-2}, \dots, 2^{-5 \cdot 10^1}$
 $h = 2^{-6}, k = \infty$




A (pic), $c(u, v) = \int_{\Omega} (1 + \sin(10x)) u v \, dx$

Numerical experiment



$$H = 2^{-1}, 2^{-2}, \dots, 2^{-5} \quad |\lambda_h - \lambda_H^{\text{ms}}| / |\lambda_h| \text{ vs. } \# \text{dof}$$

$$h = 2^{-6}, k = \infty$$



$$\pi c(u, v) = \int_{\Omega} \left(\text{atan}(10x - 5) + \frac{\pi}{2} \right) A \nabla u \nabla v \, dx$$

QEP: Analysis

We note that the operator $B : V \times V \rightarrow V \times V$ defined by,

$$a(Bu, v) = b(u, v),$$

has eigenvalues $Bu = \mu u = \lambda^{-1} u$ since,

$$b(u, v) = a(Bu, v) = a(\mu u, v) = \mu a(u, v).$$

Furthermore, a is coercive, bounded, symmetric and b is bounded if $|c(u, v)| \leq C\|u\|_V\|v\|_V$.

 J. Descloux, N. Nassif, and J. Rappaz.

On spectral approximation I & II. RAIRO 1978. (non compact)

QEP: Analysis

We note that the operator $B : V \times V \rightarrow V \times V$ defined by,

$$a(Bu, v) = b(u, v),$$

has eigenvalues $Bu = \mu u = \lambda^{-1} u$ since,

$$b(u, v) = a(Bu, v) = a(\mu u, v) = \mu a(u, v).$$

Conjecture

Let $c(u, v) = \int_{\Omega} \alpha_0 A \nabla u \cdot \nabla v \, dx + \int_{\Omega} \alpha_1 uv \, dx$, with $\alpha_i \in W^{1,\infty}(\Omega)$ and let λ be an eigenvalue with algebraic multiplicity m . Then,

$$\left| \lambda - \left(\frac{1}{m} \sum_{j=1}^m (\lambda_{H,j}^{ms})^{-1} \right)^{-1} \right| \leq CH^2.$$

The Gross-Pitaevskii equation

Consider the Gross-Pitaevskii equation: find $u \in V$, $\|u\|_{L^2(\Omega)} = 1$, and $\lambda \in \mathbb{R}$ such that

$$(A\nabla u, \nabla v) + (bu, v) + (u^3, v) = \lambda(u, v), \quad \forall v \in V.$$

The equation describes the quantum states of a boson gas cooled down to an ultra-low temperature.

- We reuse the same discrete space $V_{H,\ell}^{\text{ms},h}$ i.e. we ignore the low order non-linearity on the fine scale.
- We then solve the upscaled non-linear eigenvalue problem on the coarse scale.

 P. Henning, A. Målqvist, and D. Peterseim.

Two-level discretization techniques for **ground state** computations of Bose-Einstein condensates.
SIAM Journal on Numerical Analysis 2014.

The Gross-Pitaevskii equation

Consider the Gross-Pitaevskii equation: find $u \in V$, $\|u\|_{L^2(\Omega)} = 1$, and $\lambda \in \mathbb{R}$ such that

$$(A\nabla u, \nabla v) + (bu, v) + (u^3, v) = \lambda(u, v), \quad \forall v \in V.$$

The equation describes the quantum states of a boson gas cooled down to an ultra-low temperature.

Theorem

$$\begin{aligned} \lambda &\leq \lambda_h^p \leq \lambda + CH^2\|u - u_h\|_{H^1(\Omega)} + CH^4, \\ \|u - u_h^p\|_{H^1(\Omega)} &\leq C\|u - u_h\|_{H^1(\Omega)} + CH^3. \end{aligned}$$

for the ground state, with C independent on the regularity of u and variations in A .

- 1 Model problem
- 2 Upscaling technique and error analysis
- 3 Numerical experiments
- 4 Applications to non-linear eigenvalue problems
- 5 **Conclusions**

Conclusion

- The Localized Orthogonal Decomposition (LOD) technique preserves the low spectrum of the operator. In particular the eigenvalue error is proportional to H^4 after postprocessing H^6 .
- Numerical experiments indicates even higher rates possibly due to additional regularity in the solution that is not taken advantage of in the analysis.
- The technique is applicable also for non-linear eigenvalue problems without pre-asymptotic effects in the convergence.
- More work is needed in the analysis for the quadratic eigenvalue problem.
- Numerical tests with more complicated damping is needed.

Thank you for your attention!